



Prior Authorization

Kentucky Medicaid



MedImpact Clinical Document

Kentucky Medicaid Prior Authorization Criteria

VERSION: DECEMBER 2025




Table of Contents

Table of Contents	ii
INTRODUCTION	1
CARDIOVASCULAR: ANGIOTENSIN RECEPTOR MODULATORS	2
CARDIOVASCULAR: ANTI-ANGINAL & ANTI-ISCHEMIC AGENTS	4
CARDIOVASCULAR: ANTIARRHYTHMICS (ORAL ANTI-ARRHYTHMICS)	5
CARDIOVASCULAR: BETA BLOCKERS	6
CARDIOVASCULAR: CALCIUM CHANNEL BLOCKERS	7
CARDIOVASCULAR: ANTICOAGULANTS	9
CARDIOVASCULAR: PLATELET AGGREGATION INHIBITORS	10
CARDIOVASCULAR: PULMONARY ARTERIAL HYPERTENSION (PAH) AGENTS, ORAL AND INHALED ...	11
CARDIOVASCULAR: LIPOTROPICS	13
GASTROINTESTINAL: ANTIEMETICS AND ANTIVERTIGO AGENTS	17
GASTROINTESTINAL: ANTIDIARRHEALS	20
GASTROINTESTINAL: ANTISPASMODICS/ANTICHOLINERGICS	21
GASTROINTESTINAL: ANTI-ULCER PROTECTANTS	23
GASTROINTESTINAL: BILE SALTS	24
GASTROINTESTINAL: HELICOBACTER PYLORI (H. PYLORI) TREATMENT	28
GASTROINTESTINAL: HISTAMINE II (H2) RECEPTOR BLOCKERS	30
GASTROINTESTINAL: LAXATIVES AND CATHARTICS	31
GASTROINTESTINAL: GASTROINTESTINAL MOTILITY AGENTS	32
GASTROINTESTINAL: PROTON PUMP INHIBITORS	35
GASTROINTESTINAL: ULCERATIVE COLITIS AGENTS	38
RESPIRATORY: ANTIBIOTICS, INHALED	39
RESPIRATORY: ANTIHISTAMINES, MINIMALLY SEDATING	41
RESPIRATORY: INTRANASAL RHINITIS AGENTS	42
RESPIRATORY: LEUKOTRIENE MODIFIERS	43
RESPIRATORY: BRONCHODILATORS, BETA-AGONIST	44
RESPIRATORY: GLUCOCORTICOIDS, INHALED	46
RESPIRATORY: EPINEPHRINE, SELF-INJECTABLE	47
RESPIRATORY: CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD) AGENTS	48
CENTRAL NERVOUS SYSTEM: ALZHEIMER'S AGENTS	50
CENTRAL NERVOUS SYSTEM: ANXIOLYTICS	51
CENTRAL NERVOUS SYSTEM: ANTICONVULSANTS	53
CENTRAL NERVOUS SYSTEM: ANTIPSYCHOTICS: FIRST GENERATION (TYPICAL)	58
CENTRAL NERVOUS SYSTEM: ANTIPSYCHOTICS: SECOND GENERATION (ATYPICAL) AND INJECTABLE	59
CENTRAL NERVOUS SYSTEM: DOPAMINE RECEPTOR AGONISTS	67
CENTRAL NERVOUS SYSTEM: PARKINSON'S DISEASE (ANTIPARKINSON'S AGENTS)	68
CENTRAL NERVOUS SYSTEM: MOVEMENT DISORDERS	71

CENTRAL NERVOUS SYSTEM: ANTIDEPRESSANTS	74
CENTRAL NERVOUS SYSTEM: ANTI-MIGRAINE AGENTS, TRIPTANS	79
CENTRAL NERVOUS SYSTEM: ANTI-MIGRAINE AGENTS, CGRP INHIBITORS	81
CENTRAL NERVOUS SYSTEM: STIMULANTS AND RELATED AGENTS	85
CENTRAL NERVOUS SYSTEM: NARCOLEPSY AGENTS.....	90
CENTRAL NERVOUS SYSTEM: NEUROPATHIC PAIN	94
CENTRAL NERVOUS SYSTEM: SEDATIVE HYPNOTICS.....	96
CENTRAL NERVOUS SYSTEM: SKELETAL MUSCLE RELAXANTS.....	99
CENTRAL NERVOUS SYSTEM: TOBACCO CESSATION	101
SPINAL MUSCULAR ATROPHY.....	102
ANALGESICS: NARCOTICS, LONG-ACTING.....	107
ANALGESICS: NARCOTICS, SHORT-ACTING	110
ANALGESICS: CLINICAL CRITERIA FOR SHORT-ACTING AND LONG-ACTING OPIOIDS.....	113
ANALGESICS: NARCOTICS, FENTANYL CITRATE PRODUCTS.....	116
ANALGESICS: NARCOTIC AGONISTS/ANTAGONISTS.....	117
ANALGESICS: NON-STEROIDAL ANTI-INFLAMMATORY DRUGS (NSAIDS)	118
ANALGESICS: OPIATE DEPENDENCE TREATMENTS.....	120
ANTI-INFECTIVE: ORAL ANTIFUNGALS.....	122
ANTI-INFECTIVE: ORAL ANTIVIRALS.....	124
ANTI-INFECTIVE: ORAL ANTIBIOTICS	126
ANTI-INFECTIVE: VAGINAL ANTIBIOTICS	131
ANTIRETROVIRALS: HUMAN IMMUNODEFICIENCY VIRUS/ACQUIRED IMMUNODEFICIENCY SYNDROME (HIV/AIDS)	132
HEPATITIS B AGENTS	137
HEPATITIS C AGENTS: INTERFERONS AND RIBAVIRINS.....	138
HEPATITIS C AGENTS: DIRECT-ACTING ANTIVIRALS	139
DIABETES: INSULINS AND RELATED AGENTS	142
DIABETES: GLUCAGON-LIKE PEPTIDE-1 (GLP-1) RECEPTOR AGONISTS.....	144
DIABETES: DIPEPTIDYL PEPTIDASE-4 (DPP-4) INHIBITORS.....	148
DIABETES: SODIUM-GLUCOSE COTRANSORTER-2 (SGLT2) INHIBITORS	150
DIABETES: ALPHA-GLUCOSIDASE INHIBITORS	152
DIABETES: METFORMINS	153
DIABETES: MEGLITINIDES.....	154
DIABETES: SULFONYLUREAS	155
DIABETES: THIAZOLIDINEDIONES (TZDS).....	156
ENDOCRINE AND METABOLIC AGENTS: GLUCAGON AGENTS.....	157
ENDOCRINE AND METABOLIC AGENTS: GROWTH HORMONES	158
ENDOCRINE AND METABOLIC AGENTS: GLUCOCORTICOIDS, ORAL (ORAL STEROIDS).....	161
ENDOCRINE AND METABOLIC AGENTS: PANCREATIC ENZYMES	162
ENDOCRINE AND METABOLIC AGENTS: PROGESTINS FOR CACHEXIA	163
ENDOCRINE AND METABOLIC AGENTS: ANDROGENIC AGENTS	164

ENDOCRINE AND METABOLIC AGENTS: BONE RESORPTION SUPPRESSION AND RELATED AGENTS	165
ENDOCRINE AND METABOLIC AGENTS: UTERINE DISORDER TREATMENTS	172
IMMUNOSUPPRESSANTS	173
IMMUNOLOGIC AND GENETIC: MULTIPLE SCLEROSIS AGENTS	176
IMMUNOLOGIC AND GENETIC: CYTOKINE AND CAM ANTAGONISTS	181
IMMUNOLOGIC AND GENETIC: IMMUNOMODULATORS, ASTHMA	199
IMMUNOLOGIC AND GENETIC: MUSCULAR DYSTROPHY AGENTS	206
IMMUNOMODULATORS, ATOPIC DERMATITIS	210
BLOOD MODIFIERS: ANTIHYPERURICEMICS	217
BLOOD MODIFIERS: COLONY STIMULATING FACTORS	219
BLOOD MODIFIERS: ERYTHROPOIESIS STIMULATING AGENTS	221
BLOOD MODIFIERS: PHOSPHATE BINDERS	224
BLOOD MODIFIERS: SICKLE CELL ANEMIA TREATMENTS	225
BLOOD MODIFIERS: THROMBOPOIESIS STIMULATING PROTEINS	227
OPHTHALMIC ANTIBIOTICS AND ANTIVIRALS	231
OPHTHALMICS FOR ALLERGIC CONJUNCTIVITIS	233
OPHTHALMICS: GLAUCOMA AGENTS	234
OPHTHALMICS: NON-STEROIDAL ANTI-INFLAMMATORY DRUGS (NSAIDS)	236
OPHTHALMICS: ANTI-INFLAMMATORY STEROIDS	237
OPHTHALMICS: IMMUNOMODULATORS	238
OPHTHALMIC: MYDRIATIC & CYCLOPLEGICS	240
OTICS	241
RENAL AND GENITOURINARY: ALPHA BLOCKERS FOR BPH & 5-ALPHA REDUCTASE INHIBITORS	242
BLADDER RELAXANTS	243
DERMATOLOGICS: TOPICAL ANTIBIOTIC AGENTS	245
DERMATOLOGICS: TOPICAL ANTIPARASITICS	246
DERMATOLOGICS: ORAL ANTIPSORIATICS	247
DERMATOLOGICS: ORAL ACNE AGENTS	248
DERMATOLOGICS: TOPICAL ACNE AGENTS	249
DERMATOLOGICS: TOPICAL ROSACEA AGENTS	251
DERMATOLOGICS: TOPICAL ANTIFUNGAL AGENTS	252
DERMATOLOGICS: TOPICAL ANTIVIRAL AGENTS	253
DERMATOLOGICS: TOPICAL ANTIPSORIATICS	254
DERMATOLOGICS: TOPICAL STEROIDS	256



Prior Authorization Criteria

Kentucky Medicaid

INTRODUCTION

The following document contains prior authorization criteria for agents on the Kentucky Medicaid Preferred Drug List. MedImpact's Clinical Team works closely with the Kentucky Pharmacy and Therapeutics (P&T) Committee to provide regular Therapeutic Class Reviews (TCR) to determine the State's Preferred Drug List (PDL) pursuant to KY statute 205.564. MedImpact follows an evidence-based approach when evaluating medication literature and developing recommendations for the P&T Committee. In addition to evaluating available clinical evidence, cost analyses are performed, and recommendations are developed within each therapeutic class to keep overall net costs manageable. Please note that all non-preferred agents on the PDL must meet the criteria for non-preferred agents in addition to brand or generic medically necessary criteria when applicable.

During regularly occurring meetings, the Kentucky Medicaid P&T Committee reviews information provided by MedImpact related to prior authorization criteria for new to market agents as well as recommendations on changes to the PDL. Once their review is complete, the P&T Committee submits their recommendations to the Kentucky Commissioner for final approval. MedImpact also works closely with the Pharmacy Director for the Department of Medicaid Services (DMS), or their designee, to develop and implement prior authorization criteria for medications already on the market. Clinical criteria may be updated if deemed appropriate when new information becomes available.

For a Managed Care Organization (MCO) member, prescribing providers may request a peer-to-peer review and/or an internal (first level) appeal upon denial of an initial prior authorization request. Appeal requests received from a prescribing provider, member, and/or member's representative (with the member's permission) may be submitted via phone, fax, or US mail within sixty (60) days of the receipt of the initial denial letter. Internal (first level) appeals will be reviewed and determined within 30 days. External (second level) appeals review by a third-party will be processed in accordance with 907 KAR 17:035 and must be made within thirty (30) days of an upheld internal (first level) appeal decision.

For a Fee-for service (FFS) member, providers may request a peer-to-peer review, a reconsideration, or an internal appeal on behalf of the member. Peer-to-peer requests may be initiated by phone or fax.

If needed, clinical support is available to assist with questions 8:00AM – 7:00PM Eastern Standard Time seven days per week and may be reached at:

- For MCO members:
 - Phone: (844) 336-2676
 - Fax: (858) 357-2612
- For FFS members:
 - Phone: (877) 403-6034
 - Fax: (858) 357-2612

CARDIOVASCULAR: ANGIOTENSIN RECEPTOR MODULATORS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires ≥ 30 -day trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Epaned ^{CC} Qbrelis ^{CC, QL}	<ul style="list-style-type: none">NPD criteria; ORUnable to swallow whole or consume crushed generic tablets.

CURRENT PDL STATUS

ACE INHIBITORS

Preferred Agents	Non-Preferred Agents
benazepril	Accupril
enalapril tablets	Altace
enalapril solution	captopril
lisinopril	Epaned ^{CC}
quinapril	fosinopril
ramipril	Lotensin
	moexipril
	perindopril
	Qbrelis ^{CC, QL}
	trandolapril
	Zestril

ACE INHIBITORS + DIURETIC COMBINATIONS

Preferred Agents	Non-Preferred Agents
benazepril/HCTZ	Accuretic
enalapril/HCTZ	captopril/HCTZ
fosinopril/HCTZ	Lotensin HCT
lisinopril/HCTZ	quinapril/HCTZ
	Zestoretic

ANGIOTENSIN RECEPTOR BLOCKERS (ARB)

Preferred Agents	Non-Preferred Agents
Entresto ^{QL}	Arbli ^{QL}
irbesartan	Atacand
losartan	Avapro
olmesartan	Benicar
valsartan tablet	candesartan
	Cozaar
	Diovan
	Edarbi
	Entresto Sprinkle
	eprosartan
	Micardis
	sacubitril/valsartan ^{QL}
	telmisartan
	valsartan solution

ARB + DIURETIC COMBINATIONS

Preferred Agents	Non-Preferred Agents
irbesartan/HCTZ	Atacand HCT
losartan/HCTZ	Avalide
olmesartan/HCTZ	Benicar HCT
valsartan/HCTZ	candesartan/HCTZ
	Diovan HCT
	Edarbyclor
	Hyzaar
	Micardis HCT
	telmisartan/HCTZ

DIRECT RENIN INHIBITORS

Preferred Agents	Non-Preferred Agents
N/A	aliskiren
	Tektuma

CARDIOVASCULAR: ANTI-ANGINAL & ANTI-ISCHEMIC AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Corlanor ^{CC} ivabradine ^{CC}	<ul style="list-style-type: none">• Diagnosis of chronic heart failure that is symptomatic; AND• Documentation (e.g., progress note) of:<ul style="list-style-type: none">◦ Left ventricular ejection fraction (LVEF) $\leq 35\%$; AND◦ Resting heart rate ≥ 70 beats per minute (bpm); AND• Used in combination with maximally tolerated doses of a beta blocker (e.g., bisoprolol, carvedilol, or metoprolol succinate), OR• Documentation (e.g., progress note) of clinical rationale preventing use of a beta-blocker.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
ranolazine ER	Aspruzyo Sprinkle ER ^{QL} Corlanor ^{CC} ivabradine ^{CC}

CARDIOVASCULAR: ANTIARRHYTHMICS (ORAL ANTI-ARRHYTHMICS)

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Sotylize ^{CC}	<ul style="list-style-type: none">NPD criteria; ORUnable to swallow sotalol/sotalol AF tablets

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
amiodarone 100, 200 mg	amiodarone 400 mg
disopyramide	Betapace
dofetilide	Betapace AF
flecainide	Multaq
mexiletine	Norpace
propafenone	Norpace CR
Sorine	Pacerone
sotalol	propafenone SR/ER
sotalol AF	quinidine sulfate

Preferred Agents	Non-Preferred Agents
	quinidine gluconate ER
	Rythmol SR
	Sotylize ^{CC}
	Tikosyn

CARDIOVASCULAR: BETA BLOCKERS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Hemangeol ^{CC}	<ul style="list-style-type: none">Diagnosis of Infantile Hemangioma (ICD-10 Disease Group D18)

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

BETA BLOCKERS

Preferred Agents	Non-Preferred Agents
Atenolol	acebutolol
atenolol/chlorthalidone	betaxolol
bisoprolol 2.5 mg ^{QL}	Bystolic
bisoprolol 5, 10 mg	carvedilol ER
bisoprolol/HCTZ	Coreg CR
Carvedilol	Coreg
Hemangeol ^{CC}	Inderal LA, XL
Labetalol	Innopran XL
metoprolol succinate ER	Kaspargo
metoprolol tartrate	Lopressor solution, tablet
Nadolol	metoprolol/HCTZ
Nebivolol	pindolol
propranolol ER	propranolol/HCTZ
propranolol solution	Tenoretic
propranolol tablet	Tenormin
	timolol
	Toprol XL

CARDIOVASCULAR: CALCIUM CHANNEL BLOCKERS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
nifedipine IR ^{CC}	<ul style="list-style-type: none">• Diagnosis of premature labor; OR• NPD criteria
nimodipine ^{CC}	<ul style="list-style-type: none">• Diagnosis of subarachnoid hemorrhage; OR• NPD criteria
Nymalize ^{CC}	<ul style="list-style-type: none">• Diagnosis of subarachnoid hemorrhage; AND• Unable to swallow capsules

CURRENT PDL STATUS

CALCIUM CHANNEL BLOCKERS

Preferred Agents	Non-Preferred Agents
amlodipine	Calan SR
Cartia XT	diltiazem ER 12HR capsule
diltiazem	Diltiazem ER (LA) tablet
diltiazem CD capsule	felodipine ER
diltiazem ER 24HR capsule	isradipine
diltiazem XR	Katerzia
Dilt-XR	levamlodipine
nifedipine ER	Matzim
Taztia XT	nicardipine
Tiadyt ER	nifedipine IR
verapamil tablet	nimodipine
verapamil ER tablet	nisoldipine ER
	Norliqva
	Norvasc
	Nymalize solution
	Nymalize syringe

Preferred Agents	Non-Preferred Agents
	Procardia XL
	Sular ER
	verapamil ER capsule
	verapamil ER PM capsule
	verapamil SR capsule
	Verelan PM

ANGIOTENSIN MODULATOR AND CALCIUM CHANNEL BLOCKER COMBINATIONS

Preferred Agents	Non-Preferred Agents
amlodipine/benazepril	amlodipine/valsartan/HCTZ
amlodipine/olmesartan	Azor
amlodipine/valsartan	Exforge HCT
	Exforge
	Lotrel
	Olmesartan/Amlodipine/HCTZ
	telmisartan/amlodipine
	trandolapril/verapamil
	Tribenzor

CARDIOVASCULAR: ANTICOAGULANTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Eliquis tablet, dose pack	Arixtra syringe
Enoxaparin syringe, vial	Dabigatran capsule
Jantoven tablet	Eliquis tablet for suspension
Pradaxa capsule	Eliquis sprinkle
Warfarin tablet	Fondaparinux sodium syringe
Xarelto dose pack, tablet	Fragmin syringe, vial
	Lovenox syringe, vial
	Pradaxa pellet pack pellet pack
	rivaroxaban tablet, suspension
	Savaysa tablet
	Xarelto suspension suspension

CARDIOVASCULAR: PLATELET AGGREGATION INHIBITORS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Brilinta	aspirin/dipyridamole
cilostazol	Effient
clopidogrel	Plavix
dipyridamole	ticagrelor
prasugrel	

CARDIOVASCULAR: PULMONARY ARTERIAL HYPERTENSION (PAH) AGENTS, ORAL AND INHALED

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Alyq ^{CC, QL} ambrisentan ^{CC} sildenafil suspension ^{CC} sildenafil tablet ^{CC} tadalafil ^{CC, QL} Tracleer tablet ^{CC, QL}	<ul style="list-style-type: none">Diagnosis of pulmonary hypertension (ICD-10 Disease Group I27)

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Opsynvi ^{CC, QL}	<p>Approval Duration: 1 year</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none">Diagnosis of pulmonary arterial hypertension (PAH) World Health Organization (WHO) Group 1; ANDPatient is WHO functional class (FC) 2 or 3; ANDPrescribed by, or in consultation with, a cardiologist, pulmonologist, or other specialist in the treatment of pulmonary arterial hypertension (PAH); ANDPatient has had at least a 30-day trial and failure, allergy, or contraindication (including potential drug-drug interactions with other medications) or intolerance of the following agents:<ul style="list-style-type: none">ambrisentan; ANDsildenafil or tadalafil; ANDPatient meets the minimum age recommended by the package insert for use in PAH; ANDPatient will not be using with other phosphodiesterase-5 inhibitors, e.g., sildenafil, tadalafil.

Agent(s) Subject to Criteria	Criteria for Approval
	Renewal Criteria: <ul style="list-style-type: none"> Prescriber attestation of clinically significant improvement or stabilization in clinical signs and symptoms.
Tracleer tablet suspension ^{CC}	Quantity Limit: 1 tablet per day <ul style="list-style-type: none"> PDP criteria; AND Unable to swallow Tracleer tablet.
Tyvaso, Tyvaso DPI ^{CC} Yutrepia ^{CC}	Approval Duration: 1 year Initial Approval Criteria: <p><i>Pulmonary Arterial Hypertension (PAH)</i></p> <ul style="list-style-type: none"> Diagnosis of Pulmonary Arterial Hypertension (PAH) WHO Group 1 Prescribed by, or in consultation with, a cardiologist or a pulmonologist Patient has trial and failure, allergy, contraindication, or intolerance to 2 or more preferred agents for at least 1 month <p><i>Pulmonary Hypertension Associated with Interstitial Lung Disease</i></p> <ul style="list-style-type: none"> Diagnosis of Pulmonary Hypertension Associated with Interstitial Lung Disease WHO Group 3 Prescribed by, or in consultation with, a cardiologist or a pulmonologist Baseline forced vital capacity < 70% for patients with connective tissue disease Patient had a right heart catheterization (documentation required) Results of the right heart catheterization confirm the diagnosis of WHO Group 3 interstitial lung disease associated with pulmonary hypertension Renewal Criteria: <ul style="list-style-type: none"> Patient has a documented response to therapy

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Alyq ^{CC, QL}	Adcirca ^{QL}
ambrisentan ^{CC}	Adempas ^{QL}
sildenafil suspension ^{CC}	bosentan tablet
sildenafil tablet ^{CC}	bosentan tablet suspension ^{QL}
tadalafil ^{CC, QL}	Letairis
Tracleer tablet ^{CC}	Liqrev
	Opsumit ^{QL}
	Opsynvi ^{CC, QL}
	Orenitram ER
	Revatio suspension ^{CC}
	Revatio tablet ^{CC}
	Tadliq
	Tracleer tablet suspension ^{CC, QL}
	Tyvaso ^{CC}
	Tyvaso DPI ^{CC}
	Upravi ^{QL}
	Ventavis
	Yutrepia

CARDIOVASCULAR: LIPOTROPICS

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Repatha ^{CC}	<p>Approval Duration: 6 months initial; 1 year renewal</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none">• Prescribed initially by, or in consultation with a cardiologist, lipid specialist, endocrinologist, vascular medicine or other applicable specialist; AND• Documentation of low-density lipoprotein cholesterol (LDL-C) prior to/without PCSK9 inhibitor therapy; AND• Documentation of trial and failure to achieve LDL goal after 3 months of maximum dose statin therapy recommended for patient age; OR• Prescriber attestation that patient does not tolerate statins (\geq 2 statin trials of any length were unsuccessful due to adverse effects); AND• Medication is prescribed for ONE of the following:<ul style="list-style-type: none">○ To reduce the risk of Major Adverse Cardiovascular Events (MACE) (e.g., cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) in a patient at increased risk for these events; OR○ Diagnosis of hyperlipidemia including Heterozygous and Homozygous familial hypercholesterolemia (HoFH); AND• Prescriber attestation that maximum tolerated doses of lipid-lowering therapies (e.g., statin, ezetimibe, omega-3-acid ethyl esters) to reduce LDL-C will continue to be used in combination with PCSK9 therapy; AND• Prescriber attestation that patient has been counseled to initiate and maintain diet and exercise modifications to reduce LDL-C in combination with PCSK9 therapy; AND• Patient meets the minimum age recommended by the package insert for the provided indication. <p>Renewal Criteria:</p> <ul style="list-style-type: none">• Patient must continue to meet initial authorization criteria; AND• Documentation (e.g., progress note or lab report) that demonstrate a reduction in LDL-C when compared to the baseline values.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 3 months** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
amlodipine/atorvastatin ^{CC, QL}	<ul style="list-style-type: none">• Trial and failure (e.g., poor adherence) of individual, generic components
Juxtapid ^{CC}	<p>Approval Duration: 6 months initial; 12 months renewal</p> <ul style="list-style-type: none">• Diagnosis of homozygous familial hypercholesterolemia (HoFH); AND• Prescribed by a cardiologist, lipid specialist, endocrinologist, vascular medicine or other applicable specialist; AND• Documentation (e.g., lab report) of cholesterol panel, including low density lipoprotein, cholesterol (LDL-C) prior to initiation; AND• Failure to achieve LDL-C goal on at least 3 of the following, unless contraindication:<ul style="list-style-type: none">○ Maximally tolerated or high-dose statin (e.g. atorvastatin 80mg, rosuvastatin 40mg)○ Ezetimibe○ PCSK9 inhibitor (e.g., alirocumab, evolocumab)○ Bempedoic acid <p>Renewal Criteria:</p> <ul style="list-style-type: none">• Documentation (e.g., progress note or lab report) that demonstrate a reduction in LDL-C when compared to the baseline values
Nexletol ^{CC, AE, QL} Nexlizet ^{CC, AE, QL}	<ul style="list-style-type: none">• Prescribed initially by, or in consultation with a cardiologist, lipid specialist, endocrinologist, vascular medicine or other applicable specialist; AND• Patient will be using Nexletol or Nexlizet for ONE of the following:<ul style="list-style-type: none">○ Diagnosis of primary hyperlipidemia, including heterozygous familial hypercholesterolemia (HeFH); OR○ To reduce the risk of myocardial infarction and coronary revascularization for those unable to take recommended statin therapy (including those not taking a statin) with established cardiovascular disease (CVD) or a high risk for a CVD event but without established CVD; AND• Patient meets at least ONE of the following:<ul style="list-style-type: none">○ Trial and failure to achieve LDL goal after 3 months of high intensity statin therapy (e.g., rosuvastatin 40 mg daily); OR○ Patient does not tolerate statins (≥ 2 statin trials of any length were unsuccessful due to adverse effects). <p>Age Limit: ≥ 18 years Quantity Limit: 1 per day</p>
Leqvio ^{CC, AE}	<p>Approval Duration: 6 months initial; 1 year renewal</p>

Agent(s) Subject to Criteria	Criteria for Approval
	<p>Initial Criteria</p> <ul style="list-style-type: none"> • Diagnosis of hypercholesterolemia, including heterozygous familial hypercholesterolemia (HeFH); AND • Prescribed initially by, or in consultation with, a cardiologist, lipid specialist, endocrinologist, vascular medicine or other applicable specialist; AND • Documentation of low-density lipoprotein cholesterol (LDL-C) prior to/without PCSK9 inhibitor therapy; AND • One of the following: <ul style="list-style-type: none"> ○ Trial and failure to achieve LDL goal after 3 months of high intensity statin therapy (e.g., rosuvastatin 40 mg daily); OR ○ Patient does not tolerate statins (≥ 2 statin trials of any length were unsuccessful due to adverse effects); AND • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to Repatha. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Documentation (e.g., progress note or lab report) that demonstrate a reduction in LDL-C when compared to the baseline values. <p>Age Limit: ≥ 18 years</p>
Praluent ^{CC}	<p>Approval Duration: 6 months initial; 1 year renewal</p> <p>Initial Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of primary hyperlipidemia, including heterozygous and homozygous familial hypercholesterolemia (HeFH); AND • Prescribed initially by, or in consultation with a cardiologist, lipid specialist, endocrinologist, vascular medicine or other applicable specialist; AND • Documentation of low-density lipoprotein cholesterol (LDL-C) prior to/without PCSK9 inhibitor therapy; AND • One of the following: <ul style="list-style-type: none"> ○ Trial and failure to achieve LDL goal after 3 months of high intensity statin therapy (e.g., rosuvastatin 40 mg daily); OR ○ Patient does not tolerate statins (≥ 2 statin trials of any length were unsuccessful due to adverse effects); AND • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to Repatha. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Documentation (e.g., progress note or lab report) that demonstrate a reduction in LDL-C when compared to the baseline values.

CURRENT PDL STATUS

LIPOTROPICS: OTHER

Preferred Agents	Non-Preferred Agents
cholestyramine powder, powder packet	colesevelam powder packet
cholestyramine light powder, powder packet	Colesevelam HCL tablet
colestipol tablet	Colestid granules, packet, tablet
ezetimibe tablet	colestipol granules, packet
fenofibrate capsule (generic Lofibra)	fenofibrate capsule (generic Lipofen)
fenofibrate nanocrystallized (generic Tricor)	fenofibrate tablet
fenofibric acid DR capsule	fenofibric acid tablet
gemfibrozil tablet	Fibracor tablet
niacin ER tablet	icosapent ethyl capsule
omega-3 acid ethyl esters capsules	Juxtapid capsule ^{CC}
Prevalite powder, powder packet	Leqvio syringe ^{CC, AE}
Repatha Pushtronex, SureClick, syringe ^{CC}	Lipofen capsule
	Lipid tablet
	Lovaza capsule
	Nexletol tablet ^{CC, AE, QL}
	Nexlizet tablet ^{CC, AE, QL}
	Praluent pen ^{CC}
	Questran powder, powder packet
	Questran Light powder
	TriCor tablet
	Trilipix DR capsule
	Vascepa capsule
	WelChol powder packet, tablet
	Zetia tablet

LIPOTROPICS: STATINS

Preferred Agents	Non-Preferred Agents
atorvastatin ^{QL}	Altoprev ^{QL}
lovastatin ^{QL}	amlodipine/atorvastatin ^{CC, QL}
pravastatin ^{QL}	Atorvaliq
rosuvastatin ^{QL}	Caduet ^{QL}
simvastatin ^{QL}	Crestor ^{QL}
	Ezallor Sprinkle ^{QL}
	ezetimibe/simvastatin ^{QL}
	Flolipid ^{QL}
	fluvastatin ^{QL}
	fluvastatin ER ^{QL}
	Lescol XL ^{QL}
	Lipitor ^{QL}
	Livalo ^{QL}
	pitavastatin
	Vytorin ^{QL}
	Zocor ^{QL}
	Zypitamag ^{QL}

GASTROINTESTINAL: ANTIEMETICS AND ANTIVERTIGO AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
dronabinol ^{CC, QL}	<ul style="list-style-type: none">• Diagnosis of nausea and/or vomiting (N/V) associated with cancer chemotherapy; AND<ul style="list-style-type: none">◦ Trial and failure of ≥ 1 non-cannabinoid antiemetic (e.g. ondansetron); OR• Diagnosis of anorexia associated with weight loss in a patient with acquired immune deficiency syndrome (AIDS) or cancer.
Diclegis ^{CC, QL}	<ul style="list-style-type: none">• Patient is a pregnant female; AND• Diagnosis of nausea and vomiting of pregnancy; AND• Documentation (e.g., progress note) of trial and failure of dietary and lifestyle modifications without adequate control of symptoms.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
doxylamine/pyridoxine ^{CC, QL}	<ul style="list-style-type: none">• Patient is a pregnant female; AND• Diagnosis of nausea and vomiting of pregnancy; AND• Documentation (e.g., progress note) of trial and failure of dietary and lifestyle modifications without adequate control of symptoms.
Cesamet ^{CC, QL}	<ul style="list-style-type: none">• NPD Criteria; AND• Dronabinol is one of the NPD drug trials.
Sancuso ^{CC, QL}	<ul style="list-style-type: none">• NPD criteria; OR• Used for preventing nausea and vomiting associated with moderately- or highly- emetogenic cancer chemotherapy.
Gimoti ^{CC, QL}	<p>Criteria for Initial Approval (duration 8 weeks):</p> <ul style="list-style-type: none">• Diagnosis of diabetic gastroparesis; AND• Prescribed by an endocrinologist, gastroenterologist or other specialist in the diagnosis and treatment of diabetic gastroparesis; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Prescriber attests that patient does NOT meet ANY of the following conditions: <ul style="list-style-type: none"> History of signs or symptoms of tardive dyskinesia (TD); History of a dystonic reaction to metoclopramide; Known or suspected circumstances where stimulation of gastrointestinal (GI) motility could be dangerous (e.g., GI hemorrhage, mechanical obstruction, or perforation); Known or suspected pheochromocytoma or other catecholamine-releasing paraganglioma; Diagnosis of epilepsy or any other seizure disorder; Hypersensitivity to metoclopramide (e.g., angioedema, bronchospasm); Moderate or severe renal impairment (creatinine clearance [CrCl] < 60 mL/minute); Moderate or severe hepatic impairment (Child-Pugh B or C); AND Prescriber attests that each course of treatment, with all dosage forms and routes of administration of metoclopramide, will NOT extend beyond 12 weeks; AND Adequate (e.g., 2-4 week) trial and failure of oral (e.g., tablet, solution, orally disintegrating tablet) or injectable (e.g., intramuscular) metoclopramide; OR NOT a candidate for oral metoclopramide (e.g., demonstrated or documented erratic absorption of oral medications). <p>Renewal Criteria (duration 8 weeks)</p> <ul style="list-style-type: none"> Must continue to meet initial authorization criteria; AND At least 2 weeks have passed (i.e., drug holiday) since completion of a previous course of metoclopramide treatment of any dosage form; AND Demonstrated improvement in signs and symptoms of diabetic gastroparesis (e.g., nausea, vomiting, early satiety, postprandial fullness, bloating, upper abdominal pain); AND Prescriber attestation that the patient is being monitored for extrapyramidal symptoms (e.g., tardive dyskinesia, dystonia) or other serious adverse events (e.g., suicidal ideation, fluid retention). <p>Age Limit: ≥ 18 years</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
aprepitant capsule dose pack ^{QL}	Akynzeo capsule ^{QL}
aprepitant capsule ^{QL}	Antivert chewable tablet
Bonjesta tablet	Antivert tablet
Diclegis tablet ^{CC, QL}	Anzemet tablet
dronabinol capsule ^{CC, QL}	Compro suppository
meclizine tablet	doxylamine/pyridoxine tablet ^{CC, QL}
metoclopramide solution	Emend capsule dose pack ^{QL}
metoclopramide tablet	Emend capsule ^{QL}
ondansetron solution	Emend suspension ^{QL}
ondansetron tablet	Gimoti nasal spray ^{CC, QL, AE}
ondansetron ODT	granisetron tablet
prochlorperazine tablet	Marinol capsule ^{CC, QL}
promethazine 12.5 mg, 25 mg suppository	prochlorperazine suppository
promethazine syrup	Promethegan 50 mg suppository
promethazine tablet	Reglan tablet
Promethegan 12.5 mg, 25 mg, suppository	Sancuso patch ^{CC, QL}

Preferred Agents	Non-Preferred Agents
scopolamine patch	Transderm-Scop patch trimethobenzamide capsule

GASTROINTESTINAL: ANTIDIARRHEALS

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Mytesi ^{CC, QL}	<p>Approval Duration: 6 months initial; 1 year renewal</p> <ul style="list-style-type: none">• Diagnosis of human immunodeficiency virus (HIV); AND• Current use of antiretroviral therapy for the treatment of HIV; AND• Active infection has been ruled out via fecal collection and microbiologic culture; AND• Trial and failure of 2 preferred agents. <p>Renewal Criteria:</p> <ul style="list-style-type: none">• Documented reduction in the frequency and quantity of liquid stool volume for the previous 6 months; AND• Documented monitoring/follow-up plan that includes re-culture for microbiologic agents if breakthrough diarrhea occurs while on crofelemer therapy. <p>Age Limit: ≥ 18 years Quantity Limit: 2 per day</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
diphenoxylate with atropine tablets	diphenoxylate with atropine liquid
loperamide capsule	Lomotil tablet
	Motofen tablet
	Mytesi tablet ^{CC, QL, AE}
	Opium tincture

GASTROINTESTINAL: ANTISPASMODICS/ANTICHOLINERGICS

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Cuvposa ^{CC}	<ul style="list-style-type: none"> BMN criteria; OR Inability to swallow whole or consume crushed glycopyrrolate tablets.
glycopyrrolate solution ^{CC}	<ul style="list-style-type: none"> NPD criteria; OR Inability to swallow whole or consume crushed glycopyrrolate tablets.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
dicyclomine capsule, solution, tablet	chlordiazepoxide/clindinium capsule
ED-Spaz ODT	Cuvposa solution ^{CC}
glycopyrrolate tablet	Dartisla ODT
hyoscyamine sulfate drops	Donnatal elixir
hyoscyamine sulfate elixir	Glycate tablet
hyoscyamine sulfate ER tablet	glycopyrrolate solution
hyoscyamine sulfate ODT tablet	Hyosyne drops
hyoscyamine sulfate SL tablet	Hyosyne elixir
hyoscyamine sulfate tablet	Levsin tablet
methoscopolamine tablet	Levsin/SL tablet
NuLev ODT	phenobarbital/hyoscyamine/atropine/scopolamine elixir
Oscimin SL tablet	phenobarbital/hyoscyamine/atropine/scopolamine tablet
Oscimin tablet	Phenohydro elixir
	Phenohydro tablet
	Robinul Forte tablet
	Robinul tablet

GASTROINTESTINAL: ANTI-ULCER PROTECTANTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Carafate suspension	Carafate tablet
misoprostol tablet	Cytotec tablet
sucralfate suspension	
sucralfate tablets	

GASTROINTESTINAL: BILE SALTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Bylvay ^{CC, QL}	<p>Approval Duration: 1 year</p> <p>Initial Approval Criteria <i>Progressive familial intrahepatic cholestasis (PFIC)</i></p> <ul style="list-style-type: none">• Patient is diagnosed with progressive familial intrahepatic cholestasis (PFIC) type 1 or type 2, confirmed by a genetic test; AND• Odevixibat is prescribed by or in consultation with a specialist (e.g., gastroenterologist, hepatologist, dermatologist); AND• Patient has elevated serum bile acid concentration; AND• Patient experiences persistent moderate to severe pruritus; AND• Patient has a history of trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to at least 1 pruritus treatment (e.g., ursodiol, cholestyramine, rifampin, naloxone, naltrexone, antihistamine) <i>Note: use of these agents is off-label.</i> <p>Renewal Criteria</p> <ul style="list-style-type: none">• Patient has experienced a reduction in serum bile acids from baseline; AND• Patient has experienced an improvement in pruritus <p>Initial Approval Criteria <i>Alagille syndrome</i></p> <ul style="list-style-type: none">• Patient is diagnosed with Alagille syndrome; AND• Odevixibat is prescribed by or in consultation with a specialist (e.g., gastroenterologist, hepatologist, dermatologist); AND• Patient has evidence of cholestasis, as evidenced by ≥ 1 of the following:<ul style="list-style-type: none">○ Serum bile acid > 3 times upper limit of normal (ULN) for age○ Conjugated bilirubin > 1 mg/dL○ Gamma glutamyl transferase (GGT) > 3 times ULN for age○ Fat soluble vitamin deficiency not otherwise explained

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ Intractable pruritus only explained by liver disease; AND • Patient experiences persistent moderate to severe pruritus; AND • Patient has a history of trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to at least 1 pruritic treatment (e.g., ursodiol, cholestyramine, rifampin, naloxone, naltrexone, antihistamine). <i>Note: use of these agents is off-label</i> <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient has experienced a reduction in serum bile acids from baseline <p>Quantity Limit:</p> <ul style="list-style-type: none"> • 200 mcg oral pellets: 2 per day; 60 per 30 days • 400 mcg capsule: 2 per day; 60 per 30 days • 600 mcg oral pellets: 5 per day; 150 per 30 days • 1,200 mcg capsule: 6 per day; 180 per 30 days
Iqirvo ^{CC, QL}	<p>Approval Duration: 1 year</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of primary biliary cholangitis (PBC); AND • Prescribed by, or in consultation with a gastroenterologist, hepatologist, or other disease state specialist; AND • Patient meets one of the following: <ul style="list-style-type: none"> ○ Patient has had a 12-month trial and failure of ursodiol, and will take Iqirvo in addition to current therapy; OR ○ Patient has a contraindication or intolerance to ursodiol and will take Iqirvo as monotherapy; AND • Patient has an alkaline phosphatase (ALP) level greater than 200 IU/L; AND • Patient does not have decompensated cirrhosis; AND • Patient meets the minimum age recommended by the package insert. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Documentation (e.g., progress notes, labs) of improvement or stabilization in alkaline phosphatase (ALP); AND • Patient meets one of the following: <ul style="list-style-type: none"> ○ Patient has had a 12-month trial and failure of ursodiol, and will take Iqirvo in addition to current therapy; OR ○ Patient has a contraindication or intolerance to ursodiol and will take Iqirvo as monotherapy <p>Quantity Limit: 1 tablet per day</p>
Livdelz ^{AE, CC, QL}	<p>Approval Duration: 1 year</p> <p>Initial Approval Criteria:</p> <p>Diagnosis of primary biliary cholangitis (PBC); AND</p> <p>Prescribed by, or in consultation with, a gastroenterologist, hepatologist, or other disease state specialist; AND</p> <ul style="list-style-type: none"> • Patient meets one of the following: <ul style="list-style-type: none"> ○ Patient has had a 12-month trial and failure of ursodiol, and will take Livdelzi in addition to current therapy; OR ○ Patient has a contraindication or intolerance to ursodiol and will take Livdelzi as monotherapy; AND • Patient has an alkaline phosphatase (ALP) level greater than 200 IU/L; AND • Patient does not have decompensated cirrhosis; AND

Agent(s) Subject to Criteria	Criteria for Approval
Livmarli ^{CC, QL}	<ul style="list-style-type: none"> Patient meets the minimum age recommended by the package insert for the provided <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Documentation (e.g., progress notes, labs) of improvement or stabilization in alkaline phosphatase (ALP); AND Patient meets one of the following: Patient has had a 12-month trial and failure of ursodiol and will take Livdelzi in addition to current therapy; OR Patient has a contraindication or intolerance to ursodiol and will take Livdelzi as monotherapy. <p>Age Limit: 18 years of age or older Quantity Limit: 1 capsule per day Approval Duration: 1 year</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> Diagnosis of one of the following conditions: <ul style="list-style-type: none"> Alagille Syndrome; OR Progressive familial intrahepatic cholestasis (PFIC) type 1 or type 2, confirmed by a genetic test; AND Maralixibat is prescribed by or in consultation with a specialist (e.g., gastroenterologist, hepatologist, dermatologist); AND Patient has evidence of cholestasis, as evidenced by ≥ 1 of the following: <ul style="list-style-type: none"> Serum bile acid > 3 times upper limit of normal (ULN) for age Conjugated bilirubin > 1 mg/dL Gamma glutamyl transferase (GGT) > 3 times ULN for age Fat soluble vitamin deficiency not otherwise explained Intractable pruritus only explained by liver disease; AND Patient experiences persistent moderate to severe pruritus; AND Patient has a history of trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to at least 1 pruritus treatment (e.g., ursodiol, cholestyramine, rifampin, naloxone, naltrexone, antihistamine, sertraline). Note: use of these agents is off-label; AND Patient must meet the minimum age recommended by the package insert for the FDA-approved indication. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Prescriber attests that the patient has experienced an improved clinical response (e.g., a reduction in serum bile acids from baseline, an improvement in pruritus). <p>Quantity Limit:</p> <ul style="list-style-type: none"> 9.5 mg/mL oral solution: 3 mL per day 19 mg/mL oral solution: 2 mL per day 10 mg, 15 mg, 20 mg oral tablet: 2 per day 30 mg oral tablet: 1 per day

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
ursodiol capsule	Bylvay capsule ^{CC, QL}
ursodiol tablet	Bylvay pellet ^{CC, QL}
	Chenodal tablet
	Cholbam capsule
	Iqirvo tablet ^{CC, QL}

Preferred Agents	Non-Preferred Agents
	Livdelzi ^{AE, CC, QL}
	Livmarli solution ^{CC, QL}
	Livmarli tablet ^{CC, QL}
	Reltone capsule
	Urso Forte tablet
	Urso tablet

GASTROINTESTINAL: HELICOBACTER PYLORI (H. PYLORI) TREATMENT

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure of a complete course of therapy, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of the preferred agent OR combination therapy comprised of individual, generic agents (e.g. lansoprazole and amoxicillin and clarithromycin).

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Voquezna Dual Pak ^{AE, CC, QL} Voquezna Triple Pak ^{AE, CC, QL}	<p>Approval Duration: 30 days</p> <ul style="list-style-type: none">Diagnosis of diagnostically confirmed <i>H. pylori</i> infection; ANDPrescribed by, or in consultation with, a gastroenterologist or other specialist in the diagnosis and treatment of <i>H. pylori</i>; ANDPatient has had a ≥ 2-week trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to Pylera. <p>Age Limit: ≥ 18 years of age</p> <p>Quantity Limit:</p> <ul style="list-style-type: none">Voquezna Dual Pak: 1 carton of 28 tablets and 84 capsules per 14-day supplyVoquezna Triple Pak: 1 carton of 56 tablets and 56 capsules per 14-day supply

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Pylera capsule ^{QL}	bismuth subcitrate potassium/metronidazole/ tetracycline capsule ^{QL} lansoprazole/amoxicillin/clarithromycin pack ^{QL}

Preferred Agents	Non-Preferred Agents
	Omeclamox-Pak ^{QL}
	Talicia capsule
	Voquezna Dual Pak ^{AE, CC, QL}
	Voquezna Triple Pak ^{AE, CC, QL}

GASTROINTESTINAL: HISTAMINE II (H₂) RECEPTOR BLOCKERS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
famotidine suspension famotidine tablet	cimetidine solution, tablet nizatidine capsule

GASTROINTESTINAL: LAXATIVES AND CATHARTICS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure or failed bowel preparation course, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
constulose solution	alvimopan capsule
enulose solution	Clenpiq solution
Gavilyte-C	GoLyteLy solution
Gavilyte-G	Kristalose packet
Gavilyte-N	lactulose packet
generlac solution	PEG 3350/Sod Sul/NaCl/KCl/AsbC powder packet
lactulose solution	Sodium Sulfate/Potassium Sulfate/Magnesium Sulfate solution
PEG 3350/Electrolyte solution	Suflave solution
PEG-3350 and Electrolytes	Suprep solution
	Sutab tablet

GASTROINTESTINAL: GASTROINTESTINAL MOTILITY AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Lubiprostone ^{AE, CC, QL}	<ul style="list-style-type: none"> • Diagnosis of one of the following conditions: <ul style="list-style-type: none"> ○ Chronic idiopathic constipation (CIC); OR ○ Irritable bowel syndrome with constipation (IBS-C); OR ○ Opioid-induced constipation (OIC) associated with the treatment of chronic, non-cancer pain [including patients with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation]; AND ○ Trial and failure of ≥ 1 over-the-counter (OTC) laxative (e.g., polyethylene glycol 3350). <p>Age Limit: ≥ 18 years Quantity Limit: 2 per day</p>
Linzess ^{AE, CC, QL}	<ul style="list-style-type: none"> • Patient is at least 6 years old; AND <ul style="list-style-type: none"> ○ Diagnosis of functional constipation (FC); OR • Patient is at least 18 years old; AND <ul style="list-style-type: none"> ○ Diagnosis of one of the following conditions: <ul style="list-style-type: none"> ▪ Chronic idiopathic constipation (CIC); OR ▪ Irritable bowel syndrome with constipation (IBS-C). • Trial and failure of ≥ 1 over-the-counter (OTC) laxative (e.g., polyethylene glycol 3350). <p>Age Limit: 72 mcg capsule (≥ 6 years old); 145 mcg (> 7 years old); and 290 mcg capsule (≥ 18 years old) Quantity Limit: 1 per day</p>
Movantik ^{AE, CC, QL}	<ul style="list-style-type: none"> • Diagnosis of opioid-induced constipation (OIC) associated with the treatment chronic, non-cancer pain [including patients with chronic pain related to prior cancer or its treatment who do NOT require frequent (e.g., weekly) opioid dosage escalation]; AND • Trial and failure of ≥ 1 over-the-counter (OTC) laxative (e.g., polyethylene glycol 3350). <p>Age Limit: ≥ 18 years Quantity Limit: 1 per day</p>

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
alosetron ^{AE, CC, QL} Lotronex ^{AE, CC, QL}	<ul style="list-style-type: none">Diagnosis of severe irritable bowel syndrome with diarrhea (IBS-D); ANDPatient is female; ANDTrial and failure of the specified length of, contraindication or intolerance to, ≥ 3 agents among the following drug classes (used separately or in combination):<ul style="list-style-type: none">Antidiarrheals (e.g., diphenoxylate/atropine, loperamide) for ≥ 1 monthBile acid sequestrants for ≥ 1 monthAntispasmodics (e.g., dicyclomine, hyoscyamine) for ≥ 1 month <p>Age Limit: ≥ 18 years</p>
Amitiza ^{AE, CC, QL}	<ul style="list-style-type: none">Diagnosis of one of the following conditions:<ul style="list-style-type: none">Chronic idiopathic constipation (CIC); ORIrritable bowel syndrome with constipation (IBS-C); OROpioid-induced constipation (OIC) associated with the treatment of chronic, non-cancer pain [including patients with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation]; ANDTrial and failure of ≥ 1 over-the-counter (OTC) laxative (e.g., polyethylene glycol 3350); ANDPatient has had at least a 1-month trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents; ANDPatient has had a trial and failure (e.g., allergy or intolerance to an inactive ingredient) with at least TWO manufacturers (if available) of the corresponding generic. <p>Age Limit: ≥ 18 years Quantity Limit: 2 per day</p>
Ibsrela ^{AE, CC, QL}	<ul style="list-style-type: none">Diagnosis of severe irritable bowel syndrome with constipation (IBS-C); ANDPatient does NOT have known or suspected mechanical GI obstruction; ANDPatient does NOT have severe diarrhea; ANDPatient has failed on 1 of the following regimens:<ul style="list-style-type: none">Osmotic laxatives; ORAntispasmodics; AND

Agent(s) Subject to Criteria	Criteria for Approval
Symproic AE, CC, QL	<ul style="list-style-type: none"> Patient has had at least a 1-month trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents. <p>Age Limit: ≥ 18 years Quantity Limit: 60 tablets per 30 days</p>
Viberzi AE, CC, QL	<ul style="list-style-type: none"> Diagnosis of opioid-induced constipation (OIC) related to chronic non-cancer pain [including patients with chronic pain related to prior cancer or its treatment who do NOT require frequent (e.g., weekly) opioid dosage escalation]; AND Patient has been using opioids for at least 150 days within past 180 days; AND Trial and failure of ≥ 2 preferred agents in this class; AND Trial and failure of ≥ 2 different laxative drug classes, such as: <ul style="list-style-type: none"> Stool softeners (e.g., docusate) Stimulant laxatives (e.g., bisacodyl, sennosides) Osmotic or saline laxatives (e.g., polyethylene glycol 3350) Bulk forming laxatives (e.g., psyllium) Lubricant laxatives (e.g., mineral oil) Patient does NOT have any the following conditions: <ul style="list-style-type: none"> Known or suspected gastrointestinal obstruction Pregnancy Severe hepatic impairment (Child-Pugh Class C) <p>Age Limit: ≥ 18 years Quantity Limit: 1 per day</p>
	<ul style="list-style-type: none"> Diagnosis of irritable bowel syndrome with diarrhea (IBS-D); AND Trial and failure of the specified length of, contraindication or intolerance to, ≥ 3 agents among the following drug classes (used separately or in combination): <ul style="list-style-type: none"> Antidiarrheals (e.g., diphenoxylate/atropine, loperamide) for ≥ 1 month Bile acid sequestrants for ≥ 1 month Antispasmodics (e.g., dicyclomine, hyoscyamine) for ≥ 1 month <p>Age Limit: ≥ 18 years Quantity Limit: 2 per day</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Linzess AE, CC, QL	alosetron tablet AE, CC, QL
lubiprostone AE, CC, QL	Amitiza AE, CC, QL
Movantik AE, CC, QL	Ibsrela AE, CC, QL
Trulance	Lotronex AE, CC, QL
	Motegrity AE, QL
	prucalopride AE, QL
	Symproic AE, CC, QL
	Relistor
	Viberzi AE, CC, QL

GATROINTESTINAL: PROTON PUMP INHIBITORS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 2-week** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Konvomep ^{CC, QL}	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none">Patient had at least a 2-week trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to 2 preferred agents; ORFor G-tube administration, the patient had at least a 2-week trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to the preferred liquid agent of Nexium suspension packets [Rx only]. <p><i>For twice daily dosing</i></p> <ul style="list-style-type: none">Patient has at least one of the following:<ul style="list-style-type: none">Diagnosis of <i>H. pylori</i>; AND/ORHad at least a 2-week trial and failure of once daily dosing. <p>Renewal Criteria:</p> <ul style="list-style-type: none">Patient has been on this medication and the request is for once daily dosing; ORPatient has a reconfirmation diagnosis of <i>H. pylori</i> infection or a trial of once daily dosing after initial course of therapy; ORPatient has undergone diagnostic testing (i.e., endoscopy, UGI, EGD with biopsy, serum gastrin or serum secretin stimulation test) to confirm one of the following diagnoses<ul style="list-style-type: none">Barrett's EsophagusSchatzki's Ring

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ A hypersecretory condition (including but not limited to Zollinger-Ellison Syndrome, Multiple Endocrine Adenoma, Systemic Mastocytosis) ○ Hyperacidity in Cystic Fibrosis Recipients, Achalasia, CREST syndrome, Scleroderma, Sarcoid; OR • Patient has attempted 'step-down' therapy at any time from multiple to single daily dosing for at least 2 weeks.
Voquezna ^{AE, CC, QL}	<p>For Erosive Esophagitis: Approval Duration: 8 weeks initial approval, 6 months renewal Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of diagnostically confirmed erosive esophagitis; AND • Prescribed by, or in consultation with, a gastroenterologist or other specialist in the diagnosis and treatment of erosive esophagitis; AND • Patient has had a ≥ 2-week trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to 2 preferred agents in this PDL class. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of diagnostically confirmed erosive esophagitis; AND • Prescribed by, or in consultation with, a gastroenterologist or other specialist in the diagnosis and treatment of erosive esophagitis; AND • Patient has experienced symptom improvement or control during initial treatment course. <p>For Non-Erosive Gastroesophageal Reflux Disease: Approval Duration: 4 weeks Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of non-erosive gastroesophageal reflux disease; AND • Prescribed by, or in consultation with, a gastroenterologist; AND • Patient has had a ≥ 2-week trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to 2 preferred agents in this PDL class. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of non-erosive gastroesophageal reflux disease; AND • Prescribed by, or in consultation with, a gastroenterologist; AND • Patient has received no more than one 20-week authorization in the past 365 days; AND • Patient has experienced symptom improvement or control during initial treatment course. <p>Age Limit: ≥ 18 years of age Quantity Limit: 1 tablet per day</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
esomeprazole capsule ^{QL}	Aciphex tablet ^{QL}
lansoprazole capsule ^{QL}	Dexilant capsule ^{QL}
Nexium suspension ^{QL}	dexlansoprazole DR capsule ^{QL}

Preferred Agents	Non-Preferred Agents
omeprazole capsule ^{QL}	esomeprazole suspension ^{QL}
pantoprazole tablets ^{QL}	Konvomep suspension ^{CC, QL}
	lansoprazole ODT ^{QL}
	Nexium Capsule ^{QL}
	omeprazole/sodium bicarbonate capsule ^{QL}
	omeprazole/sodium bicarbonate packet ^{QL}
	pantoprazole suspension ^{QL}
	Prevacid capsule ^{QL}
	Prevacid tablet ^{QL}
	Prilosec suspension ^{QL}
	Protonix suspension ^{QL}
	Protonix tablet ^{QL}
	rabeprazole tablet ^{QL}
	Voquezna tablet ^{AE, CC, QL}

GATROINTESTINAL: ULCERATIVE COLITIS AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
balsalazide capsule	Asacol HD tablet
Lialda tablet	Azulfidine tablet
mesalamine enema	Azulfidine EN-Tabs
mesalamine kit	budesonide ER tablet
mesalamine suppository	budesonide rectal foam
Pentasa capsule	Canasa suppository
sulfasalazine DR tablet	Delzicol capsule
sulfasalazine tablet	Dipentum capsule
	mesalamine DR capsule
	mesalamine DR tablet
	mesalamine ER capsule
	Rowasa enema

RESPIRATORY: ANTIBIOTICS, INHALED

GUIDELINES FOR USE

Approval Duration: 1 year; Arikayce – 3 months initial; 1 year renewal

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Arikayce ^{AE, CC, QL}	<ul style="list-style-type: none">Diagnosis of mycobacterium avium complex (MAC) lung disease as determined by the following:<ul style="list-style-type: none">Chest radiography or high-resolution computed tomography (HRCT) scan; ANDAt least 2 positive sputum cultures; ANDOther conditions such as tuberculosis and lung malignancy have been ruled out; ANDPatient has failed a multi-drug regimen with a macrolide (clarithromycin or azithromycin), rifampin, and ethambutol. (Failure is defined as continual positive sputum cultures for MAC while adhering to a multi-drug treatment regimen for a minimum duration of 6 months); ANDPatient has documented failure or intolerance to aerosolized administration of amikacin solution for injection, including pretreatment with a bronchodilator (cavitary or advanced/severe bronchiectatic or macrolide-resistant MAC pulmonary disease ONLY); ANDArikayce will be prescribed in conjunction with a multi-drug antimycobacterial regimen. <p>Age Limit: ≥ 18 years Quantity Limit: 1 kit per 28 days (1 vial per day)</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Bethkis ^{QL}	Arikayce ^{AE, CC, QL}

Preferred Agents	Non-Preferred Agents
Kitabis Pak ^{QL}	Cayston ^{QL}
tobramycin inhalation solution ^{QL} (generic for TOBI)	TOBI ^{QL}
	TOBI Podhaler ^{QL}
	tobramycin inhalation solution ^{QL} (generic for Bethkis and Kitabis Pak)

RESPIRATORY: ANTIHISTAMINES, MINIMALLY SEDATING

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
cetirizine solution	Clarinet tablet
levocetirizine tablet	Clarinet-D 12 HR tablet
	desloratadine ODT
	desloratadine tablet
	levocetirizine solution

RESPIRATORY: INTRANASAL RHINITIS AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Xhance ^{CC}	<ul style="list-style-type: none">Diagnosis of chronic rhinosinusitis with or without nasal polyps; ANDTrial and failure of high-dose generic fluticasone nasal spray.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
azelastine spray	azelastine/fluticasone nasal spray ^{QL}
Dymista nasal spray ^{QL}	Beconase AQ nasal spray ^{QL}
fluticasone propionate spray ^{QL}	flunisolide nasal spray ^{QL}
ipratropium bromide spray	mometasone nasal spray ^{QL}
olopatadine nasal spray	Omnaris nasal spray ^{QL}
	Patanase nasal spray
	Qnasl Children HFA ^{QL}
	Qnasl HFA ^{QL}
	Ryaltris nasal spray
	Xhance nasal spray ^{CC}
	Zetonna HFA ^{QL}

RESPIRATORY: LEUKOTRIENE MODIFIERS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
montelukast granules ^{AE, CC, QL}	<ul style="list-style-type: none">Under 6 years of age: no authorization required.6 years of age and older: clinical rationale that chewable or tablet cannot be used.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
montelukast granules ^{AE, CC, QL}	Accolate tablet ^{QL}
montelukast chewable tablet ^{QL}	Singulair granules ^{QL}
montelukast tablet ^{QL}	Singulair chewable tablet ^{QL}
	Singulair tablet ^{QL}
	zafirlukast ^{QL}
	zileuton ER ^{QL}
	Zyflo ^{QL}

RESPIRATORY: BRONCHODILATORS, BETA-AGONIST

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 14 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent within the same sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
AirDuo Respiclick ^{CC, QL, AE} AirDuo Digihaler ^{CC, QL}	<ul style="list-style-type: none">Trial and failure of at least two preferred agents, one of which must be Advair Diskus or Advair HFA.Age Limit: ≥ 12 years
arformoterol ^{CC, QL} formoterol ^{CC, QL}	<ul style="list-style-type: none">Diagnosis of chronic obstructive pulmonary disorder (COPD); ANDDocumentation of spirometry measurement; ANDNOT using any other long-acting beta adrenergic agonists (LABAs); ANDMust have a prescription for rescue therapy.Age Limit: ≥ 18 years

CURRENT PDL STATUS

BETA AGONISTS: COMBINATION PRODUCTS

Preferred Agents	Non-Preferred Agents
Advair Diskus ^{QL}	AirDuo Digihaler ^{CC, QL}
Advair HFA ^{QL}	AirDuo Respiclick ^{CC, QL, AE}
Dulera HFA ^{QL}	Breo Ellipta ^{QL}
Symbicort HFA ^{QL}	Breyna HFA ^{QL}
	budesonide/formoterol HFA ^{QL}
	fluticasone/salmeterol inhalation powder ^{QL}
	fluticasone/salmeterol HFA ^{QL}
	fluticasone/vilanterol ^{QL}

Preferred Agents	Non-Preferred Agents
	Wixela Inhub ^{QL}

LONG-ACTING BETA₂ ADRENERGIC AGONISTS

Preferred Agents	Non-Preferred Agents
Serevent Diskus ^{QL}	Arformoterol solution ^{AE, CC, QL}
	Brovana solution ^{CC, QL}
	Formoterol solution ^{AE, CC, QL}
	Perforomist ^{CC, QL}
	Striverdi Respimat ^{QL}

SHORT-ACTING BETA₂ ADRENERGIC AGONISTS

Preferred Agents	Non-Preferred Agents
albuterol sulfate solution ^{QL}	Airsupra HFA ^{QL}
Proventil HFA ^{QL}	albuterol sulfate HFA ^{QL}
terbutaline tablets ^{QL}	albuterol sulfate syrup ^{QL}
Ventolin HFA ^{QL}	albuterol sulfate ER tablet ^{QL}
	albuterol sulfate tablet ^{QL}
	levalbuterol concentrate solution ^{QL}
	levalbuterol HFA ^{QL}
	levalbuterol solution ^{QL}
	ProAir® Digihaler ^{QL}
	ProAir Respiclick ^{QL}
	Xopenex HFA ^{QL}

RESPIRATORY: GLUCOCORTICIDS, INHALED

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
budesonide inhalation suspension ^{AE, CC, QL}	<ul style="list-style-type: none">Under 8 years of age: no prior authorization required.8 years of age and older: clinical rationale (e.g., trial and failure, comorbid condition) that a metered dose inhaler (e.g., Flovent HFA) cannot be used.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Asmanex Twisthaler ^{QL}	Alvesco ^{QL}
budesonide inhalation suspension ^{AE, CC, QL}	ArmonAir Digihaler ^{QL}
Flovent HFA ^{QL}	Arnuity Ellipta ^{QL}
fluticasone propionate HFA ^{QL}	Asmanex HFA ^{QL}
Pulmicort Flexhaler ^{QL}	Flovent Diskus ^{QL}
	fluticasone furoate ^{QL}
	Pulmicort Respules ^{QL}
	Qvar Redihaler

RESPIRATORY: EPINEPHRINE, SELF-INJECTABLE

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
epinephrine 0.3 mg autoinjector ^{QL}	Auvi-Q autoinjector ^{QL}
epinephrine 0.15 mg autoinjector ^{QL}	Neffy ^{QL}
EpiPen ^{QL}	Symjepi ^{QL}
EpiPen Jr. ^{QL}	

RESPIRATORY: CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD) AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Breztri Aerosphere ^{AE, CC, QL}	<ul style="list-style-type: none">• Diagnosis of chronic obstructive pulmonary disorder (COPD); AND• 2-week trial and failure (e.g., limited ability to use or comply with multiple devices) of triple-ingredient therapy (such as, glucocorticoid, long-acting beta agonist, and long-acting muscarinic antagonist) among single- and dual-ingredient inhalers (e.g., Flovent HFA and Anoro Ellipta or Stiolto Respimat).
roflumilast ^{CC, QL}	<p>Age Limit: ≥ 18 years</p> <ul style="list-style-type: none">• Diagnosis of chronic obstructive pulmonary disorder (COPD); AND• Documentation (e.g., progress notes) of FEV¹ ≤ 50% of predicted.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 14 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Daliresp ^{AE, CC, QL}	<ul style="list-style-type: none">• Diagnosis of chronic obstructive pulmonary disorder (COPD); AND• Trial and failure of ≥ 1 inhaled therapy; AND• Documentation (e.g., progress notes) of FEV¹ ≤ 50% of predicted. <p>Approval Duration: 6 months initial, 1 year renewal</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none">• Diagnosis of moderate to severe chronic obstructive pulmonary disorder; AND• Trial and failure of at least a 2-week trial of standard of care therapy:<ul style="list-style-type: none">○ Triple-ingredient therapy (inhaled corticosteroids [ICS], long-acting beta agonist [LABA], and long-acting muscarinic antagonist [LAMA]); OR

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Dual-ingredient therapy (long-acting beta agonist [LABA]/long-acting muscarinic agent [LAMA]) <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Clinically significant improvement or stabilization in signs and symptoms <p>Age Limit: 18 years of age or older Quantity Limit: 5 mL per day</p>
Trelegy Ellipta ^{AE, CC, QL}	<p>Asthma</p> <ul style="list-style-type: none"> Diagnosis of asthma; AND Failure (e.g., limited ability to use or comply with multiple devices) of at least a 2-week trial of triple-ingredient therapy (glucocorticoid, long-acting beta agonist, and long-acting muscarinic antagonist) among single- and dual-ingredient inhalers (e.g., Flovent HFA and Bevespi Aerosphere). <p>COPD</p> <ul style="list-style-type: none"> Diagnosis of chronic obstructive pulmonary disease (COPD); AND Patient has had at least a 30-day trial and failure of Breztri Aerosphere. <p>Age Limit: ≥ 18 years</p>
Yupelri ^{AE, CC, QL}	<ul style="list-style-type: none"> Diagnosis of chronic obstructive pulmonary disorder (COPD); AND Demonstrate treatment failure with 1 other long-acting muscarinic antagonist (LAMA) agent due to technique/delivery mechanism (e.g., cannot use inhaler) <p>Age Limit: ≥ 18 years</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
albuterol-ipratropium inhalation solution ^{QL}	Bevespi Aerosphere ^{QL}
Anoro Ellipta ^{QL}	Daliresp tablet ^{AE, CC, QL}
Atrovent HFA ^{QL}	Duaklir Pressair
Breztri Aerosphere ^{AE, CC, QL}	Incruse Ellipta ^{QL}
Combivent Respimat ^{QL}	Ohtuvayre ^{AE, CC, QL}
ipratropium inhalation solution ^{QL}	Spiriva Respimat ^{QL}
roflumilast tablet ^{CC, QL}	tiotropium ^{QL}
Spiriva Handihaler ^{QL}	Trelegy Ellipta ^{AE, CC, QL}
Stiolto Respimat ^{QL}	Tudorza Pressair ^{QL}
	umeclidinium-vilanterol ^{QL}
	Yupelri solution ^{AE, CC, QL}

CENTRAL NERVOUS SYSTEM: ALZHEIMER'S AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 1 week** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
donepezil 23 mg ^{CC}	<ul style="list-style-type: none">Use of donepezil 10 mg tablets for ≥ 90 days.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
donepezil ODT	Adlarity patch
Donepezil 5 mg, 10 mg tablet	Aricept tablet, ODT
Exelon patch	donepezil 23 mg ^{CC}
memantine tablets	galantamine
rivastigmine capsule	galantamine ER
	memantine/donepezil
	memantine ER
	memantine solution
	Namzaric

Preferred Agents	Non-Preferred Agents
	Namenda tablets
	Namenda XR
	Razadyne
	Razadyne ER
	rivastigmine patch
	Zunveyi ^{QL}

CENTRAL NERVOUS SYSTEM: ANXIOLYTICS

GUIDELINES FOR USE

Approval Duration: 1 year (non-preferred approval)

1. MAXIMUM DURATION (MD) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
alprazolam IR tablets ^{MD} chlordiazepoxide ^{MD} diazepam oral solution, tablets ^{MD} diazepam oral concentrate ^{MD} lorazepam ^{MD} alprazolam ER/XR ^{MD} alprazolam intenzol ^{CC, MD} alprazolam ODT ^{CC, MD} clorazepate ^{MD} diazepam intenzol ^{CC, MD} lorazepam intenzol ^{CC, MD} oxazepam ^{MD} Xanax ^{MD} Xanax XR ^{MD}	<p>Preferred antianxiety benzodiazepines are available without a prior authorization for up to 60-day supply (cumulative) per rolling year.</p> <p>Approve for 1 month for the following diagnosis:</p> <ul style="list-style-type: none">• Acute alcohol withdrawal <p>Approve for 6 months for the following diagnoses / situations:</p> <ul style="list-style-type: none">• Agoraphobia• Anxiety• Anxiety disorder• Chemotherapy-induced nausea & vomiting• Depression• Panic attacks or panic disorder• Social phobia• Status epilepticus <p>Approve for 1 year for the following diagnosis:</p> <ul style="list-style-type: none">• Seizures/Epilepsy <p>NOTE: Prescriber (not pharmacy) must submit prior authorization request.</p>

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
alprazolam intenzol oral concentrate ^{CC, MD} alprazolam ODT ^{CC, MD} diazepam oral concentrate ^{CC, MD} lorazepam intenzol oral concentrate ^{CC, MD}	<ul style="list-style-type: none">• Therapeutic failure to at least a ONE-month trial of the tablet formulation of the drug being requested; OR• Clinical reason why the tablets cannot be used.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
alprazolam IR tablets ^{MD}	alprazolam ER tablet ^{MD}
buspirone tablet	alprazolam intensol oral concentrate ^{CC, MD}
chlordiazepoxide capsule ^{MD}	alprazolam ODT ^{CC, MD}
diazepam oral solution ^{MD}	alprazolam XR tablet ^{MD}
diazepam oral tablet ^{MD}	Bucapsol capsule
lorazepam tablet ^{MD}	clorazepate dipotassium tablet ^{MD}
	diazepam oral concentrate ^{CC, MD}
	lorazepam intensol oral concentrate ^{CC, MD}
	lorazepam oral concentrate ^{MD}
	Loreev XR capsule ^{MD}
	meprobamate tablet
	oxazepam capsule ^{MD}
	Xanax tablet ^{MD}
	Xanax XR tablet ^{MD}

CENTRAL NERVOUS SYSTEM: ANTICONVULSANTS

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to PA Criteria	Criteria for Approval
Banzel ^{CC, QL}	<ul style="list-style-type: none">• Diagnosis of Lennox-Gastaut Syndrome (LGS); OR• Trial and failure of 1 anticonvulsant.
phenobarbital ^{CC}	<p>Approval Duration: Lifetime</p> <ul style="list-style-type: none">• Diagnosis of epilepsy or seizure/convulsive disorder (ICD-10 Disease Group G40) <p>Approval Duration: 1 month</p> <ul style="list-style-type: none">• Diagnosis of alcohol withdrawal (ICD-10 Disease Group F10)
primidone ^{CC}	<p>Approval Duration: Lifetime</p> <ul style="list-style-type: none">• Diagnosis of epilepsy (ICD-10 Disease Group G40)• Diagnosis of tremor [G25.0 (essential tremor) or R25.1 (tremor, unspecified)]
Sabril ^{CC, QL}	<ul style="list-style-type: none">• Diagnosis of infantile spasms (ICD-10 = G40.401, G40.409, G40.411, G40.419); OR• Trial and failure of 1 anticonvulsant

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Briviact ^{CC, QL}	<ul style="list-style-type: none">• Diagnosis of partial-onset seizures; AND• Trial and failure of at least 1 preferred agent AND• ≥ 1 month of age.
clobazam syringe ^{AE, CC, QL}	<p>Approval Duration: 6 months initial, 1 year renewal</p> <p>Initial Approval Criteria</p>

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Diagnosis of seizures associated with Lennox-Gastaut syndrome (LGS) AND • Provider has submitted documentation supporting medical necessity for the 10mg/4mL syringe over the preferred 2.5mg/mL suspension, due to at least ONE of the following: <ul style="list-style-type: none"> ○ Volume/dosing limitations: Patient requires smaller volumes per dose for accuracy or compatibility with specialized devices (e.g., G-tube), OR ○ Intolerance: Documented adverse effects (e.g., GI upset, flavor aversion) impacting adherence, OR ○ Measurement barriers: Inability to accurately measure doses with the 2.5mg/mL syringe due to calibration limitations. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient continues to meet the above criteria; AND • Provider attests to improvement in seizure control or other relevant clinical outcome from baseline. <p>Quantity Limit: 16ml per day Age Limit: ≥ 2 years</p>
Diacomit ^{CC, QL}	<ul style="list-style-type: none"> • Diagnosis of Dravet syndrome; AND • Prescriber is, or has a consultative relationship with, a neurology/epilepsy specialist; AND • Medication will be used in combination with clobazam; AND • Trial and failure (e.g., incomplete seizure control) of ≥ 2 anticonvulsants.
Epidiolex ^{CC}	<ul style="list-style-type: none"> • Patient is ≥ 1 year old; AND • Diagnosis of: <ul style="list-style-type: none"> ○ Lennox-Gastaut syndrome (LGS); OR ○ Dravet syndrome (DS); OR ○ Tuberous Sclerosis Complex (TSC); AND • Prescriber is, or has a consultative relationship with, a neurology/epilepsy specialist; AND • Trial and failure (e.g., incomplete seizure control) of ≥ 2 anticonvulsants; AND • Must be used in adjunct with ≥ 1 anticonvulsant
levetiracetam tablet suspension ^{CC, QL}	<ul style="list-style-type: none"> • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to levetiracetam solution, levetiracetam tablet, and Spritam tablet suspension, AND • Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with at least 2 manufacturers (if available and covered) of levetiracetam solution and levetiracetam tablet, AND • Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the brand Spritam tablet suspension. <p>Quantity Limit: 2 per day</p>
Symptazan ^{AE, CC, QL}	<ul style="list-style-type: none"> • Diagnosis of Lennox-Gastaut syndrome (LGS). • Clinical rationale that clobazam suspension or tablets cannot be used. <p>Age Limit: ≥ 2 years</p>

Agent(s) Subject to Criteria	Criteria for Approval
Vigafyde solution ^{CC, QL}	<ul style="list-style-type: none"> Diagnosis of infantile spasms (IDC-10 = G40.401, G40.409, G40.411, G40.419); OR Trial and failure of 1 anticonvulsant; AND Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in Sabril.
Xcopri ^{AE, CC, QL}	<ul style="list-style-type: none"> Diagnosis of partial-onset seizures; AND Trial and failure of ≥ 1 preferred agent; AND NOT have familial QT syndrome; AND NOT have severe hepatic impairment (Child-Pugh Class C). <p>Age Limit: ≥ 18 years</p> <p>Quantity Limits:</p> <ul style="list-style-type: none"> 1 per day: 25 mg, 50 mg, 100 mg tablets; titration blister packs 2 per day: 150 mg, 200 mg; 250 and 350 mg maintenance blister packs
Ztalmy ^{AE, CC, QL}	<p>Approval Duration: 1 year</p> <p>Initial Approval Criteria</p> <ul style="list-style-type: none"> Patient is ≥ 2 years of age; AND Patient has a diagnosis of seizures associated with cyclin dependent kinase-like 5 (CDKL5) deficiency disorder (CDD) confirmed with genetic testing; AND Patient has tried ≥ 2 other anticonvulsant medications; AND Patient will avoid concomitant therapy with moderate or strong CYP450 inducers (e.g., carbamazepine, phenobarbital, phenytoin, omeprazole), or if concomitant therapy is unavoidable, dose adjustments will be considered; AND Ganaxolone is prescribed by or in consultation with a neurologist. <p>Renewal Criteria</p> <ul style="list-style-type: none"> Patient must continue to meet the above criteria; AND Prescriber attests to stabilization of disease or reduction in seizure frequency from baseline <p>Quantity Limit: 1800mg (36mL) per day</p> <p>Age Limit: 2 years of age</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Banzel suspension ^{CC, QL}	Aptiom tablet ^{QL}
Banzel tablet ^{CC, QL}	Briviact solution ^{CC, QL}
carbamazepine ER capsule	Briviact tablet ^{CC, QL}
carbamazepine ER tablet	carbamazepine suspension
carbamazepine tablet	Carbatrol ER capsule
carbamazepine chewable tablet	clobazam syringe ^{AE, CC, QL}
Celontin capsule	clonazepam ODT ^{QL}
clobazam suspension ^{QL}	Depakote ER tablet
clobazam tablet ^{QL}	Depakote sprinkle capsule
clonazepam tablet ^{QL}	Depakote tablet
diazepam rectal gel ^{QL}	Diacomit capsule ^{CC, QL}
divalproex sodium DR sprinkle capsule	Diacomit powder packet ^{CC, QL}
divalproex sodium DR tablet	Dilantin capsule
divalproex sodium ER tablet	Dilantin chewable tablet
Equetro	Dilantin-125 suspension

Preferred Agents	Non-Preferred Agents
ethosuximide capsule	Elepsia XR tablet ^{QL}
ethosuximide solution	Epidiolex solution ^{CC}
felbamate suspension	Epitol tablet
felbamate tablet	Eprontia solution
Gabitril tablet ^{QL}	eslicarbazepine tablet ^{QL}
lacosamide solution ^{QL}	Felbatol suspension
lacosamide tablet ^{QL}	Felbatol tablet
lamotrigine tablet	Fintepla solution ^{QL}
lamotrigine chewable tablet	Fycompa suspension
levetiracetam ER tablet ^{QL}	Fycompa tablet ^{QL}
levetiracetam solution ^{QL}	Keppra solution
levetiracetam tablet ^{QL}	Keppra tablet ^{QL}
Nayzilam spray ^{AE, QL}	Keppra XR tablet ^{QL}
oxcarbazepine suspension	Klonopin tablet ^{QL}
oxcarbazepine tablet ^{QL}	Lamictal tablet dose packs
phenobarbital elixir ^{CC}	Lamictal ODT dose packs
phenobarbital tablet ^{CC}	Lamictal XR tablet dose packs
phenytoin suspension	Lamictal ODT
phenytoin sodium ER capsule	Lamictal tablet
phenytoin chewable tablet	Lamictal chewable tablet
primidone tablet	Lamictal XR tablet ^{QL}
Roweepra tablet ^{QL}	lamotrigine tablet dose packs
Sabril powder packet ^{QL, CC}	lamotrigine ODT dose packs
Sabril tablet ^{CC, QL}	lamotrigine ER tablet ^{QL}
Tegretol suspension	lamotrigine ODT
tiagabine tablet ^{QL}	levetiracetam tablet suspension ^{CC, QL}
topiramate sprinkle capsule ^{QL}	Libervant film ^{AE, QL}
topiramate tablet ^{QL}	methsuximide capsule
valproic acid capsule	Motpoly XR capsule
valproic acid solution	Onfi suspension ^{QL}
Valtoco spray ^{QL}	Onfi tablet ^{QL}
zonisamide capsule ^{QL}	oxcarbazepine ER tablet ^{QL}
	Oxtellar XR tablet ^{QL}
	perampanel tablet ^{QL}
	Phenytek capsule
	Qudexy XR sprinkle capsule ^{QL}
	rufinamide suspension ^{QL}
	rufinamide tablet ^{QL}
	Spritam suspension ^{QL}
	subvenite oral suspension ^{AE, QL}
	subvenite tablet dose packs
	subvenite tablet
	Sympazan film ^{AE, CC, QL}
	Tegretol tablet
	Tegretol XR tablet
	Topamax sprinkle capsule ^{QL}
	Topamax tablet ^{QL}
	topiramate ER capsule ^{QL}
	topiramate ER sprinkle capsule ^{QL}
	topiramate solution
	Trileptal suspension
	Trileptal tablet ^{QL}
	Trokendi XR capsule ^{QL}
	vigabatrin powder packet ^{QL}
	vigabatrin tablet ^{QL}
	vigadrone powder packet ^{QL}
	vigadrone tablet ^{QL}
	Vigafyde solution ^{CC, QL}
	Vimpat solution ^{QL}
	Vimpat tablet ^{QL}
	Xcorpi tablet dose pack ^{AE, CC, QL}

Preferred Agents	Non-Preferred Agents
	Xcorpi tablet ^{AE, CC, QL}
	Zarontin capsule
	Zarontin solution
	zonisamide suspension ^{QL}
	Ztalmy suspension ^{AE, CC, QL}

CENTRAL NERVOUS SYSTEM: ANTIPSYCHOTICS: FIRST GENERATION (TYPICAL)

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
amitriptyline/perphenazine tablet	Adasuve inhalation powder
chlorpromazine tablet	molindone tablet
chlorpromazine oral concentrate	pimozide
fluphenazine elixir	
fluphenazine oral concentrate	
fluphenazine tablet	
haloperidol oral concentrate	
haloperidol tablet	
loxapine capsule	
perphenazine tablet	
thioridazine tablet	
thiothixene capsule	
trifluoperazine tablet	

CENTRAL NERVOUS SYSTEM: ANTIPSYCHOTICS: SECOND GENERATION (ATYPICAL) AND INJECTABLE

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to PA Criteria	Criteria for Approval
aripiprazole tablets ^{CC, QL} asenapine ^{CC, QL} clozapine tablets ^{CC, QL} lurasidone ^{CC, QL} quetiapine ^{CC, QL} quetiapine ER ^{CC, QL} risperidone ^{CC, QL} Vraylar ^{AE, CC, QL} ziprasidone capsules ^{CC, QL} Abilify Maintena ^{AE, CC, QL} Aristada ER ^{AE, CC, QL} Aristada Initio ^{AE, CC, QL} fluphenazine decanoate ^{AE, CC, QL} Geodon injection ^{AE, CC, QL} haloperidol decanoate ^{AE, CC, QL} haloperidol lactate ^{AE, CC, QL} Invega Sustenna ^{AE, CC, QL} olanzapine ODT, tablet ^{CC, QL} olanzapine vial ^{AE, CC, QL} Perseris ER ^{AE, CC} Risperdal Consta ^{AE, CC, QL} Abilify Asimtufii ^{AE, CC, QL}	<ul style="list-style-type: none"> Diagnosis of any of the following conditions: <ul style="list-style-type: none"> Dementias (ICD-10 Disease Groups F01, F02, F03, F06); Dissociative and conversion disorders (ICD-10 Disease Group F44); Episodic Mood Disorders (ICD-10 Disease Groups F30, F31, F39); Huntington's disease (ICD-10 Disease Group G10); Major depressive disorder (ICD-10 Disease Groups F32, F33); Oppositional defiant disorder (ICD-10 = F91.3); Pervasive developmental disorders (ICD-10 Disease Group F84); Schizoaffective disorder (F25.9); Schizophrenic Disorders (ICD-10 Disease Group F20; ICD-10 = F60.1); Tic disorder (ICD-10 Disease Group F95); Substance use disorders and related conditions (see below for list).
	<ul style="list-style-type: none"> Patient has a diagnosis of bipolar disorder or schizophrenia <p>Age Limit: ≥ 18 years Quantity Limit: 1 syringe every 56 days</p>
Invega Hafyera ^{AE, CC, QL}	<ul style="list-style-type: none"> Patient is ≥ 18 years of age; AND Patient has a confirmed diagnosis of schizophrenia; AND Patient has received a minimum of 4 months of monthly injections with Invega Sustenna® with adequate response and acceptable patient tolerance; OR Patient has received a minimum of one 3 month injection of Invega Trinza® with adequate response and acceptable patient tolerance.
Invega Trinza ^{AE, CC, QL}	<ul style="list-style-type: none"> Patient is ≥ 18 years of age; AND Patient has a confirmed diagnosis of schizophrenia; AND Patient has received a minimum of 4 months of monthly injections with Invega Sustenna® with adequate response and acceptable patient tolerance.
Uzedy ^{AE, CC, QL}	<ul style="list-style-type: none"> Patient has a diagnosis of schizophrenia OR bipolar 1 disorder; AND If used for bipolar I disorder, will be used for either: <ul style="list-style-type: none"> Maintenance monotherapy treatment; OR Maintenance therapy as adjunct to lithium or valproate. <p>Age Limit: ≥ 18 years</p>

Agent(s) Subject to PA Criteria		Criteria for Approval
		Quantity Limit: 1 syringe per 30 days
Approvable ICD-10 Group	Approvable ICD-10	Approvable ICD-10 Description
	F60.1	Schizoid personality disorder
F01		Vascular dementia
F02		Dementia in other diseases classified elsewhere
F03		Unspecified dementia
F06		Other mental disorders due to known physiological condition
F20		Schizophrenia, schizotypal and delusional, and other non-mood psychotic disorders
	F25.9	Schizoaffective disorder, unspecified
F30		Manic episode
F31		Bipolar disorder
F32		Major depressive disorder, single episode
F33		Major depressive disorder, recurrent
F39		Unspecified mood [affective] disorders
F44		Dissociative and conversion disorders
F84		Pervasive developmental disorders
	F11.150	Opioid abuse with opioid-induced psychotic disorder with delusions
	F11.250	Opioid dependence with opioid-induced psychotic disorder with delusions
	F11.950	Opioid use, unspecified with opioid-induced psychotic disorder with delusions
	F12.150	Cannabis abuse with psychotic disorder with delusions
	F12.250	Cannabis dependence with psychotic disorder with delusions
	F12.950	Cannabis use, unspecified with psychotic disorder with delusions
	F13.150	Sedative, hypnotic or anxiolytic abuse with sedative, hypnotic or anxiolytic-induced psychotic disorder with delusions
	F13.250	Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced psychotic disorder with delusions
	F13.950	Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced psychotic disorder with delusions
	F14.150	Cocaine abuse with cocaine-induced psychotic disorder with delusions
	F14.250	Cocaine dependence with cocaine-induced psychotic disorder with delusions

Approvable ICD-10 Group	Approvable ICD-10	Approvable ICD-10 Description
	F14.950	Cocaine use, unspecified with cocaine-induced psychotic disorder with delusions
	F15.150	Other stimulant abuse with stimulant-induced psychotic disorder with delusions
	F15.250	Other stimulant dependence with stimulant-induced psychotic disorder with delusions
	F15.950	Other stimulant use, unspecified with stimulant-induced psychotic disorder with delusions
	F16.150	Hallucinogen abuse with hallucinogen-induced psychotic disorder with delusions
	F16.250	Hallucinogen dependence with hallucinogen-induced psychotic disorder with delusions
	F16.950	Hallucinogen use, unspecified with hallucinogen-induced psychotic disorder with delusions
	F18.150	Inhalant abuse with inhalant-induced psychotic disorder with delusions
	F18.250	Inhalant dependence with inhalant-induced psychotic disorder with delusions
	F18.950	Inhalant use, unspecified with inhalant-induced psychotic disorder with delusions
	F19.150	Other psychoactive substance abuse with psychoactive substance-induced psychotic disorder with delusions
	F19.250	Other psychoactive substance dependence with psychoactive substance-induced psychotic disorder with delusions
	F19.950	Other psychoactive substance use, unspecified with psychoactive substance-induced psychotic disorder with delusions
	F11.151	Opioid abuse with opioid-induced psychotic disorder with hallucinations
	F11.251	Opioid dependence with opioid-induced psychotic disorder with hallucinations
	F11.951	Opioid use, unspecified with opioid-induced psychotic disorder with hallucinations
	F12.151	Cannabis abuse with psychotic disorder with hallucinations
	F12.251	Cannabis dependence with psychotic disorder with hallucinations
	F12.951	Cannabis use, unspecified with psychotic disorder with hallucinations
	F13.151	Sedative, hypnotic or anxiolytic abuse with sedative, hypnotic or anxiolytic-induced psychotic disorder with hallucination
	F13.251	Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced psychotic disorder with hallucination
	F13.951	Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced psychotic disorder with hallucination
	F14.151	Cocaine abuse with cocaine-induced psychotic disorder with hallucinations
	F14.251	Cocaine dependence with cocaine-induced psychotic disorder with hallucinations
	F14.951	Cocaine use, unspecified with cocaine-induced psychotic disorder with hallucinations

Approvable ICD-10 Group	Approvable ICD-10	Approvable ICD-10 Description
	F15.151	Other stimulant abuse with stimulant-induced psychotic disorder with hallucinations
	F15.251	Other stimulant dependence with stimulant-induced psychotic disorder with hallucinations
	F15.951	Other stimulant use, unspecified with stimulant-induced psychotic disorder with hallucinations
	F16.151	Hallucinogen abuse with hallucinogen-induced psychotic disorder with hallucinations
	F16.251	Hallucinogen dependence with hallucinogen-induced psychotic disorder with hallucinations
	F16.951	Hallucinogen use, unspecified with hallucinogen-induced psychotic disorder with hallucinations
	F18.151	Inhalant abuse with inhalant-induced psychotic disorder with hallucinations
	F18.251	Inhalant dependence with inhalant-induced psychotic disorder with hallucinations
	F18.951	Inhalant use, unspecified with inhalant-induced psychotic disorder with hallucinations
	F19.151	Other psychoactive substance abuse with psychoactive substance-induced psychotic disorder with hallucinations
	F19.251	Other psychoactive substance dependence with psychoactive substance-induced psychotic disorder with hallucinations
	F19.951	Other psychoactive substance use, unspecified with psychoactive substance-induced psychotic disorder with hallucinations
	F11.159	Opioid abuse with opioid-induced psychotic disorder, unspecified
	F91.3	Oppositional defiant disorder
F95		Tic disorder
G10		Huntington's disease

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 2 week** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Abilify MyCite ^{CC, QL} Caplyta ^{AE, CC, QL}	<ul style="list-style-type: none"> • Trial and failure of, or intolerance/contraindication to, ≥ 1 long-acting antipsychotic <p>Initial Criteria:</p> <ul style="list-style-type: none"> • Patient has a confirmed diagnosis of bipolar I or II disorder (bipolar depression) AND medication will be used as monotherapy or adjunctive therapy with lithium or valproate; AND • Trial and failure of ≥ 2 preferred antipsychotics. OR • Patient has a confirmed diagnosis of schizophrenia AND • Trial and failure of ≥ 2 preferred antipsychotics. OR • Patient has a diagnosis of moderate to severe major depressive disorder; AND • Medication will be used as adjunctive therapy with antidepressant; AND • Prescribed by, or in consultation with a psychiatrist or psychiatric mental health nurse practitioner (PMHNP); AND <ul style="list-style-type: none"> • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred antidepressants; AND • 6-week trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of aripiprazole or Vraylar. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Attestation or documentation (e.g., progress note) of disease improvement and/or stabilization <p>Age Limit: ≥ 18 years Quantity Limit: 1 per day</p>
Cobenfy ^{AE, CC, QL}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none"> • Diagnosis of schizophrenia; AND • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to one preferred agent; AND • Prescriber attests that liver enzymes and bilirubin were measured prior to initiation; AND • Patient meets the minimum age recommended by the package insert for the provided indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Prescriber attestation of clinically significant improvement or stabilization in clinical signs and symptoms. <p>Age Limit: 18 years of age or older Quantity Limit: 2 capsules per day</p>
Lybalvi ^{AE, CC, QL}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none"> • Patient has a diagnosis of schizophrenia OR bipolar I disorder; AND • If used for bipolar I disorder, will be used for either: <ul style="list-style-type: none"> ◦ acute treatment of manic or mixed episodes as monotherapy or as adjunct to lithium or valproate; OR ◦ maintenance monotherapy treatment; AND • Patient is NOT currently using opioids; AND • Patient is NOT undergoing acute opioid withdrawal; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Patient has a history of trial and therapeutic failure, allergy, contraindication or intolerance of ≥ 1 preferred second-generation (atypical) antipsychotic. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient must continue to meet the above criteria; AND • Patient must have disease improvement and/or stabilization <p>Age Limit: ≥ 18 years of age Quantity Limit: 30 tablets/30 days</p>
Nuplazid ^{AE, CC, QL}	<ul style="list-style-type: none"> • Diagnosis of Parkinson's Disease; AND • Trial of dose adjustment or withdrawal of anti-Parkinson's medications prior to treatment with this agent, (ex; anticholinergics, amantadine, dopamine agents, COMT inhibitors, selegiline) because these are known to cause hallucinations. <p>Age Limit: ≥ 18 years Quantity Limit: 2 tablets per day (60 tablets per 30 days)</p>
Opipza ^{CC, QL}	<ul style="list-style-type: none"> • Diagnosis of any of the following conditions: <ul style="list-style-type: none"> ◦ Dementias (ICD-10 Disease Groups F01, F02, F03, F06); ◦ Dissociative and conversion disorders (ICD-10 Disease Group F44); ◦ Episodic Mood Disorders (ICD-10 Disease Groups F30, F31, F39); ◦ Huntington's disease (ICD-10 Disease Group G10); ◦ Major depressive disorder (ICD-10 Disease Groups F32, F33); ◦ Oppositional defiant disorder (ICD-10 = F91.3); ◦ Pervasive developmental disorders (ICD-10 Disease Group F84); ◦ Schizoaffective disorder (F25.9); ◦ Schizophrenic Disorders (ICD-10 Disease Group F20; ICD-10 = F60.1); ◦ Tic disorder (ICD-10 Disease Group F95); ◦ Substance use disorders and related conditions; AND • Patient has had at least a 2-week trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to both aripiprazole tablet AND aripiprazole oral solution. <p>Quantity Limit:</p> <ul style="list-style-type: none"> • 2 mg film: 2 per day • 5 mg film: 1 per day • 10 mg film: 1 per day
olanzapine/fluoxetine capsule ^{CC, QL} Symbyax ^{CC, QL}	<ul style="list-style-type: none"> • Diagnosis of depressive episodes associated with bipolar disorder; AND • Trial and failure on one of the following: lithium, lamotrigine, bupropion, paroxetine. <p>OR</p> <ul style="list-style-type: none"> • Diagnosis of treatment-resistant depression; AND • Trial and failure on one of the following: selective serotonin reuptake inhibitor (SSRI), serotonin norepinephrine reuptake inhibitor (SNRI), new generation antidepressant, tricyclic antidepressant, monoamine oxidase inhibitor. <p>Quantity Limit: 1 capsule per day</p>

6. THERAPEUTIC DUPLICATION/MULTIPLE AGENTS CRITERIA

Prior authorization when ≥ 3 atypical antipsychotics are used may be approved under the following conditions:

- Approve for 1 year when it is continuation of current therapy and member is stable on 3 or more agents;
OR
- A maximum of two months to allow patients to taper to dual therapy (if one of the previous will be discontinued); **OR**
- Additional agents may be added to existing dual therapy after a two-week trial at the maximum tolerated dose of each agent.

CURRENT PDL STATUS

ANTIPSYCHOTICS: SECOND GENERATION (ATYPICAL)

Preferred Agents	Non-Preferred Agents
aripiprazole tablet ^{CC, QL}	Abilify MyCite starter kit ^{CC, QL}
asenapine tablet ^{CC, QL}	Abilify MyCite maintenance kit ^{CC, QL}
clozapine tablet ^{CC, QL}	Abilify tablet ^{QL}
lurasidone tablet ^{CC, QL}	aripiprazole ODT
olanzapine ODT ^{CC, QL}	aripiprazole solution
olanzapine tablet ^{CC, QL}	Caplyta capsule ^{AE, CC, QL}
quetiapine tablet ^{CC, QL}	clozapine ODT ^{QL}
quetiapine ER tablet ^{CC, QL}	Clozaril tablet ^{QL}
risperidone ODT ^{CC, QL}	Cobenfy capsule ^{AE, CC, QL}
risperidone solution ^{CC, QL}	Fanapt tablet dose pack ^{QL}
risperidone tablet ^{CC, QL}	Fanapt tablet ^{QL}
Vraylar capsule dose pack ^{AE, CC, QL}	Geodon capsule ^{QL}
Vraylar capsule ^{AE, CC, QL}	Invega ER tablet ^{QL}
ziprasidone capsule ^{CC, QL}	Latuda tablet ^{QL}
	Lybalvi tablet ^{AE, CC, QL}
	Nuplazid capsule ^{AE, CC, QL}
	Nuplazid tablet ^{AE, CC, QL}
	olanzapine/fluoxetine capsule ^{CC, QL}
	Opipza film ^{CC, QL}
	paliperidone ER tablet ^{QL}
	Rexulti tablet ^{QL}
	Risperdal solution ^{QL}
	Risperdal tablet ^{QL}
	Saphris SL tablet ^{QL}
	Secuado patch ^{QL}
	Seroquel tablet ^{QL}
	Seroquel XR tablet ^{QL}
	Symbyax capsule ^{CC, QL}
	Versacloz suspension ^{QL}
	Zyprexa tablet ^{QL}
	Zyprexa Zydis ODT ^{QL}

ANTIPSYCHOTICS: INJECTABLE

Preferred Agents	Non-Preferred Agents
Abilify Asimtufii ^{AE, CC, QL}	Erzofri syringe ^{AE, QL}
Abilify Maintena syringe ^{AE, CC, QL}	Haldol Decanoate ampule ^{AE, QL}
Abilify Maintena vial ^{AE, CC, QL}	risperidone ER vial ^{AE, QL}
Aristada syringe ^{AE, CC, QL}	Rykindo vial ^{AE, QL}
Aristada Initio syringe ^{AE, CC, QL}	ziprasidone mesylate vial ^{AE, QL}
fluphenazine decanoate vial ^{AE, CC, QL}	Zyprexa Relprevv vial ^{AE, QL}
Geodon vial ^{AE, CC, QL}	Zyprexa vial ^{AE, QL}
haloperidol decanoate ampule ^{AE, CC, QL}	
haloperidol decanoate vial ^{AE, CC, QL}	
haloperidol lactate syringe ^{AE, CC, QL}	
haloperidol lactate vial ^{AE, CC, QL}	
Invega Hafyera syringe ^{AE, CC, QL}	
Invega Sustenna syringe ^{AE, CC, QL}	
Invega Trinza syringe ^{AE, CC, QL}	
olanzapine vial ^{AE, CC, QL}	
Perseris suspension ^{AE, CC}	
Risperdal Consta vial ^{AE, CC, QL}	
Uzedy suspension ^{AE, CC, QL}	

CENTRAL NERVOUS SYSTEM: DOPAMINE RECEPTOR AGONISTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
pramipexole tablet	bromocriptine capsule
ropinirole tablet	bromocriptine tablet
	Mirapex ER tablet
	Neupro patch
	pramipexole ER tablet
	ropinirole ER tablet

CENTRAL NERVOUS SYSTEM: PARKINSON'S DISEASE (ANTIPARKINSON'S AGENTS)

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 1 week** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Nourianz ^{CC, QL}	<ul style="list-style-type: none">Diagnosis of Parkinson's disease (PD); ANDReceiving PD therapy with carbidopa/levodopa; ANDExperiencing "off" episodes with carbidopa/levodopa; ANDTrial and failure of at least 2 adjunctive therapies, such as:<ul style="list-style-type: none">Dopamine agonists (e.g., pramipexole, ropinirole);Monoamine oxidase-B inhibitors (e.g., selegiline)Catechol-O-methyltransferase inhibitors (e.g., entacapone); ANDNONE of the following contraindications:<ul style="list-style-type: none">Severe hepatic impairment (Child-Pugh C); OREnd-stage renal disease, including dialysis; ORPregnant; ORMajor psychiatric disorder. <p>Renewal Criteria</p> <ul style="list-style-type: none">Patient has clinically meaningful response of treatment (e.g., patient shows a reduction in time of "off" episodes) <p>Age Limit: ≥ 18 years Quantity Limit: 1 per day</p>
Ongentys ^{CC, QL}	<ul style="list-style-type: none">Diagnosis of Parkinson's disease (PD); ANDReceiving PD therapy with carbidopa/levodopa; ANDExperiencing "off" episodes with carbidopa/levodopa for at least 2 hours per day; ANDTrial and failure of at least 2 adjunctive therapies, such as:<ul style="list-style-type: none">Dopamine agonists (e.g., pramipexole, ropinirole);Monoamine oxidase-B inhibitors (e.g., selegiline)

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ Catechol-O-methyltransferase inhibitors (e.g., entacapone); AND • NONE of the following contraindications: <ul style="list-style-type: none"> ○ Severe hepatic impairment (Child-Pugh C); OR ○ End-stage renal disease (creatinine clearance \leq 15 ml/min); OR ○ Use with a monoamine oxidase-B (MAO-B) inhibitor. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient has clinically meaningful response of treatment (e.g., patient shows a reduction in time of “off” episodes) <p>Age Limit: \geq 18 years Quantity Limit: 1 per day</p>
Vyalev ^{AE, CC, QL}	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of Parkinson’s disease (PD); AND • Receiving PD therapy with carbidopa/levodopa; AND • Experiencing “off” episodes with carbidopa/levodopa for at least 2 hours per day; AND <ul style="list-style-type: none"> ○ Trial and failure of at least 2 adjunctive therapies, such as: Dopamine agonists (e.g., pramipexole, ropinirole) ○ Monoamine oxidase-B inhibitors (e.g., selegiline) ○ Catechol-O-methyltransferase inhibitors (e.g., entacapone); AND • Patient will not take within two weeks of a non-selective monoamine oxidase (MAO) inhibitor (e.g., phenelzine, isocarboxazid, tranylcypromine); AND • Patient meets the minimum age recommended by the package insert for the provided indication. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Patient has clinically meaningful response of treatment (e.g., patient shows a reduction in time of “off” episodes). <p>Age Limit: 18 years of age or older Quantity Limit: 2 vials (20 mL) per day</p>
Xadago ^{CC, QL}	<ul style="list-style-type: none"> • Diagnosis of Parkinson’s disease (PD); AND • Receiving PD therapy with carbidopa/levodopa; AND • Experiencing “off” episodes with carbidopa/levodopa; AND • Does not have severe hepatic impairment (Child-Pugh Score $>$ 9); AND • Not taking ANY the following medications: <ul style="list-style-type: none"> ○ Dextromethorphan; OR ○ MAOIs (e.g., or other drugs that are potent inhibitors of monoamine oxidase (e.g., linezolid); OR ○ Other serotonergic drugs (e.g., SNRIs, SSRIs, TCAs, St. John’s wort, cyclobenzaprine); OR ○ Opioids (e.g., meperidine, methadone, propoxyphene, tramadol); OR ○ Sympathomimetic medications (e.g., methylphenidate, amphetamine). <p>Age Limit: \geq 18 years Quantity Limit: 1 tablet per day</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
amantadine	Azilect tablet
benztropine tablet	carbidopa tablet
carbidopa/levodopa ta	carbidopa/levodopa ER capsule
carbidopa/levodopa ER tablet	Crexont
carbidopa/levodopa ODT	Dhivy
carbidopa/levodopa/entacapone	Duopa
entacapone	Gocovri
selegiline	Inbrija inhalation
trihexyphenidyl	Nourianz ^{AE, CC QL}
	Ongentys ^{AE, CC, QL}
	Osmolex ER
	Rytary
	Sinemet
	Xadago ^{AE, CC, QL}

CENTRAL NERVOUS SYSTEM: MOVEMENT DISORDERS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to PA Criteria	Criteria for Approval
Austedo ^{CC, AE, QL}	<p>Huntington's Chorea</p> <ul style="list-style-type: none"> • Patient is diagnosed with chorea related to Huntington's disease; AND • Patient does NOT have the following conditions: <ul style="list-style-type: none"> ○ Hepatic impairment or hepatic disease; AND ○ History of, or current, untreated or inadequately treated depression; OR ○ Suicidal ideation; AND • Patient has tried and failed tetrabenazine <p>Tardive Dyskinesia</p> <ul style="list-style-type: none"> • Diagnosis of tardive dyskinesia; AND • Prescribed by or in consultation with a neurologist or psychiatrist (or other mental health provider), provided patient has reasonable access; AND • Documentation or claims history of current or former chronic patient use of a dopamine antagonist (e.g., antipsychotic, metoclopramide, prochlorperazine, droperidol, promethazine, etc.) <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Patient continues to meet criteria defined for initial approval; AND • Documentation (e.g., progress note) of improvement in symptoms associated with respective condition (i.e., tardive dyskinesia or Huntington's chorea).
Ingrezza ^{AE, CC, QL}	<p>Huntington's Chorea</p> <ul style="list-style-type: none"> • Patient is diagnosed with chorea related to Huntington's disease; AND • Patient does NOT have the following conditions: <ul style="list-style-type: none"> ○ History of, or current, untreated or inadequately treated depression; OR ○ Suicidal ideation; AND • Patient is NOT concurrently using any of the following: <ul style="list-style-type: none"> ○ Monoamine oxidase (MAO) inhibitors (e.g., isocarboxazid, phenelzine, rasagiline, safinamide, selegiline, tranylcypromine, etc.) within 14 days; OR ○ Strong CYP3A4 inducers (e.g., carbamazepine, phenytoin, phenobarbital, rifampin and related agents, St. John's wort, etc.); OR ○ Another VMAT2 inhibitor (e.g., tetrabenazine, valbenazine); AND • Patient has tried and failed tetrabenazine. <p>Tardive Dyskinesia</p> <ul style="list-style-type: none"> • Diagnosis of tardive dyskinesia (TD); AND

Agent(s) Subject to PA Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Prescribed by or in consultation with a neurologist or psychiatrist (or other mental health provider), provided patient has reasonable access; AND • Documentation or claims history of current or former chronic patient use of a dopamine antagonist (e.g., antipsychotic, metoclopramide, prochlorperazine, droperidol, promethazine, etc.); AND • Patient is NOT concurrently using any of the following: <ul style="list-style-type: none"> ○ Monoamine oxidase (MAO) inhibitors (e.g., isocarboxazid, phenelzine, rasagiline, safinamide, selegiline, tranylcypromine, etc.) within 14 days; OR ○ Strong CYP3A4 inducers (e.g., carbamazepine, phenytoin, phenobarbital, rifampin and related agents, St. John's wort, etc.); OR ○ Another VMAT2 inhibitor (e.g., tetrabenazine, valbenazine). <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Patient continues to meet criteria defined for initial approval; AND • Documentation (e.g., progress note) of improvement in symptoms associated with respective condition (i.e., tardive dyskinesia or Huntington's chorea). <p>Age Limit: ≥ 18 years Quantity Limit: 1 per day</p>

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Not applicable.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to PA Criteria	Criteria for Approval
Austedo XR ^{CC, QL}	<p>Huntington's Chorea</p> <ul style="list-style-type: none"> • Patient is diagnosed with chorea related to Huntington's disease; AND • Patient does NOT have the following conditions: <ul style="list-style-type: none"> ○ Hepatic impairment or hepatic disease; AND ○ History of, or current, untreated or inadequately treated depression; OR ○ Suicidal ideation; AND • Patient has tried and failed tetrabenazine <p>Tardive Dyskinesia</p> <ul style="list-style-type: none"> • Diagnosis of tardive dyskinesia; AND • Prescribed by or in consultation with a neurologist or psychiatrist (or other mental health provider), provided patient has reasonable access; AND

Agent(s) Subject to PA Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Documentation or claims history of current or former chronic patient use of a dopamine antagonist (e.g., antipsychotic, metoclopramide, prochlorperazine, droperidol, promethazine, etc.) <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Patient continues to meet criteria defined for initial approval; AND Documentation (e.g., progress note) of improvement in symptoms associated with respective condition (i.e., tardive dyskinesia or Huntington's chorea).

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Austedo tablet ^{CC, AE, QL}	Austedo XR tablet ^{CC, QL}
Ingrezza capsule ^{AE, CC, QL}	Austedo XR tablet titration kit ^{CC}
Ingrezza sprinkle capsule ^{AE, CC, QL}	Xenazine
tetrabenazine tablet	

CENTRAL NERVOUS SYSTEM: ANTIDEPRESSANTS

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

5. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

6. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Auvelity ^{CC, AE, QL}	<p>Approval Duration: 1 year</p> <p>Initial Approval Criteria</p> <ul style="list-style-type: none">Diagnosis of major depressive disorder; ANDPatient is not pregnant, breastfeeding, or planning to become pregnant; ANDPatient as tried and failed, unless allergic, contraindicated or intolerant to 2 preferred agents in any sub-class; ORPatient has suicidal ideations with severe depression based on an objective measure [e.g., Patient Health Questionnaire-9 (PHQ-9), Hamilton Rating Scale for Depression (HDRS), Montgomery-Asberg Depression Rating Scale (MADRS), Clinically Useful Depression Outcome Scale (CUDOS), or Quick Inventory of Depressive Symptomatology – Self Report 16 Item (QIDS-SR16)]

Agent(s) Subject to Criteria	Criteria for Approval
	<p>Renewal Criteria</p> <ul style="list-style-type: none">Patient must continue to meet the above criteria; ANDPatient must have disease improvement and/or stabilization of disease; AND <p>Quantity Limit: 60 tablets per 30 days Age Limit: ≥ 18 years old</p>

Agent(s) Subject to CriteriaRaldesyl^{AE, CC}**Criteria for Approval****Approval Duration:** 6 months initial; 1 year renewal**Initial Approval Criteria**

- Diagnosis of major depressive disorder (MDD); **AND**
- Prescribed by, or in consultation with, psychiatrist, neurologist, or another qualified healthcare provider experienced in treating depression or related conditions; **AND**
- Patient meets at least ONE of the following criteria:
 - Unable to tolerate, swallow, or absorb oral tablet trazodone; **OR**
 - Tried and failed two preferred agents, one being trazodone IR; **AND**
- Patient must meet the minimum age recommended by the package insert for the provided indication.

Renewal Criteria

- Patient has experienced disease improvement and/or stabilization such as improvement in depressive symptoms, as assessed by the prescriber; **AND**
- Patient continues to require an oral solution.

Age Limit: ≥ 18 years of ageSpravato^{CC, QL}**Approval Duration:** 4 weeks initial; 1 year renewal (treatment resistant depression only)**Initial Approval Criteria****Treatment-resistant depression**

- Diagnosis of depression considered treatment resistant as evidenced by BOTH of the following:
 - Trial and failure (defined as < 50% reduction in symptom severity using any validated depression rating scale) of ≥ 2 antidepressants from different classes for a duration of ≥ 6 weeks each at generally accepted doses in the current depressive episode, unless contraindicated or clinically significant adverse effects are experienced; **AND**
 - Trial and failure of antidepressant augmentation therapy for a duration of ≥ 6 weeks in the current depressive episode with ≥ 1 of the following, unless contraindicated or clinically significant adverse effects are experienced:
 - An atypical antipsychotic; **OR**
 - Lithium; **OR**
 - An antidepressant from a different class; **AND**
- Prescribed by, or in consultation with, a psychiatrist or psychiatric mental health nurse practitioner (PMHNP); **AND**
- Prescriber has performed baseline depression assessment using any validated rating scale; **AND**
- If female of childbearing potential, NOT pregnant or planning to become pregnant, **AND**
- Prescriber attests that:
 - An accessible treatment center certified in the Spravato Risk Evaluation and Mitigation Strategies (REMS) program has been identified; **AND**
 - Dosing schedule has been reviewed with patient; **AND**
 - Patient understands and is committed to dosing schedule and requirements (e.g., office visits, transportation).

Major depressive disorder with acute suicidal ideations

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Diagnosis of MDD with acute suicidal ideation or behavior; AND • Used in conjunction with another oral antidepressant medication (not to be used as monotherapy); AND • Prescribed by, or in consultation with, a psychiatrist or psychiatric mental health nurse practitioner (PMHNP); AND • Prescriber has performed baseline depression assessment using any validated rating scale; AND • If female of childbearing potential, NOT pregnant or planning to become pregnant, AND • Prescriber attests that: <ul style="list-style-type: none"> ○ An accessible treatment center certified in the Spravato Risk Evaluation and Mitigation Strategies (REMS) program has been identified; AND ○ Dosing schedule has been reviewed with patient; AND ○ Patient understands and is committed to dosing schedule and requirements (e.g., office visits, transportation). <p>Renewal Criteria (not applicable when used for acute suicidal ideation)</p> <p>Treatment-resistant depression</p> <ul style="list-style-type: none"> • Continue to meet initial approval criteria for treatment resistant depression; AND • Prescriber attestation that patient has been compliant with doses/appointments; AND • Prescriber provides documentation of disease improvement or stabilization as evidenced by improvement on a validated depression rating scale. <p>Age Limit: ≥ 18 years old Quantity Limit: 1 kit (56 or 84 mg) per week; overrides allowed for twice weekly)</p>
Zurzuva ^{CC, QL}	<p>Approval Duration: 6 months with limit of 2 courses of treatment (28 days)</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of postpartum depression (PPD) in adults • Within one year of giving birth. <p>Quantity Limit: maximum 14 day supply per fill, maximum 2 fills per 180 days</p>

CURRENT PDL STATUS

ANTIDEPRESSANTS: OTHER

Preferred Agents	Non-Preferred Agents
bupropion tablet	Auvelity tablet ^{CC, AE, QL}
bupropion SR tablet	bupropion XL 450 mg tablet
bupropion XL 150 mg, 300 mg tablet	Forfivo XL tablet
mirtazapine ODT	nefazodone tablet
mirtazapine tablet	Raldesy solution ^{AE, CC}
trazodone tablet	Remeron Soltab
	Remeron tablet
	Spravato spray ^{CC, AE, QL}
	Trintellix tablet

Preferred Agents	Non-Preferred Agents
	Viibryd tablet dose pack
	Viibryd tablet
	Vilazodone tablet
	Wellbutrin SR tablet
	Wellbutrin XL tablet
	Zurzuva capsule ^{CC, QL}

ANTIDEPRESSANTS: SNRIS

Preferred Agents	Non-Preferred Agents
desvenlafaxine succinate ER tablet	desvenlafaxine ER base tablet
venlafaxine tablet	Effexor XR capsule
venlafaxine ER capsule	Fetzima ER capsule
venlafaxine ER tablet	Fetzima ER capsule dose pack
	Pristiq ER tablet
	venlafaxine besylate ER tablet

ANTIDEPRESSANTS: SSRIS

Preferred Agents	Non-Preferred Agents
citalopram 10 mg/5 mL solution	Celexa tablet
citalopram tablet	citalopram capsule
escitalopram tablet	citalopram 20 mg/10 mL solution
fluoxetine capsule	escitalopram capsules, solution
fluoxetine solution	fluoxetine 90 mg DR capsule ^{QL}
paroxetine tablet	fluoxetine tablet
sertraline oral concentrate	fluvoxamine ER capsule
sertraline tablet	fluvoxamine tablet
	Lexapro tablet
	paroxetine CR tablet
	paroxetine ER tablet
	paroxetine mesylate capsule
	paroxetine suspension
	Paxil CR tablet
	Paxil suspension
	Paxil tablet
	Pexeva tablet
	Prozac capsule
	sertraline capsule
	Zoloft oral concentrate
	Zoloft tablet

ANTIDEPRESSANTS: TRICYCLICS

Preferred Agents	Non-Preferred Agents
amitriptyline tablet	amoxapine tablet
clomipramine capsule	Anafranil capsule
doxepin capsule	desipramine tablet
doxepin oral concentrate	imipramine pamoate capsule
imipramine tablet	Norpramin tablet
nortriptyline capsule	nortriptyline solution
	Pamelor capsule
	protriptyline tablet
	trimipramine capsule

ANTIDEPRESSANTS: MAOIs

Preferred Agents	Non-Preferred Agents
	Emsam patch
	Marplan tablet
	Nardil tablet
	phenelzine tablet
	tranylcypromine tablet

CENTRAL NERVOUS SYSTEM: ANTI-MIGRAINE AGENTS, TRIPTANS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Zembrace SymTouch ^{CC, QL}	<ul style="list-style-type: none">Trial and failure of a corresponding generic sumatriptan formulation (e.g., nasal spray, injection)

6. QUANTITY LIMIT CRITERIA

One-time approval when all of the following circumstances are true

- NOT using triptans in combination with an MAOI (e.g., Parnate, Marplan, or Nardil); **AND**
- Patient must **NOT** have a history of ischemic heart disease; **AND**
- Prescriber has counseled the member about the negatives (e.g., causes additional migraines) of daily use and/or overuse of triptans and will attempt to taper the quantity of triptan medication used monthly; **AND**
- Current use of any oral or injectable prophylactic agent, such as (though not limited to):
 - Antiepileptic drugs (AEDs): divalproex sodium, sodium valproate, topiramate
 - Beta Blockers: metoprolol, propranolol, timolol, atenolol, nadolol
 - Antidepressants: amitriptyline, venlafaxine
 - NSAIDs: fenoprofen, ibuprofen, ketoprofen, naproxen

- CGRP inhibitor: Ajoovy, Emgality 120 mg/mL
- Botulinum toxin: Botox

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
rizatriptan ODT ^{QL}	almotriptan tablet ^{QL}
rizatriptan tablet ^{QL}	eletriptan tablet ^{QL}
sumatriptan spray ^{QL}	Frova tablet ^{QL}
sumatriptan tablet ^{QL}	frovatriptan tablet ^{QL}
sumatriptan vial ^{QL}	Imitrex cartridge ^{QL}
	Imitrex pen ^{QL}
	Imitrex tablet ^{QL}
	Maxalt-MLT ODT ^{QL}
	Maxalt tablet ^{QL}
	naratriptan tablet ^{QL}
	Onzetra Xsail ^{QL}
	Relpax tablet ^{QL}
	sumatriptan cartridge ^{QL}
	sumatriptan injector ^{QL}
	sumatriptan/naproxen tablet ^{QL}
	Symbravo ^{QL}
	Tosymra spray
	Zembrace SymTouch ^{CC, QL}
	zolmitriptan ODT ^{QL}
	zolmitriptan spray ^{QL}
	zolmitriptan tablet ^{QL}
	Zomig spray ^{QL}
	Zomig tablet ^{QL}

CENTRAL NERVOUS SYSTEM: ANTI-MIGRAINE AGENTS, CGRP INHIBITORS

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria		Criteria for Approval							
Aimovig ^{AE, CC, QL} Emgality 120 mg/mL ^{AE, CC, QL}		<p>Approval Duration: 3 months initial; 1 year renewal</p> <ul style="list-style-type: none">• Diagnosis of migraine with or without aura; AND• Patient has tried and failed a ≥ 1-month trial (at maximally tolerated doses) of two medications listed below from the 2012 American Academy of Neurology/American Headache Society guidelines. At least ONE medication must be level A or B recommendation, unless ALL are contraindicated: <table><tr><th>Level A</th><th>Level B</th><th>Level C</th></tr><tr><td><ul style="list-style-type: none">• divalproex sodium• sodium valproate• topiramate• metoprolol• propranolol• timolol</td><td><ul style="list-style-type: none">• amitriptyline• venlafaxine• atenolol• nadolol</td><td><ul style="list-style-type: none">• clonidine• guanfacine• lisinopril• candesartan• carbamazepine• cyproheptadine• nebivolol• pindolol</td></tr></table> <p>Renewal Criteria</p> <ul style="list-style-type: none">• Patient has an overall improvement in function with therapy. <p>Age Limit: ≥ 18 years</p>		Level A	Level B	Level C	<ul style="list-style-type: none">• divalproex sodium• sodium valproate• topiramate• metoprolol• propranolol• timolol	<ul style="list-style-type: none">• amitriptyline• venlafaxine• atenolol• nadolol	<ul style="list-style-type: none">• clonidine• guanfacine• lisinopril• candesartan• carbamazepine• cyproheptadine• nebivolol• pindolol
Level A	Level B	Level C							
<ul style="list-style-type: none">• divalproex sodium• sodium valproate• topiramate• metoprolol• propranolol• timolol	<ul style="list-style-type: none">• amitriptyline• venlafaxine• atenolol• nadolol	<ul style="list-style-type: none">• clonidine• guanfacine• lisinopril• candesartan• carbamazepine• cyproheptadine• nebivolol• pindolol							
Ajovy ^{AE, CC, QL}		<p>Approval Duration: 3 months initial; 1 year renewal</p> <ul style="list-style-type: none">• Patient is 6 to 17 years of age or older and weighs 45 kg or more:<ul style="list-style-type: none">○ Diagnosis of episodic migraine; AND○ If patient is 12 years or older, patient has a 2-month trial and failure of topiramate, unless contraindicated or clinically significant adverse effects are experienced; OR• Patient is 18 years of age or older:<ul style="list-style-type: none">○ Diagnosis of migraine with or without aura; AND○ Patient has tried and failed a ≥ 1-month trial (at maximally tolerated doses) of two medications listed below from the 2012 American Academy of Neurology/American Headache Society guidelines. At least ONE medication must be level A or B recommendation, unless ALL are contraindicated: <table><tr><th>Level A</th><th>Level B</th><th>Level C</th></tr><tr><td><ul style="list-style-type: none">• divalproex sodium• sodium valproate• topiramate• metoprolol• propranolol</td><td><ul style="list-style-type: none">• amitriptyline• venlafaxine• atenolol• nadolol</td><td><ul style="list-style-type: none">• clonidine• guanfacine• lisinopril• candesartan• carbamazepine• cyproheptadine</td></tr></table>		Level A	Level B	Level C	<ul style="list-style-type: none">• divalproex sodium• sodium valproate• topiramate• metoprolol• propranolol	<ul style="list-style-type: none">• amitriptyline• venlafaxine• atenolol• nadolol	<ul style="list-style-type: none">• clonidine• guanfacine• lisinopril• candesartan• carbamazepine• cyproheptadine
Level A	Level B	Level C							
<ul style="list-style-type: none">• divalproex sodium• sodium valproate• topiramate• metoprolol• propranolol	<ul style="list-style-type: none">• amitriptyline• venlafaxine• atenolol• nadolol	<ul style="list-style-type: none">• clonidine• guanfacine• lisinopril• candesartan• carbamazepine• cyproheptadine							

Agent(s) Subject to Criteria	Criteria for Approval		
	<ul style="list-style-type: none">timolol		<ul style="list-style-type: none">nebivololpindolol
	<p>Renewal Criteria</p> <ul style="list-style-type: none">Patient has an overall improvement in function with therapy. <p>Age Limit: ≥ 6 years</p>		
Nurtec ODT AE, CC, QL	<p>Acute treatment of migraine</p> <ul style="list-style-type: none">Diagnosis of migraine, with or without aura; ANDTrial and failure, or contraindication to, 2 triptans. <p>Renewal Criteria:</p> <ul style="list-style-type: none">Attestation or documentation of resolution in headache pain or reduction in headache severity, as assessed by prescriber. <p>Age Limit: > 18 years Quantity Limit: 18 tablets per 30 days</p>		
Qulipta AE, CC, QL Nurtec ODT AE, CC, QL (for prevention of episodic migraine)	<p>Approval Duration: 3 months initial; 1 year renewal</p> <p>Initial Approval Criteria</p> <p><i>Episodic migraine</i></p> <ul style="list-style-type: none">Patient has diagnosis of migraine with or without aura based on International Classification of Headache Disorders (ICHD-III) diagnostic criteria; ANDPatient has tried and failed a ≥ 1-month trial of a preferred injectable CGRP agent (i.e., Aimovig, Ajovy, Emgality 120 mg/mL); ANDPatient has experienced ≥ 4 migraine days per month; ANDPatient has not experienced > 15 headache days per month during the prior 6 months; ANDMedication overuse has been ruled out. <p><i>Chronic Migraine</i></p> <ul style="list-style-type: none">Patient has diagnosis of migraine with or without aura based on International Classification of Headache Disorders (ICHD-III) diagnostic criteria; ANDPatient has tried and failed a ≥ 1-month trial of a preferred injectable CGRP agent (i.e. Aimovig, Ajovy, Emgality 120 mg/mL); ANDPatient has experienced ≥ 8 migraine days per month during the last 3 months; ANDPatient has experienced ≥ 15 headache days per month during the prior 3 months; ANDMedication overuse has been ruled out. <p>Renewal Criteria</p> <ul style="list-style-type: none">Patient demonstrated significant decrease in the number, frequency, and/or intensity of headaches <p>Age Limit: ≥ 18 years Quantity Limit:</p> <p><u>Qulipta</u></p> <ul style="list-style-type: none">30mg tablet and 60mg tablet: 30 tablets per 30 days10mg tablet: 60 tablets per 30 days <p>Nurtec ODT</p>		

Agent(s) Subject to Criteria	Criteria for Approval
Ubrelvy ^{AE, CC, QL}	<ul style="list-style-type: none"> 18 tablets per 30 days <ul style="list-style-type: none"> Diagnosis of migraine, with or without aura; AND Trial and failure, or contraindication to, 2 triptans (e.g., sumatriptan). <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Attestation or documentation of resolution in headache pain or reduction in headache severity, as assessed by prescriber. <p>Age Limit: > 18 years Quantity Limit:</p> <ul style="list-style-type: none"> 50 mg tablet: 10 tablets (1 package) per 30 days 100 mg tablet: 16 tablets (1 package) per 30 days One-time fill of 20 tablets (2 packages) per 30 days allowed with prior authorization: current use of any oral or injectable prophylactic agent listed below.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers if available and covered) of the corresponding generic.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Emgality 100 mg/mL ^{CC, AE, QL}	<p>Approval Duration: 3 months initial; 1 year renewal</p> <ul style="list-style-type: none"> Diagnosis of episodic cluster headache as evidenced by a history of ≥ 2 cluster periods lasting from ≥ 7 days to ≤ 1 year each and separated by ≥ 3 months; AND Prescribed by, or in consultation with, a neurologist or headache/pain specialist; AND NOT to be used in combination with any other injectable CGRP (e.g., Ajovy) or botulinum toxin (e.g., Botox); <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Patient has an overall improvement in function with therapy compared with previous cluster periods; AND Member has not received more than 12 months of consecutive treatment OR it has been at least 3 months since last treatment with Emgality 100 mg/ML. <p>Age Limit: ≥ 18 years Quantity Limit: 300 mg (3 mL) per 30 days</p>
Reyvow ^{CC, AE, QL}	<ul style="list-style-type: none"> Diagnosis of migraine, with or without aura; AND NOT have severe hepatic impairment (Child-Pugh C); AND Trial and failure of at least one of the following: NSAID, non-opioid

Agent(s) Subject to Criteria	Criteria for Approval
	<p>analgesic, acetaminophen, OR caffeinated analgesic combination; AND</p> <ul style="list-style-type: none"> • Trial and failure, or contraindication to, ≥ 2 triptans; AND • Prescriber attests patient has been educated about need to refrain from driving or operating machinery for ≥ 8 hours after dose. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Attestation or documentation of resolution in headache pain or reduction in headache severity, as assessed by prescriber. <p>Age Limit: ≥ 18 years</p> <p>Quantity Limit: 8 tablets (1 package) per 30 days – no exceptions</p>
Zavzpret ^{CC, AE, QL}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none"> • Patient has a diagnosis of migraine with or without aura; AND • Prescriber attestation will NOT be used for preventive treatment of migraine or for chronic migraine; AND • Patient must have tried and failed or have a contraindication or intolerance to 2 triptans; AND • Patient must have tried and failed or have a contraindication or intolerance to 1 preferred CGRP antagonist <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient must continue to meet the above criteria; AND • Patient must demonstrate symptom improvement (e.g., resolution in headache pain or reduction in headache severity), as assessed by the prescriber. <p>Quantity Limit: 8 nasal spray devices per 30 days</p> <p>Age Limit: ≥ 18 years old</p>

6. QUANTITY LIMIT CRITERIA FOR NURTEC ODT AND UBRELVY

Current use of any oral or injectable prophylactic agent, such as (though not limited to):

- Antiepileptic drugs (AEDs): divalproex sodium, sodium valproate, topiramate
- Beta Blockers: metoprolol, propranolol, timolol, atenolol, nadolol
- Antidepressants: amitriptyline, venlafaxine
- NSAIDs: fenoprofen, ibuprofen, ketoprofen, naproxen
- CGRP inhibitor: Ajovy, Emgality 120 mg/mL
- Botulinum toxin: Botox

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Aimovig ^{CC, AE, QL}	Emgality 100 mg/mL & 100 mg/mL ^{CC, AE, QL}
Ajovy ^{CC, AE, QL}	Reyvow ^{CC, AE, QL}
Emgality 120 mg/mL ^{CC, AE, QL}	Zavzpret ^{CC, AE, QL}
Nurtec ODT ^{CC, AE, QL}	
Qulipta ^{CC, AE, QL}	
Ubrelyvy ^{CC, AE, QL}	

CENTRAL NERVOUS SYSTEM: STIMULANTS AND RELATED AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to PA Criteria	Criteria for Approval
Adderall XR ^{CC, QL} atomoxetine ^{CC, QL} clonidine ER ^{CC, QL} Concerta ^{CC, QL} dexamethylphenidate ^{CC, QL} dexamethylphenidate ER ^{CC, QL} dextroamphetamine sulfate tablet 5mg, 10mg, 15mg ^{CC, QL} dextroamphetamine/amphetamine ^{CC, QL} dextroamphetamine/amphetamine ER ^{CC, QL} guanfacine ER ^{CC, QL} Methylin solution ^{CC, QL} methylphenidate solution ^{CC, QL} methylphenidate ER tablets 10 mg & 20 mg ^{CC, QL} (generic Metadate ER) methylphenidate ER tablet 18 mg, 27 mg, 36 mg, 54 mg ^{CC, QL} (generic Concerta) methylphenidate tablets ^{CC, QL} mixed amphetamine salts tablets ^{CC, QL}	<ul style="list-style-type: none"> Diagnosis of: <ul style="list-style-type: none"> Add/ADHD (ICD-10 Disease Group F90); OR Narcolepsy (ICD-10 Codes G47.419, G47.411, G47.421, G47.429); OR Sleep apnea (ICD-10 Code G47.30); OR Circadian rhythm (shift work) sleep disorder (ICD-10 Codes G47.20, G47.21, G47.22, G47.23, G47.24, G47.25, G47.26, G47.27, G47.29) Idiopathic hypersomnia (ICD-10 Codes G47.11, G47.12)
Qelbree ^{CC, QL}	<ul style="list-style-type: none"> Diagnosis of ADHD (ICD-10 Disease Group F90); AND Trial and failure of at least one other preferred agent.
Vyvanse capsules, chewable tablets ^{CC, QL}	<ul style="list-style-type: none"> Diagnosis of: <ul style="list-style-type: none"> ADD/ADHD (ICD-10 Disease Group F90); OR Narcolepsy (ICD-10 Codes G47.419, G47.411, G47.421, G47.429); OR Sleep apnea (ICD-10 Code G47.30); OR Circadian rhythm (shift work) sleep disorder (ICD-10 Codes G47.20, G47.21, G47.22, G47.23, G47.24, G47.25, G47.26, G47.27, G47.29); OR Patient has a diagnosis of moderate to severe binge eating disorder based on DSM V diagnostic criteria; AND <ul style="list-style-type: none"> Prescriber attests or provides documentation that a comprehensive evaluation has been performed, including, physical exam and any necessary labs; AND The patient been counseled on the benefits of cognitive behavioral therapy (CBT) and referred if appropriate. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Documentation of disease response [e.g., reduction in the number of binge-eating days per week, improvement of the

Agent(s) Subject to PA Criteria	Criteria for Approval
	Yale-Brown Obsessive Compulsive Scale Modified for Binge Eating (Y-BOCS-BE) total score]

Approvable ICD-10 Group	Approvable ICD-10	Approvable ICD-10 Description
F90		Attention-deficit hyperactivity disorders
	G47.411	Narcolepsy with cataplexy
	G47.419	Narcolepsy without cataplexy
	G47.421	Narcolepsy in conditions classified elsewhere with cataplexy
	G47.429	Narcolepsy in conditions classified elsewhere without cataplexy
	G47.30	Sleep apnea, unspecified
	G47.20	Circadian rhythm sleep disorder, unspecified type
	G47.21	Circadian rhythm sleep disorder, delayed sleep phase type
	G47.22	Circadian rhythm sleep disorder, advanced sleep phase type
	G47.23	Circadian rhythm sleep disorder, irregular sleep wake type
	G47.24	Circadian rhythm sleep disorder, free running type
	G47.25	Circadian rhythm sleep disorder, jet lag type
	G47.26	Circadian rhythm sleep disorder, shift work type
	G47.27	Circadian rhythm sleep disorder in conditions classified elsewhere
	G47.29	Other circadian rhythm sleep disorder
	G47.11	Idiopathic hypersomnia with long sleep time
	G47.12	Idiopathic hypersomnia without long sleep time

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent, unless otherwise specified. Preferred with PA (PDP) criteria must be met.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Adzenys XR-ODT ^{AE, CC, QL} amphetamine ER ODT ^{AE, CC, QL}	<ul style="list-style-type: none"> • Diagnosis of: <ul style="list-style-type: none"> ○ Attention Deficit Disorder (ADD)/ Attention-deficit hyperactivity disorder (ADHD) (ICD-10 Disease Group F90); OR ○ Narcolepsy (ICD-10 Codes G47.419, G47.411, G47.421, G47.429); OR ○ Sleep apnea (ICD-10 Code G47.30); OR ○ Circadian rhythm (shift work) sleep disorder (ICD-10 Codes G47.20, G47.21, G47.22, G47.23, G47.24, G47.25, G47.26, G47.27, G47.29); OR ○ Idiopathic hypersomnia (ICD-10 Codes G47.11, G47.12); AND • One of the following: <ul style="list-style-type: none"> ○ Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications), or intolerance to TWO preferred agents; OR ○ Patient has a swallowing disorder and cannot be given tablets or capsules. <p>Age Limit: ≥ 6 years Quantity Limit: 1 per day</p>
Xelstrym ^{CC, QL} Daytrana patch ^{CC, QL} methylphenidate patch ^{CC, QL}	<ul style="list-style-type: none"> • Diagnosis of: <ul style="list-style-type: none"> ○ Attention Deficit Disorder (ADD)/ Attention-deficit hyperactivity disorder (ADHD) (ICD-10 Disease Group F90); OR ○ Narcolepsy (ICD-10 Codes G47.419, G47.411, G47.421, G47.429); OR ○ Sleep apnea (ICD-10 Code G47.30); OR ○ Circadian rhythm (shift work) sleep disorder (ICD-10 Codes G47.20, G47.21, G47.22, G47.23, G47.24, G47.25, G47.26, G47.27, G47.29); OR ○ Idiopathic hypersomnia (ICD-10 Codes G47.11, G47.12); AND • One of the following: <ul style="list-style-type: none"> ○ Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications), or intolerance to ONE preferred agents; OR ○ Inability to swallow or tolerate medications. <p>Quantity Limit: 1 per day</p>
methylphenidate chewable tablet ^{CC, QL}	<ul style="list-style-type: none"> • Diagnosis of: <ul style="list-style-type: none"> ▪ Attention Deficit Disorder (ADD)/ Attention-deficit hyperactivity disorder (ADHD) (ICD-10 Disease Group F90); OR ▪ Narcolepsy (ICD-10 Codes G47.419, G47.411, G47.421, G47.429); OR ▪ Sleep apnea (ICD-10 Code G47.30); OR ▪ Circadian rhythm (shift work) sleep disorder (ICD-10 Codes G47.20, G47.21, G47.22, G47.23, G47.24, G47.25, G47.26, G47.27, G47.29); OR ▪ Idiopathic hypersomnia (ICD-10 Codes G47.11, G47.12); AND • One of the following: <ul style="list-style-type: none"> ▪ Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications), or intolerance to ONE preferred agents; OR

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Inability to swallow tablets or use the oral solution. <p>Quantity Limit: 3 per day</p>
ProCentra solution ^{CC, QL} dextroamphetamine solution ^{CC, QL}	<ul style="list-style-type: none"> Diagnosis of: <ul style="list-style-type: none"> Attention Deficit Disorder (ADD)/ Attention-deficit hyperactivity disorder (ADHD) (ICD-10 Disease Group F90); OR Narcolepsy (ICD-10 Codes G47.419, G47.411, G47.421, G47.429); OR Sleep apnea (ICD-10 Code G47.30); OR Circadian rhythm (shift work) sleep disorder (ICD-10 Codes G47.20, G47.21, G47.22, G47.23, G47.24, G47.25, G47.26, G47.27, G47.29); OR Idiopathic hypersomnia (ICD-10 Codes G47.11, G47.12); AND One of the following: <ul style="list-style-type: none"> Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications), or intolerance to ONE preferred agents; OR Inability to swallow tablets or capsules whole Quantity Limit: 60mL per day
lisdexamfetamine capsule ^{CC, QL} lisdexamfetamine chewable tablet ^{CC, QL}	<ul style="list-style-type: none"> Drug-specific PDP criteria; AND NPD criteria; AND GMN criteria <p>Quantity Limit: 1 per day</p>
Intuniv ER tablet ^{CC, QL}	<ul style="list-style-type: none"> Diagnosis of: <ul style="list-style-type: none"> Attention Deficit Disorder (ADD)/ Attention-deficit hyperactivity disorder (ADHD) (ICD-10 Disease Group F90); OR Narcolepsy (ICD-10 Codes G47.419, G47.411, G47.421, G47.429); OR Sleep apnea (ICD-10 Code G47.30); OR Circadian rhythm (shift work) sleep disorder (ICD-10 Codes G47.20, G47.21, G47.22, G47.23, G47.24, G47.25, G47.26, G47.27, G47.29); OR Idiopathic hypersomnia (ICD-10 Codes G47.11, G47.12); AND BMN criteria <p>Quantity Limit: 1 per day</p>

6. THERAPEUTIC DUPLICATION

Patients are limited to **one** long-acting and **one** short-acting CNS stimulant at a time within the quantity/dosing limits. Overrides may be approved:

- For the date of service when patients are switching from one agent to another.
- For 1 year when the member requires two different strengths of one medication because the prescribed dosage cannot be achieved otherwise (e.g., Concerta 18 mg and Concerta 27 mg taken together once daily).
- A clinical pharmacist may approve with clinical justification when:
 - Prescriber requests more than one long-acting or more than one short-acting;
 - Prescriber is splitting the dosage of a long-acting agent for the patient (e.g., Concerta 18 mg AM and Concerta 27 mg at 4:00 PM).

- **NOTES:**

- Intuniv (guanfacine ER) and Kapvay (clonidine ER) are exempt from this edit.
- Strattera (atomoxetine) is included in this edit.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Adderall XR capsule ^{CC, QL}	Adderall tablet ^{QL}
atomoxetine capsule ^{CC, QL}	Adzenys XR-ODT ^{AE, CC, QL}
clonidine ER tablet ^{CC, QL}	amphetamine ER ODT ^{AE, CC, QL}
Concerta tablet ^{CC, QL}	amphetamine sulfate tablet ^{QL}
dexmethylphenidate ER capsule ^{CC, QL}	Aptensio XR capsule ^{QL}
dexmethylphenidate tablet ^{CC, QL}	Azstarys capsule ^{QL}
dextroamphetamine sulfate tablet 5 mg, 10 mg, 15 mg ^{CC, QL}	Cotempla XR-ODT tablet ^{AE, QL}
dextroamphetamine/amphetamine ER capsule ^{CC, QL}	Daytrana patch ^{CC, QL}
dextroamphetamine/amphetamine tablet ^{CC, QL}	Desoxyn tablet ^{QL}
guanfacine ER tablet ^{CC, QL}	Dexedrine ER capsule ^{QL}
Methylin solution ^{CC, QL}	dextroamphetamine ER capsule ^{QL}
methylphenidate solution ^{CC, QL}	dextroamphetamine solution ^{CC, QL}
methylphenidate ER tablet 10 mg, 20 mg ^{CC, QL} (generic Metadate ER)	dextroamphetamine sulfate tablet 2.5 mg, 7.5 mg, 20 mg, 30 mg ^{QL}
methylphenidate ER tablet 18 mg, 27 mg, 36 mg, 54 mg ^{CC, QL} (generic Concerta)	Dyanavel XR suspension ^{AE, QL}
methylphenidate tablet ^{CC, QL}	Dyanavel XR tablet ^{AE, QL}
Qelbree ER capsule ^{CC, QL}	Evekeo ODT ^{QL}
Vyvanse capsule ^{CC, QL}	Evekeo tablet ^{QL}
Vyvanse chewable tablet ^{CC, QL}	Focalin tablet ^{QL}
	Focalin XR capsule ^{QL}
	Intuniv ER tablet ^{CC, QL}
	Jornay PM capsule ^{AE, QL}
	lisdexamfetamine capsule ^{CC, QL}
	lisdexamfetamine chewable tablet ^{CC, QL}
	methamphetamine tablet ^{QL}
	methylphenidate CD capsule ^{QL}
	methylphenidate ER capsule ^{QL}
	methylphenidate ER tablet 63 mg, 72 mg tablet ^{QL} (generic for Relexxii)
	methylphenidate ER sprinkle capsule ^{QL}
	methylphenidate LA capsule ^{QL}
	methylphenidate ER OROS ^{QL}
	methylphenidate chewable tablet ^{CC, QL}
	methylphenidate patch ^{CC, QL}
	Mydayis ER capsule ^{AE, QL}
	Onyda XR suspension ^{AE, QL}
	ProCentra solution ^{CC, QL}
	QuilliChew ER tablet ^{AE, QL}
	Quillivant XR suspension ^{QL}
	Relexxii tablet ^{QL}
	Ritalin LA capsule ^{QL}
	Ritalin tablet ^{QL}
	Strattera capsule ^{QL}
	Xelstryl patch ^{CC, QL}
	Zenzedi tablet ^{QL}

CENTRAL NERVOUS SYSTEM: NARCOLEPSY AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to PA Criteria	Criteria for Approval
armodafinil tablet ^{CC, QL} Provigil ^{CC, QL}	<ul style="list-style-type: none"> Diagnosis of: <ul style="list-style-type: none"> Narcolepsy (ICD-10 Codes G47.419, G47.411, G47.421, G47.429); OR Sleep apnea (ICD-10 Code G47.30); OR Shift work sleep disorder (ICD-10 Codes G47.20, G47.21, G47.22, G47.23, G47.24, G47.25, G47.26, G47.27, G47.29). Idiopathic hypersomnia (ICD-10 Codes G47.11, G47.12)

Approvable ICD-10 Group	Approvable ICD-10	Approvable ICD-10 Description
	G47.411	Narcolepsy with cataplexy
	G47.419	Narcolepsy without cataplexy
	G47.421	Narcolepsy in conditions classified elsewhere with cataplexy
	G47.429	Narcolepsy in conditions classified elsewhere without cataplexy
	G47.30	Sleep apnea, unspecified
	G47.20	Circadian rhythm sleep disorder, unspecified type
	G47.21	Circadian rhythm sleep disorder, delayed sleep phase type
	G47.22	Circadian rhythm sleep disorder, advanced sleep phase type
	G47.23	Circadian rhythm sleep disorder, irregular sleep wake type
	G47.24	Circadian rhythm sleep disorder, free running type
	G47.25	Circadian rhythm sleep disorder, jet lag type
	G47.26	Circadian rhythm sleep disorder, shift work type
	G47.27	Circadian rhythm sleep disorder in conditions classified elsewhere
	G47.29	Other circadian rhythm sleep disorder
	G47.11	Idiopathic hypersomnia with long sleep time
	G47.12	Idiopathic hypersomnia without long sleep time

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent, unless otherwise specified. Preferred with PA (PDP) criteria must be met.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Sunosi ^{CC, QL}	<ul style="list-style-type: none">• Prescriber is a neurologist, sleep medicine, or other specialist in the treatment of narcolepsy; AND• Prescriber attestation or documentation that member's blood pressure is adequately controlled ($\leq 140/90$ mmHg); AND• Trial and failure/intolerance of, or contraindication to, ≥ 1 narcolepsy agent (e.g., modafinil); AND• Diagnosis of excessive daytime sleepiness associated with narcolepsy; AND<ul style="list-style-type: none">▪ Trial and failure of ≥ 1 stimulant (e.g., amphetamine); OR• Diagnosis of excessive daytime sleepiness associated with obstructive sleep apnea (OSA); AND<ul style="list-style-type: none">▪ Member is using constant positive airway pressure (CPAP).
Wakix ^{CC, QL}	<ul style="list-style-type: none">• Prescriber is a neurologist, sleep medicine, or other specialist in the treatment of narcolepsy; AND• Diagnosis of cataplexy and/or excessive daytime sleepiness associated with narcolepsy; AND• Documentation of a multiple sleep latency test (MSLT) confirming narcolepsy; AND• Trial and failure/intolerance of, contraindication to, ≥ 1 narcolepsy agent (e.g., modafinil); trial can be waived if member has a history of substance abuse; AND• Trial and failure/intolerance of, contraindication to, of ≥ 1 stimulant (e.g., amphetamine); trial can be waived if member has a history of substance abuse; OR• Trial and failure/intolerance of, contraindication to, of ≥ 1 antidepressant (e.g., imipramine, citalopram) for cataplexy symptoms.
Xyrem ^{CC, QL}	<ul style="list-style-type: none">• Prescriber is a neurologist, sleep medicine, or other specialist in the treatment of narcolepsy; AND• Diagnosis of cataplexy and/or excessive daytime sleepiness associated with narcolepsy; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Documentation of a multiple sleep latency test (MSLT) confirming narcolepsy; AND Trial and failure/intolerance of, contraindication to, ≥ 1 narcolepsy agent (e.g., modafinil); AND Trial and failure/intolerance of, contraindication to, of ≥ 1 stimulant (e.g., amphetamine) for excessive daytime sleepiness symptoms; OR Trial and failure/intolerance of, contraindication to, of ≥ 1 antidepressant (e.g., imipramine, citalopram) for cataplexy symptoms; AND If requesting Xywav: failure of Xyrem due to intolerance or adverse outcome (e.g., hyponatremia) suspected to be caused by high sodium content of Xyrem.
Xywav ^{CC, QL}	<p><i>Cataplexy and excessive daytime sleepiness associated with narcolepsy</i></p> <ul style="list-style-type: none"> Prescriber is a neurologist, sleep medicine, or other specialist in the treatment of narcolepsy; AND Diagnosis of cataplexy and/or excessive daytime sleepiness associated with narcolepsy; AND Documentation of a multiple sleep latency test (MSLT) confirming narcolepsy; AND Trial and failure/intolerance of, contraindication to, ≥ 1 narcolepsy agent (e.g., modafinil); AND Trial and failure/intolerance of, contraindication to, of ≥ 1 stimulant (e.g., amphetamine) for excessive daytime sleepiness symptoms; OR Trial and failure/intolerance of, contraindication to, of ≥ 1 antidepressant (e.g., imipramine, citalopram) for cataplexy symptoms; AND If requesting Xywav: failure of Xyrem due to intolerance or adverse outcome (e.g., hyponatremia) suspected to be caused by high sodium content of Xyrem. <p><i>Idiopathic Hypersomnia</i></p> <ul style="list-style-type: none"> Prescriber is a neurologist, sleep medicine, or other specialist in the treatment of sleep disorders; AND Patient is at least 18 years of age; AND Diagnosis of idiopathic hypersomnia; AND Documentation of a multiple sleep latency test (MSLT) confirming idiopathic hypersomnia; AND Other causes of sleep disorder have been ruled out; AND Trial and failure/intolerance of, contraindication to modafinil; AND Trial and failure/intolerance of, contraindication to, of ≥ 1 stimulant (e.g., amphetamine) for excessive daytime sleepiness symptoms; AND The requested dose does not exceed FDA approved dosing for diagnosis.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
armodafinil tablet ^{CC, QL}	modafinil tablet ^{CC, QL}

Preferred Agents	Non-Preferred Agents
Provigil tablet ^{CC, QL}	Nuvigil tablet ^{CC, QL}
	sodium oxybate solution ^{CC, QL}
	Sunosi tablet ^{CC, QL}
	Wakix tablet ^{CC, QL}
	Xyrem solution ^{CC, QL}
	Xywav solution ^{AE, CC, QL}

CENTRAL NERVOUS SYSTEM: NEUROPATHIC PAIN

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Gabarone ^{CC, QL}	<ul style="list-style-type: none">Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to gabapentin capsule AND gabapentin oral solutionTrial and failure (e.g., allergy or intolerance to an inactive ingredient) with at least 2 manufacturers (if available and covered) of gabapentin capsule AND gabapentin oral solution.
ZTlido ^{CC, QL}	<ul style="list-style-type: none">Diagnosis of post-herpetic neuralgiaTrial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to BOTH of the following:<ul style="list-style-type: none">lidocaine 5% patch; ANDcapsaicin (OTC)Quantity Limit: 3 per day

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
duloxetine DR capsule (generic Cymbalta)	Cymbalta DR capsule
gabapentin capsule ^{QL}	Drizalama sprinkle capsule
gabapentin solution ^{QL}	duloxetine DR capsule (generic Irenka)
gabapentin tablet ^{QL}	Gabarone tablet ^{CC, QL}
Lidocaine patch ^{QL}	Gralise tablet (brand and generic)
pregabalin capsule ^{QL}	Horizant tablet
pregabalin solution ^{QL}	Lyrica capsule ^{QL}
	Lyrica CR tablet ^{QL}
	Lyrica solution ^{QL}

Preferred Agents	Non-Preferred Agents
	Neurontin capsule ^{QL}
	Neurontin solution ^{QL}
	Neurontin tablet ^{QL}
	pregabalin ER tablet ^{QL}
	Savella tablet dose pack
	Savella tablet
	ZTlido patch ^{CC, QL}

CENTRAL NERVOUS SYSTEM: SEDATIVE HYPNOTICS

GUIDELINES FOR USE

Approval Duration: 1 year

1. MAXIMUM DURATION (MD) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
temazepam 15, 30 mg ^{MD, QL} zolpidem ^{MD, QL} Ambien ^{MD, QL} Ambien CR ^{MD, QL} Belsomra ^{MD, QL} Dayvigo ^{MD, QL} Doral ^{MD, QL} doxepin ^{QL} (generic Silenor) Edluar ^{CC, MD, QL} estazolam ^{MD, QL} eszopiclone ^{MD, QL} flurazepam ^{MD, QL} Halcion ^{MD, QL} Lunesta ^{MD, QL} Quviviq ^{AE,CC, MD, QL} ramelteon ^{CC, MD, QL} Restoril ^{MD, QL} Rozerem ^{CC, MD, QL} Silenor ^{QL} temazepam 7.5, 22.5 mg ^{MD, QL} triazolam ^{MD, QL} zaleplon ^{MD, QL} zolpidem ER ^{MD, QL} zolpidem SL ^{MD, QL}	Approval Duration: 6 months <ul style="list-style-type: none"> • Patient has been evaluated for signs and symptoms of abuse, dependency, misuse, or overuse of controlled substances including KASPER monitoring; AND • Patient has had a trial (at least 3 weeks) of non-pharmacological therapies (e.g., stimulus control, sleep restriction, sleep hygiene measures, and relaxation therapy); AND • Patient has a diagnosis of severe or refractory insomnia; AND/OR • Patient has a comorbid condition (e.g., psychiatric disorder, chronic pain) which causes and/or exacerbates insomnia; AND/OR • Patient requires use of a sedative hypnotic medication to maintain compliance with nighttime breathing apparatus (e.g., CPAP); OR • A Clinical Pharmacist may approve the request if there is another valid medical reason why the recipient requires long-term use of the requested medication. • Approval of requests beyond 60 days should be limited to non-benzodiazepine agents (e.g., eszopiclone, suvorexant, zaleplon, zolpidem) wherever possible due to the higher potential for abuse, dependency, and withdrawal associated with benzodiazepines. • Benzodiazepine sedative hypnotics (e.g., estazolam, flurazepam, temazepam, triazolam) should only be approved for long-term use when: <ul style="list-style-type: none"> ▪ Patient has tried and failed a non-benzodiazepine sedative hypnotic (e.g., eszopiclone, suvorexant, zaleplon, zolpidem) or is unable to use these agents due to allergy or contraindication which does not apply to benzodiazepine sedative hypnotics; AND ▪ Patient meets all other above criteria for exceeding the duration limit.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Edluar ^{CC, MD, QL}	<ul style="list-style-type: none">Diagnosis of dysphagia; ORTrial and failure of 2 sedative hypnotics, ONE of which must be zolpidem.
Hetlioz ^{CC, QL} tasimelteon ^{CC, QL}	<ul style="list-style-type: none">Diagnosis of Non-24-hour sleep-wake disorder (“non-24”) in adults ORUsed for the treatment of nighttime sleep disturbances in a patient age ≥ 3 years that has been diagnosed with Smith-Magenis syndrome (SMS).
Igalmi ^{AE, CC, QL}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none">Patient has agitation associated with a confirmed diagnosis of schizophrenia or bipolar disorder, defined as meeting DSM-5 criteria for schizophrenia, schizoaffective, or schizophreniform disorder or bipolar I or II disorder; ANDAgitation is NOT due to acute intoxication; ANDPrescriber attestation that patient will be monitored by a healthcare provider, including an assessment of vital signs and alertness to prevent falls and syncope; ANDPatient is NOT taking medications known to prolong the QT interval; ANDPrescriber attestation that patient has been advised to avoid activities requiring mental alertness for at least 8 hours following administration. <p>Renewal Criteria</p> <ul style="list-style-type: none">Patient must continue to meet the above criteria; ANDPrescriber attestation of response (patient not requiring alternative agents following treatment of mild to moderate agitation); AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none">Patient has not experienced any treatment-restricting adverse effects (e.g., syncope, orthostatic hypotension, fall, QT prolongation, symptomatic bradycardia). <p>Age Limit: ≥ 18 years Quantity Limit: 2 per day</p>

Agent(s) Subject to Criteria	Criteria for Approval
Quviviq ^{AE, CC, MD, QL}	<ul style="list-style-type: none"> Trial and therapeutic failure, allergy, contraindication (including potential drug- drug interactions with other medications) or intolerance of 1 preferred agent. <p>Age Limit: ≥ 18 years Quantity Limit: 1 per day</p>
Rozerem ^{CC, MD, QL}	<ul style="list-style-type: none"> Trial of preferred agents can be waived if there is a history of substance abuse
temazepam 7.5 mg, 22.5 mg ^{MD, QL}	<ul style="list-style-type: none"> Trial and failure of 15 mg dose; OR Prescriber requests 7.5 mg starting dose

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
eszopiclone tablet ^{MD, QL}	Ambien CR tablet ^{MD, QL}
ramelteon tablet ^{MD, QL}	Ambien tablet ^{MD, QL}
temazepam 15 mg, 30 mg capsule ^{MD, QL}	Belsomra tablet ^{MD, QL}
zolpidem ER tablet ^{MD, QL}	Dayvigo tablet ^{MD, QL}
	Doral tablet ^{MD, QL}
	doxepin tablet ^{QL}
	Edluar SL tablet ^{CC, MD, QL}
	estazolam tablet ^{MD, QL}
	flurazepam capsule ^{MD, QL}
	Halcion tablet ^{MD, QL}
	Hetlioz capsule ^{CC, QL}
	Hetlioz LQ suspension ^{CC, QL}
	Igalmi film ^{AE, CC, QL}
	Lunesta tablet ^{MD, QL}
	quazepam tablet ^{MD, QL}
	Quviviq tablet ^{AE, CC, MD, QL}
	Restoril capsule ^{MD, QL}
	Rozerem tablet ^{CC, MD, QL}
	tasimelteon capsule ^{CC, QL}
	temazepam 7.5 mg, 22.5 mg capsule ^{MD, QL}
	triazolam tablet ^{MD, QL}
	zaleplon capsule ^{MD, QL}
	zolpidem capsule ^{MD, QL}
	zolpidem SL tablet ^{MD, QL}

CENTRAL NERVOUS SYSTEM: SKELETAL MUSCLE RELAXANTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 1 week** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **2 preferred agents**.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
dantrolene ^{QL, CC}	<ul style="list-style-type: none">NPD criteria; ORPrescribed for prophylaxis against malignant hyperthermia
Amrix ^{QL, MD} carisoprodol ^{QL, MD} carisoprodol compound ^{QL, MD} Fexmid ^{QL, MD} Soma ^{QL, MD}	<ul style="list-style-type: none">Limited to 21 days of therapy per rolling 30 days; UNLESSPatient has a diagnosis of the following conditions:<ul style="list-style-type: none">Lumbago with sciatica; ORRadiculopathy; ORCervical disc disorder; ORIntervertebral disc disorders with radiculopathy; ORPrescribed by or in consult with neurology, neurosurgery, or orthopedic specialist for another chronic condition.
tizanidine capsules ^{QL}	<ul style="list-style-type: none">Trial and failure of tizanidine tablets at the requested dose.
Zanaflex 8mg	<ul style="list-style-type: none">At least 1-week trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications), or intolerance to generic tizanidine tablets; ANDTrial and failure (e.g., allergy or intolerance to an inactive ingredient) with at least 2 generic manufacturers of tizanidine capsules; ANDProvider has submitted clinical rationale why other dosage forms and/or strengths cannot be used
Quantity Limit: 4 per day	

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
baclofen 5 mg, 10 mg, 20 mg tablet	Amrix ER ^{QL, MD}
cyclobenzaprine ^{QL}	baclofen 15 mg table, solution ^{QL} , suspension ^{QL}
Methocarbamol 500 mg, 750 mg tablet	carisoprodol tablet ^{QL, MD}
orphenadrine ER	carisoprodol/ASA ^{QL, MD}
tizanidine tablet ^{QL}	chlorthalidone ^{QL}
	cyclobenzaprine ER ^{QL}
	Dantrium ^{QL}
	dantrolene ^{QL, CC}
	Fexmid ^{QL, MD}
	Fleqsuvy ^{QL}
	Lorzone ^{QL}
	Lyvispah ^{QL}
	metaxalone ^{QL}
	methocarbamol 1000 mg tablet
	Norgesic
	Norgesic Forte tablet
	orphenadrine/ASA/caffeine
	orphengesic forte
	Ozobax ^{QL}
	Ozobax DS ^{QL}
	Soma ^{QL, MD}
	Tanlor tablet
	tizanidine capsule ^{QL}
	Tonmya ^{AE, QL}
	Zanaflex ^{QL}

CENTRAL NERVOUS SYSTEM: TOBACCO CESSATION

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Chantix ^{AE, QL}	<ul style="list-style-type: none">Age \geq 18 years old.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires \geq 30 day trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with \geq 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
bupropion SR tablet ^{QL}	
Chantix tablet dose pack ^{AE, QL}	
Chantix tablet ^{AE, QL}	
nicotine gum ^{QL}	
nicotine lozenge ^{QL}	
nicotine lozenge mini ^{QL}	
nicotine patch ^{QL}	
Nicotrol nasal spray ^{QL}	
Varenicline dose pack ^{AE, QL}	
Varenicline tablet ^{AE, QL}	

SPINAL MUSCULAR ATROPHY

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable.

2. NON-PREFERRED (NPD) CRITERIA

Not applicable.

3. BRAND MEDICALLY NECESSARY CRITERIA

Not applicable.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Evrysdi oral solution ^{CC} Evrysdi oral tablet ^{CC, QL}	<p>Initial Approval Criteria:</p> <p>Infantile-onset (Type 1) Spinal Muscular Atrophy (SMA)</p> <ul style="list-style-type: none">• Prescribed by or in consultant with a pediatric neurologist or other specialist in the diagnosis and treatment of spinal muscular atrophy (SMA); AND• Diagnosis of spinal muscular atrophy (SMA) Type 1; AND• Genetic test results (i.e., laboratory results) confirming SMA:<ul style="list-style-type: none">○ Homozygous deletion or mutation of the survival motor neuron 1 (SMN1) gene; OR○ Compound heterozygous mutation of the SMN1 gene; AND○ At least two copies of the SMN2 gene; AND• Patient does not require permanent ventilation (defined as requiring a tracheostomy or more than 21 consecutive days of either non-invasive ventilation (≥ 16 hours per day) or intubation, in the absence of an acute reversible event); AND• Prescriber conducts and submits documentation of an assessment of baseline motor function using at least one of the following:<ul style="list-style-type: none">○ Hammersmith Infant Neurologic Exam-Part 2 (HINE-2)○ Hammersmith Functional Motor Scale Expanded (HFMSE)○ Upper Limb Module (ULM) score○ Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP-INTEND); AND• Not to be used in combination with Spinraza (nusinersen); AND• Patient has not received treatment with Zolgensma (onasemnogene abeparvovec-xioi). <p>Later-onset SMA</p> <ul style="list-style-type: none">• Prescribed by or in consultation with a neurologist or other specialist in the diagnosis and treatment of spinal muscular atrophy (SMA); AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Diagnosis of spinal muscular atrophy (SMA) Type 2 or 3; AND • Prescriber attestation/opinion that patient is non-ambulatory (e.g., requires wheelchair, not able to walk unassisted, etc.); OR • Prescriber attestation/opinion that patient is experiencing a decline in motor function/failure to achieve motor milestones; AND • Prescriber conducts and submits documentation of an assessment of baseline motor function using at least one of the following: <ul style="list-style-type: none"> ○ Hammersmith Infant Neurologic Exam-Part 2 (HINE-2) ○ Hammersmith Functional Motor Scale Expanded (HFMSE) ○ Upper Limb Module (ULM) score ○ Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP-INTEND); AND • Not to be used in combination with Spinraza (nusinersen); AND Patient has not received treatment with Zolgensma (onasemnogene abeparvovec-xioi). <p>Renewal criteria (all requests):</p> <ul style="list-style-type: none"> • Individual does not require use of invasive ventilation or tracheostomy as a result of advanced SMA disease; AND • Repeat motor function testing must be performed at every 12 month interval and must show additional motor improvement from the previous demonstrated motor improvement or that the patient demonstrates clinically significant improvements in SMA associated symptoms (such as a lack of disease progression, stabilization, or decreased decline in motor function, as compared to the natural history trajectory of the disease) evident by the comparative assessment of baseline motor function measurements using one of the following assessments: <ul style="list-style-type: none"> ○ Hammersmith Infant Neurologic Exam-Part 2 (HINE-2) must demonstrate: <ul style="list-style-type: none"> ▪ An improvement or maintenance of previous improvement of at least 2 points (or maximal score) increase in ability to kick; OR ▪ An improvement or maintenance of previous improvement of at least 1-point increase in any other HINE-2 milestone (e.g., head control, rolling, sitting, crawling, etc.) excluding voluntary grasp; AND ▪ The patient exhibited improvement, or maintenance of previous improvement in more HINE-2 motor milestones than worsening, from pretreatment baseline (net positive improvement); OR ▪ Has achieved and maintained any new motor milestones when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk, etc.). ○ Hammersmith Functional Motor Scale Expanded (HFMSE) must demonstrate: <ul style="list-style-type: none"> ▪ An improvement or maintenance of previous improvement of at least a 3-point increase in score from pretreatment baseline; OR ▪ Has achieved and maintained any new motor milestones when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk, etc.). ○ Upper Limb Module (ULM) score must demonstrate:

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ▪ An improvement or maintenance of previous improvement of at least a 2-point increase in score from pretreatment baseline; OR ▪ Has achieved and maintained any new motor milestones when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk, etc.). ○ Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP-INTEND) must demonstrate: <ul style="list-style-type: none"> ▪ An improvement or maintenance of previous improvement of at least a 4-point increase in score from pretreatment baseline; OR ▪ Has achieved and maintained any new motor milestones when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk, etc.). • Not to be used in combination with Spinraza (nusinersen); AND • Patient has not received treatment with Zolgensma (onasemnogene abeparvovec-xioi).
Spinraza ^{CC}	<p>Approval Duration: 6 months initial; 12 months renewal</p> <p>Initial Approval Criteria (must meet all requirements):</p> <ul style="list-style-type: none"> • Prescribed by or in consultant with a pediatric neurologist or other specialist in the diagnosis and treatment of spinal muscular atrophy (SMA); AND • Clinical documentation (e.g., progress notes) supporting diagnosis of Spinal Muscular Atrophy (SMA) type I, II, or III AND • Diagnosis/genetic testing results (official laboratory results) confirming 5q SMA: <ul style="list-style-type: none"> ○ Homozygous deletion or mutation of the survival motor neuron 1 (SMN1) gene; OR ○ Compound heterozygous mutation of the SMN1 gene; AND ○ At least two copies of the SMN2 gene; AND • Patient is NOT maintained on permanent assisted ventilation in the absence of an acute, reversible event prompting the respiratory support; defined as: <ul style="list-style-type: none"> ○ Tracheostomy or ventilator support for ≥ 16 hours per day for > 21 continuous days; OR ○ Use of non-invasive ventilation beyond sleep > 12 hours in a 24 hour period. AND • Prescriber agrees to assess and monitor the following laboratory values throughout treatment: <ul style="list-style-type: none"> ○ Complete blood count (CBC); AND ○ Quantitative spot urine protein testing; AND ○ Prothrombin Time (PT) or Activated Partial Thromboplastin Time (aPTT) • Prescriber conducts, and submits documentation of, an assessment of baseline motor function using at least one of the following: <ul style="list-style-type: none"> ○ Hammersmith Infant Neurologic Exam-Part 2 (HINE-2) ○ Hammersmith Functional Motor Scale Expanded (HFMSE) ○ Upper Limb Module (ULM) score ○ Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP-INTEND) • Not to be used in combination with Evrysdi (risdiplam); AND

Agent(s) Subject to Criteria**Criteria for Approval**

- Patient has not received treatment with Zolgensma (onasemnogen abeparvovec-xioi).

Renewal Criteria (must meet all requirements):

- All initial approval requirements continue to be met; **AND**
- Individual does not require use of permanent assisted ventilation as a result of advanced SMA disease; **AND**
- The patient shall be considered a Responder to therapy by showing an improvement (rather than progression or lack of improvement) in motor function in accordance with the assessments outlined below (HINE-2, HFMSE, ULM, and/or CHOP-INTEND) after the initial 5 loading doses; **AND**
- Repeat motor function testing must be performed at every 6 month interval and must show additional motor improvement from the previous demonstrated motor improvement or that the patient demonstrates clinically significant improvements in SMA associated symptoms (such as a lack of disease progression, stabilization, or decreased decline in motor function, as compared to the natural history trajectory of the disease) evident by the comparative assessment of baseline motor function measurements using one of the following assessments:
 - Hammersmith Infant Neurologic Exam-Part 2 (HINE-2) must demonstrate:
 - An improvement or maintenance of previous improvement of at least 2 points (or maximal score) increase in ability to kick; **OR**
 - An improvement or maintenance of previous improvement of at least 1-point increase in any other HINE-2 milestone (e.g., head control, rolling, sitting, crawling, etc.) excluding voluntary grasp; **AND**
 - The patient exhibited improvement, or maintenance of previous improvement in more HINE-2 motor milestones than worsening, from pretreatment baseline (net positive improvement); **OR**
 - Has achieved and maintained any new motor milestones when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk, etc.).
 - Hammersmith Functional Motor Scale Expanded (HFMSE) must demonstrate:
 - An improvement or maintenance of previous improvement of at least a 3-point increase in score from pretreatment baseline; **OR**
 - Has achieved and maintained any new motor milestones when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk, etc.).
 - Upper Limb Module (ULM) score must demonstrate:
 - An improvement or maintenance of previous improvement of at least a 2-point increase in score from pretreatment baseline; **OR**
 - Has achieved and maintained any new motor milestones when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk, etc.).
 - Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP-INTEND) must demonstrate:
 - An improvement or maintenance of previous improvement of at least a 4-point increase in score from pretreatment baseline; **OR**

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ▪ Has achieved and maintained any new motor milestones when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk, etc.). • Provider must provide clinical documentation (chart/progress notes) from the most recent office visit and evaluation; AND • Not to be used in combination with Evrysdi (risdiplam); AND • Patient has not received treatment with Zolgensma (onasemnogene abeparvovec-xioi).
Zolgensma ^{CC}	<p>Approval Duration: Date of service; once per lifetime</p> <ul style="list-style-type: none"> • Prescribed by or in consultation with a pediatric neurologist or other specialist in the diagnosis and treatment of spinal muscular atrophy (SMA); AND • Must have SMA confirmed by submission of medical records (e.g., chart notes, laboratory values): <ul style="list-style-type: none"> ○ A mutation or deletion of genes in chromosome 5q resulting in one of the following: <ul style="list-style-type: none"> - Homozygous gene deletion or mutation of SMN1 gene (e.g., homozygous deletion of exon 7 at locus 5q13); OR - Compound heterozygous mutation of SMN1 gene (e.g., deletion of SMN1 exon 7 {allele 1} and mutation of SMN1 {allele 2}); AND • Patient meets one of the following: <ul style="list-style-type: none"> ○ Symptomatic SMA diagnosed by or in consultation with a neurologist with disease expertise; OR ○ SMA diagnosis confirmed by newborn screening; AND ○ 4 copies or less of the SMN2 gene; AND • Not have advanced SMA (e.g., permanent ventilation support; complete limb paralysis); AND • Not have pre-existing hepatic insufficiency; AND • Baseline anti-AAV9 antibody titer of $\leq 1:50$ (as measured by ELISA); AND • Must be used with systemic corticosteroids (e.g., 1 mg/kg/day oral prednisone or equivalent) as directed; AND • Therapy to be administered prior to recipient's 2nd birthday; AND • Not to be used in combination with Spinraza (nusinersen); AND • Not to be used in combination with Evrysdi (risdiplam).

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Not applicable	Evrysdi oral solution ^{CC}
	Evrysdi oral tablet ^{CC, QL}
	Spinraza vial ^{CC}
	Zolgensma kit ^{CC}

ANALGESICS: NARCOTICS, LONG-ACTING

GUIDELINES FOR USE

Approval Duration: 6 months (1 year for active cancer, sickle cell anemia or hospice/palliative care)

1. PREFERRED WITH PA (PDP) CRITERIA (ALSO APPLIES TO NON-PREFERRED AGENTS)

Agent(s) Subject to Criteria	Criteria for Approval
Butrans ^{CC, QL} fentanyl transdermal 12, 25, 50, 75, 100 mcg ^{CC, QL} morphine sulfate ER (generic MS Contin) ^{CC, QL} OxyContin ER tablet ^{CC, QL} tramadol ER (generic Ultram ER) ^{CC, AE, QL}	<ul style="list-style-type: none">• Opioid Class Criteria for Initial Approval must be met; AND• Patient has severe pain requiring daily, around-the-clock, long-term pain management as evidenced by:<ul style="list-style-type: none">○ Pain lasting > 3 consecutive months; AND○ Trial and failure within the past 90 days of 1 non-opioid analgesic (i.e., NSAIDs, APAP) at maximum tolerated doses without pain relief and/or functional improvement; AND○ Trial and failure within the past 90 days of at least 1 short-acting opioid analgesic at maximum tolerated doses without adequate relief of pain.• Additional criteria as applicable:<ul style="list-style-type: none">○ Class Criteria for Naloxone Prescribing○ Class Criteria for High Morphine Milligram Equivalent (MME) Requests – Over 90 MME per Day○ Class Criteria for Approval of Very High MME Requests – Over 200 MME per Day○ Class Criteria for Opioids and Benzodiazepines

1. OPIOID CLASS CRITERIA

- a. Refer to ANALGESICS (CLINICAL CRITERIA FOR SHORT-ACTING AND LONG-ACTING OPIOIDS) section.

2. OPIOID LONG-ACTING DUPLICATE THERAPY CRITERIA

- a. Prescriber has discontinued the second long-acting opioid; **OR**
- b. Prescriber has a plan to taper off one of the long-acting opioids; **OR**
- c. Prescriber provided medical justification for continuation.

3. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents unless otherwise specified. Preferred with PA (PDP) Criteria must be met.

4. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

5. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

6. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
methadone ^{CC}	<ul style="list-style-type: none">• Approve for 30 days only in infants up to 1 year of age who are discharged from the hospital on a methadone taper for neonatal abstinence syndrome (NAS); OR• If used for pain, all of the following criteria apply:<ul style="list-style-type: none">○ Opioid Class Criteria for Initial Approval must be met; AND○ Prescriber is a pain management specialist or prescriber has proof of consultation with a pain management specialist; AND○ Severe pain requiring daily, around-the-clock, long-term pain management, defined as:<ul style="list-style-type: none">▪ Pain lasting > 6 consecutive months; AND▪ Trial and failure of one non-opioid analgesic (i.e., NSAIDs, APAP) at maximum tolerated doses without adequate relief of pain; AND▪ Trial and failure of one short-acting opioid analgesic at maximum tolerated doses without adequate relief of pain; AND○ Trial and failure of two preferred long-acting opioids; AND○ Patient does not have a history of drug or alcohol abuse/dependence or addiction (drug and alcohol toxicology screen results dated within the past month must be submitted with the PA request); AND○ If the patient is female between the ages of 18 and 45 years of age, prescriber must attest to the fact that patient has been counseled regarding the risks of becoming pregnant while on this medication, including the risk of neonatal abstinence syndrome (NAS); AND○ Patient is not presently taking any other long-acting opioids. <p>Note: Methadone will not be approved for drug addiction as a pharmacy benefit</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Butrans ^{CC, QL}	Belbucca ^{AE, QL}

Preferred Agents	Non-Preferred Agents
fentanyl patch 12, 25, 50, 75, 100 mcg ^{CC, QL}	buprenorphine patch ^{QL}
morphine sulfate ER tablet ^{CC, QL}	ConZip ER capsule ^{AE, QL}
OxyContin ER tablet ^{CC, QL}	Diskets
tramadol ER tablet (generic Ultram ER) ^{CC, AE, QL}	fentanyl patch 37.5, 62.5, 87.5 mcg ^{QL}
	hydrocodone ER capsule ^{QL}
	hydrocodone ER tablet ^{QL}
	hydromorphone ER tablet ^{QL}
	Hysingla ER tablet ^{QL}
	methadone dispersible tablet ^{CC}
	methadone intensol oral concentrate ^{CC}
	methadone oral concentrate ^{CC}
	methadone solution
	methadone tablet
	methadose oral concentrate
	methadose tablet
	morphine sulfate ER capsule ^{QL}
	MS Contin ER tablet ^{QL}
	oxycodone ER tablet ^{QL}
	oxymorphone ER tablet ^{QL}
	tramadol ER capsule ^{AE, QL}
	tramadol ER tablet (generic Ryzolt) ^{AE, QL}

ANALGESICS: NARCOTICS, SHORT-ACTING

GUIDELINES FOR USE

Approval Duration: 1 month, 3 months, or 6 months based on full opioid criteria (1 year for active cancer, sickle cell anemia or hospice/palliative care)

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
codeine/APAP ^{CC, AE, MD, QL} endocet ^{CC, MD, QL} hydrocodone/APAP ^{CC, MD, QL} hydrocodone/ibuprofen ^{CC, MD, QL} hydromorphone tablets ^{CC, MD, QL} morphine concentrate ^{CC, MD} morphine solution, tablets ^{CC, MD, QL} oxycodone solution, tablets ^{CC, MD, QL} oxycodone/APAP tablets ^{CC, MD, QL} tramadol 50 mg ^{CC, MD, AE, QL} tramadol/APAP ^{MD, AE, QL}	<p>Codeine- and tramadol-containing products: Minimum age of 18 years</p> <p>PA required when:</p> <ul style="list-style-type: none"> The claim is for > 7-day supply for members ≥ 18 years old; OR The claim is for > 3-day supply for members < 18 years old; OR The claim brings the cumulative supply of short-acting opioids in the past 90 days to > 14 days; OR Product is ≥ 30 morphine milligram equivalents (MME) in a single dosing unit (e.g., hydromorphone 8 mg tablet) or a concentrated liquid (e.g., morphine sulfate 20 mg/mL). <p>30-day approval:</p> <ul style="list-style-type: none"> Only 1 short-acting opioid will be used at a time; AND Trial and failure of, or contraindication to, at least 1 non-opioid pain medication (e.g., APAP, NSAIDs); OR Medication is prescribed by a treating physician within 14 days of: <ul style="list-style-type: none"> A major surgery, any operative or invasive procedure or a delivery; OR A significant trauma, being any acute blunt, blast, or penetrating bodily injury that has a risk of death, physical disability, or impairment; OR Other clinical justification as to why treatment with opioids should extend beyond 14 days and provide a diagnosis more specific than pain. If the request is for a high strength or concentrated dosage form, the prescriber must submit rationale why lower strength or less-concentrated products cannot be used. Additional criteria as applicable: <ul style="list-style-type: none"> Class Criteria for Naloxone Prescribing Class Criteria for High Morphine Milligram Equivalent (MME) Requests – Over 90 MME per Day; OR Class Criteria for Approval of Very High MME Requests – Over 200 MME per Day Class Criteria for Opioids and Benzodiazepines <p>3- or 6-month approval:</p> <ul style="list-style-type: none"> Opioid Class Criteria for Initial Approval must be met; AND Prescriber must submit a diagnosis more specific than pain; AND Trial and failure of, or contraindication to, at least 1 non-opioid pain medication (e.g., APAP, NSAIDs) within the past 6 months; OR Medication is prescribed by a treating physician within 30 days of:

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ A major surgery, any operative or invasive procedure or a delivery; OR ○ A significant trauma, being any acute blunt, blast, or penetrating bodily injury that has a risk of death, physical disability, or impairment; OR ○ Other clinical justification as to why treatment with opioids should extend beyond 30 days. ● If short-term pain management is expected/indicated, approve for 3 months; OR ● If long-term (e.g., > 3 months) pain management is expected/indicated OR patient is currently taking a long-acting narcotic; approve for 6 months. ● If the request is for a high strength or concentrated dosage form, the prescriber must submit rationale why lower strength or less-concentrated products cannot be used. ● Additional clinical justification will be required for doses that exceed quantity limits (e.g., 90 MME). ● Additional criteria as applicable: <ul style="list-style-type: none"> ○ Class Criteria for Naloxone Prescribing ○ Class Criteria for High Morphine Milligram Equivalent (MME) Requests – Over 90 MME per Day; OR ○ Class Criteria for Approval of Very High MME Requests – Over 200 MME per Day ○ Class Criteria for Opioids and Benzodiazepines

2. OPIOID CLASS CRITERIA

Refer to ANALGESICS (CLINICAL CRITERIA FOR SHORT-ACTING AND LONG-ACTING OPIOIDS) section.

3. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 1 week** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified.

Preferred with PA (PDP) Criteria must be met.

4. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

5. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

6. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Ascomp® with codeine ^{CC, AE, QL}	Approval Duration: 1 year
butalbital/APAP/caffeine/codeine ^{CC, QL}	<ul style="list-style-type: none"> ● Diagnosis of one of the following headache disorders: <ul style="list-style-type: none"> ○ Muscular headache; OR ○ Tension-type headache; OR ○ Migraine.
butalbital/ASA/caffeine/codeine ^{CC, AE, QL}	
butalbital compound/codeine ^{CC, AE, QL}	Age Limit: ≥ 18 years

Agent(s) Subject to Criteria	Criteria for Approval
	Quantity Limit: 1 per day (30 per 30 days) - Not meant for daily use: up to 6 per headache day; 5 headache days per month

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
codeine/APAP solution ^{CC, AE, MD, QL}	APAP/caffeine/dihydrocodeine capsule ^{MD, QL}
codeine/APAP tablet ^{CC, AE, MD, QL}	ASA/butalbital/caffeine/codeine capsule ^{CC, AE, QL}
hydrocodone/APAP solution ^{CC, MD, QL}	Ascomp with codeine capsule ^{CC, AE, QL}
hydrocodone/APAP tablet ^{CC, MD, QL}	butalbital/APAP/caffeine/codeine capsule ^{CC, AE, QL}
hydrocodone/ibuprofen tablet ^{CC, MD, QL}	butalbital/codeine capsule ^{CC, AE, QL}
hydromorphone tablet ^{CC, MD, QL}	codeine tablet ^{MD, AE, QL}
morphine concentrate ^{CC, MD}	Dilaudid liquid ^{MD, QL}
morphine solution ^{CC, MD, QL}	Dilaudid tablet ^{MD, QL}
morphine syringe 10 mg/0.5 mL ^{CC, MD}	Fioricet with codeine capsule ^{CC, AE, QL}
morphine tablet ^{CC, MD, QL}	hydromorphone liquid ^{MD, QL}
oxycodone solution ^{CC, MD, QL}	hydromorphone suppository ^{MD, QL}
oxycodone tablet ^{CC, MD, QL}	levorphanol tablet ^{MD, QL}
oxycodone/APAP tablet ^{CC, MD, QL}	meperidine solution ^{MD, QL}
tramadol 50 mg tablet ^{CC, MD, AE, QL}	meperidine tablet ^{MD, QL}
tramadol/APAP tablet ^{MD, AE, QL}	morphine suppository ^{MD, QL}
	morphine syringe 20 mg/mL ^{MD}
	Nalocet tablet ^{MD, QL}
	oxycodone capsule ^{MD, QL}
	oxycodone concentrate ^{MD, QL}
	oxycodone oral syringe ^{MD, QL}
	oxycodone/APAP solution ^{MD, QL}
	oxymorphone tablet ^{MD, QL}
	Percocet tablet ^{MD, QL}
	Prolate solution ^{MD, QL}
	Prolate tablet ^{MD, QL}
	Qdolo ^{MD, AE, QL}
	Roxicodone tablet ^{MD, QL}
	Roxybond tablet ^{MD, QL}
	Seglantis tablet ^{MD, AE, QL}
	tramadol 25 mg tablet ^{MD, AE, QL}
	tramadol 75 mg tablet ^{MD, AE, QL}
	tramadol 100 mg tablet ^{MD, AE, QL}
	tramadol solution ^{MD, AE, QL}

ANALGESICS: CLINICAL CRITERIA FOR SHORT-ACTING AND LONG-ACTING OPIOIDS

GUIDELINES FOR USE

**Approval Duration: Varies
(1 year for active cancer, sickle cell anemia or hospice/palliative care)**

NOTE: Class criteria will be waived for members receiving hospice/palliative/end-of-life care or have a diagnosis of active cancer or sickle cell anemia.

1. CLASS CRITERIA FOR INITIAL APPROVAL

Exception apply to short-acting opioids for acute pain; additional criteria may also apply to specific formulation).

- a. Prescriber has evaluated the member for risk of diversion, harm, or misuse:
 - i. Prescriber attests that KASPER report for the past 12 months has been reviewed; **AND**
 - ii. Prescriber submits urine drug screen (UDS) results dated within the past 30 days for ALL new chronic opioid (e.g., beyond 45 days of treatment) requests **UNLESS** the member is in a long-term care or skilled nursing facility. Note: UDS is not required for acute prescribing; **AND**
 - iii. If UDS is positive for illicit or unexpected substances, prescriber attests that naloxone was or will be prescribed; **AND**
- b. Prescriber submits an assessment of baseline pain and function using an objective measure; **AND**
- c. Prescriber attestation or documentation that non-opioid therapies (e.g., exercise therapy, cognitive behavioral therapy, NSAIDs, etc.) have been tried and/or are being used and optimized as appropriate; **AND**
- d. For females of child-bearing age, prescriber attests that the member has been counseled regarding the risks of becoming pregnant while on this medication, including the risk of neonatal abstinence syndrome (NAS); **AND**
- e. Patient does NOT have respiratory depression, acute or severe bronchial asthma, or hypercarbia; **AND**
- f. Patient does NOT have known or suspected GI obstruction (e.g., paralytic ileus); **AND**
- g. Up to 1 long-acting opioid and 1 short-acting opioid may be used at a time.

2. CLASS CRITERIA FOR HIGH MORPHINE MILLIGRAM EQUIVALENT (MME) REQUESTS – OVER 90 MME PER DAY

- a. Additional criteria shall apply for NEW requests where the cumulative opioid dose across all prescriptions is > 90 morphine milligram equivalents (MME):
 - i. Note: Buprenorphine products (for opioid addiction treatment or pain) are not assigned an MME value and will not be included in the calculation.
 - ii. Prescriber is, or has proof of consultation with, a Pain Management Specialist OR specialist in an appropriate discipline (e.g., orthopedist, neurologist, spine specialist, etc.) for evaluation of the source of pain and/or treatment of any underlying conditions; **AND**
 - iii. Prescriber must submit clinical justification for exceeding 90 MME/day; **AND**
 - iv. Prescriber attests that a naloxone prescription and associated counseling on its use was or will be offered to the member.

3. CLASS CRITERIA FOR APPROVAL OF VERY HIGH MME REQUESTS – OVER 200 MME PER DAY

- a. Additional criteria shall apply any requests where the cumulative opioid dose across all prescriptions is > 200 MME/day:
 - i. Note: Buprenorphine products (for opioid addiction treatment or pain) are not assigned an MME value and will not be included in the calculation.
 - ii. Prescriber is, or has proof of consultation with, a Pain Management Specialist; **AND**
 - iii. Prescriber submits clinical justification for exceeding 200 MME/day; **AND**
 - iv. Prescriber submits documentation (e.g., progress notes) showing attempts and/or plans to taper below 200 MME/day as well as other non-opioid components (e.g., NSAIDs, physical therapy, etc.) of the treatment plan; **AND**
 - v. Prescriber attests that a naloxone prescription and associated counseling on its use was or will be given to the member.

4. CLASS CRITERIA FOR OPIOIDS AND BENZODIAZEPINES

- a. Additional criteria shall apply when opioids are prescribed concurrently with benzodiazepines and/or KASPER report shows a benzodiazepine prescription in the past 12 months:
 - i. Prescriber must submit clinical justification for the concurrent use of benzodiazepines and opioids; **AND**
 - ii. Prescriber attests that the member and/or caregiver(s) has been or will be counseled about the increased risks of slowed or difficult breathing and/or excessive sedation, and the associated signs and symptoms; **AND**
 - iii. Prescriber attests that a naloxone prescription and associated counseling on its use was or will be given to the member.

5. CLASS CRITERIA FOR OPIOIDS AND BUPRENORPHINE

- a. Prescriber verifies knowledge of the patient's relapse and agrees to increase psychosocial counseling **AND** prescriber provides the dates of planned counseling sessions; **OR**
- b. Opioid(s) is being used short-term (≤ 30 days) for an acute injury leading to acute pain.

Note: Requests for 2 different strengths are considered a therapy duplication and may be overridden if total mg/day does not exceed established limits or quantity limits for each specific strength.

6. CLASS CRITERIA FOR NALOXONE PRESCRIBING

- a. Prescriber attests that a naloxone prescription and associated counseling on its use was or will be offered to the member when any of the following are true (e.g., found on KASPER report, medication list, or diagnosis list):
 - i. Opioid(s) is/are concurrently prescribed with a skeletal muscle relaxant (e.g., cyclobenzaprine); **OR**
 - ii. Opioid(s) is/are concurrently prescribed with a sedative hypnotic (e.g., zolpidem); **OR**
 - iii. Opioid(s) is/are concurrently prescribed with gabapentin or pregabalin; **OR**
 - iv. Member has a history of opioid or other controlled substance overdose; **OR**
 - v. Member has a history of substance use disorder (SUD).

7. OPIOID RENEWAL CRITERIA

- a. Prescriber must submit proof of monitoring for evidence of diversion, harm, and misuse:
 - i. Attest that KASPER report has been checked within the past 3 months; **AND**
 - ii. If the member is not in a long-term care or skilled nursing facility, require prescriber to document risk assessment and provide most recent UDS results dated within:
 - a) One year if considered “low risk”; **OR**
 - b) Six months if considered “moderate risk”; **OR**
 - c) Three months if considered “high risk”; **AND**
 - iii. Prescriber explanation is required if UDS is positive for illicit or unexpected substances; **AND**
 - iv. If UDS is positive for illicit or unexpected substances, prescriber attests that naloxone was or will be prescribed.
- b. Prescriber must submit an assessment of current pain and function using an objective measure; **AND**
- c. Recipient should demonstrate a 30% improvement from baseline to continue current dose or documentation (e.g., progress note) that includes the rationale for continued opioid therapy at the current dose; **AND**
- d. Prescriber must report whether patient has required use of opioid rescue medication (e.g., naloxone) or has been hospitalized or otherwise treated for opioid or other controlled substance overdose in the past 6 months; **AND**
 - i. If member has opioid overdose or use of naloxone within the past 6 months, the prescriber must submit documentation (e.g., progress notes) a plan for preventing future overdoses (e.g., dose reduction of opioid or opioid potentiator[s]; discontinuation of opioid potentiator[s]).

ANALGESICS: NARCOTICS, FENTANYL CITRATE PRODUCTS

GUIDELINES FOR USE

Approval Duration: 6 months

Prior Approval for the medications of this edit must be obtained by the prescriber's office ONLY.

Prescribers/prescriber's agents or LTC facility-based prescribers/prescriber's agents must initiate the following prior authorization requests. Pharmacies/LTC pharmacies and their staff are not permitted to initiate these requests. For members flagged with an LTC eligibility segment, the prescriber signature prior authorization requirements MUST be satisfied by a FACILITY-BASED prescriber/prescriber agent (for faxed requests).

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable.

2. NON-PREFERRED (NPD) CRITERIA

- a. Diagnosis of cancer pain unresponsive to any other therapy; **AND**
- b. Patients must be receiving, and be tolerant to, opioid therapy; **AND**
- c. Patients must have tried opioid doses greater than or equal to either Morphine 60 mg daily or Fentanyl Patches 50 mcg/hour for at least one week.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. tried and failed or had a contraindication to or intolerance to the generic equivalent agent before obtaining approval for the branded agent. These requests must be submitted on the Brand Medically Necessary Form; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents – PA Required	Non-Preferred Agents – PA Required
None	fentanyl citrate lozenge ^{CC, QL} fentanyl citrate tablet ^{CC, QL}

ANALGESICS: NARCOTIC AGONISTS/ANTAGONISTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial of appropriate duration and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in the Analgesics: Narcotics, Short-Acting class.

3. BRAND MEDICALLY NECESSARY CRITERIA

Not applicable.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
None	butorphanol nasal spray pentazocine/naloxone tablet ^{QL}

ANALGESICS: NON-STEROIDAL ANTI-INFLAMMATORY DRUGS (NSAIDS)

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 3 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Duexis (ibuprofen/famotidine) ^{CC} naproxen/esomeprazole ^{CC, QL} Vimovo (naproxen/esomeprazole) ^{CC, QL}	<ul style="list-style-type: none">NPD Criteria above; ORTrial and failure (e.g., poor adherence) of individual, generic components.
Elyxyb solution ^{AE, CC, QL} meloxicam capsules ^{CC} Vivlodex (meloxicam submicronized) ^{CC, QL} Zorvolex (diclofenac submicronized) ^{CC}	<ul style="list-style-type: none">NPD Criteria above; OR3- day trial and failure of two preferred agents; AND<ul style="list-style-type: none">Trial and failure of the preferred, generic formulation of the same ingredient as 1 of the 2 preferred drug trials. <p>Age Edit: ≥ 18 years (Elyxyb)</p>
diclofenac epolamine patches ^{CC} diclofenac 2% solution pump ^{CC} diclofenac topical solution ^{CC} Flector ^{CC} ketorolac nasal spray ^{CC} Licart ^{CC} Pennsaid ^{CC} Sprix ^{CC} celecoxib 400 mg ^{QL}	<ul style="list-style-type: none">NPD Criteria above; ORTrial and failure of diclofenac 1% topical gel; AND<ul style="list-style-type: none">Contraindication to oral NSAIDs; ORUnable to tolerate, swallow, or absorb oral NSAIDs. <ul style="list-style-type: none">Allow up to 17 capsules in 8 days when high dose regimen is needed for acute gout: 800 mg orally immediately, followed by 400 mg 12 hours later and then 400 mg every 12 hours for 7 days.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
celecoxib ^{QL}	Arthrotec
diclofenac sodium topical gel (1%)	Celebrex ^{QL}
diclofenac sodium DR/EC tablets	Daypro
Ibu tablet	diclofenac epolamine patch ^{CC}
ibuprofen suspension	diclofenac potassium capsule
ibuprofen tablet (except 300 mg)	diclofenac potassium powder pack
indomethacin capsule	diclofenac potassium tablet
indomethacin ER capsule	diclofenac topical solution ^{CC}
ketorolac tablet	diclofenac sodium SR/ER tablet
meloxicam tablet	diclofenac 2% solution pump ^{CC}
nabumetone tablet	diclofenac sodium/misoprostol
naproxen sodium tablet	diflunisal tablet
naproxen tablet	Dolobid tablet ^{QL}
piroxicam capsule	Duexis tablet ^{CC}
sulindac tablet	EC-Naprosyn tablet
	EC-naproxen tablet
	Elyxyb solution ^{CC, AE, QL}
	etodolac capsule
	etodolac tablet
	etodolac ER tablet
	Feldene capsule
	fenoprofen capsule
	fenoprofen tablet
	Fenopron capsule
	Flector patch ^{CC}
	flurbiprofen tablet
	ibuprofen 300 mg tablet
	ibuprofen/famotidine tablet
	indomethacin suppository
	indomethacin suspension ^{QL}
	ketoprofen ER capsule
	ketoprofen capsule
	ketorolac nasal spray ^{CC}
	Kiprofen capsule
	Licart patch ^{CC}
	Lofena tablet
	meclofenamate capsule
	mefenamic acid capsule
	meloxicam capsule ^{CC, QL}
	Nalfon tablet
	Naprelan CR tablet
	Naprosyn suspension
	naproxen DR tablet
	naproxen suspension
	naproxen sodium CR/ER tablet
	naproxen/esomeprazole DR tablet ^{CC, QL}
	oxaprozin tablet
	Pennsaid ^{CC}
	Relafen tablet
	Relafen DS tablet
	Tolectin 600 tablet
	tolmetin capsule
	Tolmetin tablet ^{QL}
	Vimovo ^{CC, QL}
	Vyscoxa suspension ^{QL}

ANALGESICS: OPIATE DEPENDENCE TREATMENTS

GUIDELINES FOR USE

Approval Duration: Date of Service Only

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Not applicable.

3. BRAND MEDICALLY NECESSARY CRITERIA

Not applicable.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

6. THERAPEUTIC DUPLICATION

- a. Buprenorphine-containing products will deny for therapeutic duplication when:
 - i. There is a claim for any opioid in the past 30 days; **OR**
 - ii. There is a claim for another buprenorphine-containing product in the past 90 days.
- b. Only the buprenorphine prescriber's office can request these overrides; they will be made aware of the narcotic in history.
- c. Date-of-Service (DOS) approvals can be granted when ONE of the following apply:
 - i. The prescriber verifies knowledge of the patient's relapse and agrees to increase psychosocial counseling. Please obtain dates of planned counseling sessions. If no planned sessions, do not approve, **OR**
 - ii. The narcotic analgesic is being used short-term (30 days or less) for an acute injury leading to acute pain.
 - iii. Requests for 2 different strengths are considered a therapy duplication. Pharmacist may override if total mg/day does not exceed established limits or exceed quantity limits for each specific strength.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Brixadi	None
Buprenorphine SL tablet ^{QL}	
buprenorphine/naloxone SL film ^{QL}	
buprenorphine/naloxone SL tablet ^{QL}	
lofexidine tablet ^{QL}	
Lucemyra tablet ^{QL}	
naltrexone tablet	
Sublocade ER syringe ^{QL}	

Preferred Agents		Non-Preferred Agents	
Suboxone film ^{QL}			
Vivitrol ER suspension			
Zubsolv SL tablet ^{QL}			

ANTI-INFECTIVE: ORAL ANTIFUNGALS

GUIDELINES FOR USE

Approval Duration: 6 months, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
itraconazole capsule ^{CC, QL}	<ul style="list-style-type: none">Prescribed for the treatment or prophylaxis (in an immunocompromised patient) of systemic fungal infection (e.g., aspergillosis, esophageal candidiasis, histoplasmosis); ORDiagnosis of onychomycosis, tinea, or other superficial fungal infection; ANDTrial and failure of intolerance or contraindication to, ≥ 1 of the following agents:<ul style="list-style-type: none">Topical antifungal (e.g., clotrimazole, ketoconazole)Oral griseofulvinOral terbinafine <p>Quantity Limit: 4 per day</p>

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure of an appropriate duration, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of or evidence of organism resistance to generic itraconazole (for systemic infection) or 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Brexafemme ^{CC, QL}	<p>Initial Approval Criteria for Vulvovaginal Candidiasis Treatment:</p> <ul style="list-style-type: none">Patient is a post-menarchal female; ANDDiagnosis of vulvovaginal candidiasis (VVC); ANDFemales of reproductive potential must have a negative pregnancy test; ANDPatient must have an adequate trial and failure, contraindication, resistance, or intolerance of single dose 150 mg oral fluconazole. <p>Renewal Criteria: Cannot be renewed for the same course of infection.</p>

Agent(s) Subject to Criteria	Criteria for Approval
Vivjoa ^{CC, QL}	Initial Approval Criteria for Vulvovaginal Candidiasis Prophylaxis: <ul style="list-style-type: none"> • Patient is a post-menarchal female; AND • Patient has a history of recurrent vulvovaginal candidiasis (RVVC, defined as ≥ 3 culture-confirmed episodes in ≤ 12 months); AND • Used for recurrent vulvovaginal candidiasis prophylaxis; AND • Females of reproductive potential must have negative pregnancy test; AND • Patient must have an adequate trial and failure, contraindication, resistance, or intolerance to oral fluconazole or other triazoles. Renewal Criteria: <ul style="list-style-type: none"> • Females of reproductive potential must have negative pregnancy test; AND • Patient must have a reduction in the recurrence of vulvovaginal candidiasis; AND • Maintenance treatment cannot exceed 6 months of therapy. Quantity Limit: 4 tablets per fill
	Approval Duration: 1 year <ul style="list-style-type: none"> • Patient has diagnosis of recurrent vulvovaginal candidiasis with ≥ 3 episodes of vulvovaginal candidiasis in ≤ 12-month period; AND • Patient is a biological female who is postmenopausal or has another reason for permanent infertility (e.g., tubal ligation, hysterectomy, salpingo-oophorectomy); AND • Patient is not pregnant; AND • Patient is not lactating; AND • Patient has tried and failed or has a contraindication or intolerance to maintenance antifungal therapy with oral fluconazole for 6 months. Quantity Limit: 18 tablets per treatment course

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
clotrimazole troche	Brexafemme ^{CC, QL}
fluconazole suspension, tablet	Cresemba
griseofulvin suspension	Diflucan
itraconazole capsule ^{CC, QL}	flucytosine
ketoconazole	griseofulvin microsize tablet, ultramicrosize tablet
nystatin suspension, tablets	itraconazole solution
terbinafine	Noxafil
	Oravig
	posaconazole
	Sporanox ^{QL}
	Tolsura
	Vfend
	Vivjoa ^{CC, QL}
	voriconazole

ANTI-INFECTIVE: ORAL ANTIVIRALS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agents from the same sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Xofluza ^{AE, CC, QL}	<ul style="list-style-type: none">Confirmed or suspected diagnosis of acute, uncomplicated, outpatient influenza; ANDPatient is symptomatic for ≤ 48 hours; ORPrescribed for post-exposure prevention of influenza after contact with an individual diagnosed with influenza; ANDAllergy, contraindication, intolerance, or other reason a preferred influenza antiviral cannot be used; ANDPatient is not:<ul style="list-style-type: none">Taking concurrent neuraminidase inhibitors (e.g., Tamiflu, Relenza); ORTaking polyvalent cation-containing laxatives, antacids, or oral supplements (e.g., calcium, iron, magnesium, selenium, or zinc); ORPregnant; ORHospitalized. <p>Age Limit: ≥ 5 years Quantity Limit: 2 tablets (1 dose) per fill</p>

CURRENT PDL STATUS

ANTIVIRALS: HERPES

Preferred Agents	Non-Preferred Agents
acyclovir famciclovir	Valtrex Zovirax oral suspension

Preferred Agents	Non-Preferred Agents
Valacyclovir	

ANTIVIRALS: INFLUENZA

Preferred Agents	Non-Preferred Agents
oseltamivir ^{QL}	Flumadine
	Relenza
	rimantadine
	Tamiflu ^{QL}
	Xofluza ^{AE, CC, QL}

ANTI-INFECTIVE: ORAL ANTIBIOTICS

GUIDELINES FOR USE

Approval Duration: Date of Service, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
vancomycin capsules, solution ^{CC}	<ul style="list-style-type: none">• Diagnosis of <i>clostridium difficile</i>-associated diarrhea (ICD-10 = A04.7); OR• Diagnosis of <i>Staphylococcal</i> enterocolitis.
linezolid tablets ^{CC, QL, MD}	<ul style="list-style-type: none">• Completion of a course of therapy begun during a hospital or healthcare facility stay; OR• Diagnosis of methicillin-resistant staph aureus (MRSA), vancomycin-resistant enterococcus (VRE); AND• Prescriber attestation that the choice of therapy is based on culture and sensitivity testing; OR• Trial and failure of another first-line antibiotic in a patient at high risk for complications. <p>Maximum Duration: 28 days Quantity Limit: 2 per day</p> <ul style="list-style-type: none">• NOTE: linezolid suspension may be approved when the above criteria are met AND the member is unable to swallow linezolid tablets.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure of an appropriate course of therapy, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Difidac tablet, suspension ^{CC, QL} fidaxomicin tablet ^{CC, QL}	<p>Approval Duration: Date of Service (total 10-day course of therapy)</p> <ul style="list-style-type: none">• Patient age ≥ 6 months; AND• Diagnosis of pseudomembranous colitis due to <i>C. difficile</i> infection; AND• Trial and failure of vancomycin. <p>Quantity Limits:</p> <ul style="list-style-type: none">• Oral tablets: 2 per day (400mg)

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> 40mg/mL suspension: 10mL per day (400mg)
Firvanq ^{CC}	<ul style="list-style-type: none"> Diagnosis of <i>clostridium difficile</i>-associated diarrhea (ICD-10 = A04.7); OR Diagnosis of <i>Staphylococcal</i> enterocolitis.
metronidazole 125 mg tablet ^{CC}	<ul style="list-style-type: none"> Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to metronidazole 250mg tablet AND metronidazole 500mg tablet.
Solosec ^{AE, CC, QL}	<ul style="list-style-type: none"> Female patient with diagnosis of bacterial vaginosis (BV); AND <ul style="list-style-type: none"> No in vitro resistance to nitroimidazole derivatives (metronidazole, tinidazole, secnidazole) or prior failure of metronidazole or tinidazole for the current course of infection; AND Trial and failure of, or contraindication to, at least one preferred non-nitroimidazole (e.g., clindamycin); OR Female patient with diagnosis of trichomoniasis caused by <i>Trichomonas vaginalis</i>; AND <ul style="list-style-type: none"> No in vitro resistance to nitroimidazole derivatives (metronidazole, tinidazole, secnidazole) or prior failure of metronidazole or tinidazole for the current course of infection; AND History of unacceptable/toxic side effects (not including hypersensitivity reactions) to at least two preferred medications not requiring prior approval. <p>Age Limit: > 12 years Quantity Limit: 1 packet per fill</p>
Nuzyra ^{AE, CC, QL}	<ul style="list-style-type: none"> Diagnosis of community-acquired bacterial pneumonia (CABP) or acute bacterial skin and skin structure infection (ABSSSI) caused by susceptible microorganism(s)*; AND If female of childbearing potential, patient is NOT pregnant; AND Patient is not a candidate (based on culture and sensitivity data) or has failed treatment with ≥ 2 preferred antibiotics from 2 different classes; AND Patient has NOT failed a tetracycline unless susceptibility results demonstrate that pathogen is NOT susceptible to other tetracyclines but is susceptible to omadacycline; AND If continuing an inpatient/hospital treatment course, prescriber attests that it would be clinically inappropriate to deescalate therapy or use alternative therapy based on susceptibility results or lack of susceptibility results in conjunction with clinical picture; AND Total treatment duration will not exceed 14 days per course. <p>Age Limit: ≥ 18 years Quantity Limit: 2 per day; override by call center for loading dose</p> <p>*CABP susceptible microorganisms include: <i>Chlamydia pneumoniae</i>, <i>Haemophilus influenzae</i>, <i>Haemophilus parainfluenzae</i>, <i>Klebsiella pneumoniae</i>, <i>Legionella pneumoniae</i>, <i>Mycoplasma pneumoniae</i>, <i>Staphylococcus aureus</i> (methicillin-susceptible isolates; MSSA), <i>Streptococcus pneumoniae</i>.</p> <p>ABSSSI susceptible microorganisms include: <i>Enterobacter cloacae</i>, <i>Enterococcus faecalis</i>, <i>Klebsiella pneumoniae</i>, <i>Staphylococcus aureus</i> (methicillin-susceptible and -resistant</p>

Agent(s) Subject to Criteria	Criteria for Approval
Vowst ^{AE, CC, QL}	<p>isolates; MSSA and MRSA), <i>Streptococcus lugdunensis</i>, <i>Streptococcus anginosus</i> group (includes <i>S. anginosus</i>, <i>S. intermedius</i>, and <i>S. constellatus</i>), <i>Streptococcus pyogenes</i>.</p> <p>Approval Duration: 30 days (Limit to 1 fill per approval)</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of recurrent <i>Clostridioides difficile</i> infection (CDI); AND • Prescribed by, or in consultation with, a gastroenterologist or infectious disease specialist; AND • Patient has completed at least 3 full courses of antibiotic treatment with two or more of the following guideline recommended agents: <ul style="list-style-type: none"> ○ Vancomycin oral ○ Difcid ○ Metronidazole oral; AND • Treatment with Vowst will be initiated between 48 and 96 hours of completion of the most recent course of antibiotics; AND • At least 8 hours prior to the first dose of Vowst, the patient will receive an appropriate bowel cleansing regimen (e.g., magnesium citrate or polyethylene glycol) <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of recurrent <i>Clostridioides difficile</i> infection (CDI); AND • Prescribed by, or in consultation with, a gastroenterologist or infectious disease specialist; AND • Patient had treatment failure defined as the presence of CDI diarrhea within 8 weeks of the first dose of Vowst AND a positive stool test for <i>C. difficile</i>; AND • Patient has not previously received more than 1 treatment course of Vowst; AND • Previous course of Vowst was at least 12 days ago but no more than 8 weeks ago. <p>Age Limit: ≥ 18 years of age Quantity Limit: 12 capsules over 3 days</p>

CURRENT PDL STATUS

ANTIBIOTICS: CEPHALOSPORINS 1ST GENERATION

Preferred Agents	Non-Preferred Agents
cefadroxil capsule, suspension	cefadroxil tablet
cephalexin capsule, suspension	cephalexin tablet

ANTIBIOTICS: CEPHALOSPORINS 2ND GENERATION

Preferred Agents	Non-Preferred Agents
cefaclor capsule	cefaclor suspension
cefprozil suspension, tablet	cefaclor ER tablet
cefuroxime axetil tablet	

ANTIBIOTICS: CEPHALOSPORINS 3RD GENERATION

Preferred Agents	Non-Preferred Agents
cefdinir capsule, suspension	cefixime capsule, suspension
	cefpodoxime suspension, tablet
	Suprax capsule, chewable tablet, suspension

ANTIBIOTICS: GASTROINTESTINAL

Preferred Agents	Non-Preferred Agents
metronidazole 250 mg, 500 mg tablet	Aemcolo
neomycin	Dificid suspension, tablet ^{CC, QL}
tinidazole	fidaxomicin ^{CC, QL}
vancomycin capsule, solution ^{CC}	Firvanq ^{CC}
	Flagyl
	Likmez
	metronidazole capsule
	metronidazole 125 mg tablet ^{CC}
	nitazoxanide
	Solosec ^{AE, CC, QL}
	Vancocin
	Vowst ^{AE, CC, QL}

ANTIBIOTICS: MACROLIDES

Preferred Agents	Non-Preferred Agents
azithromycin	clarithromycin ER
clarithromycin	E.E.S 400 Filmtab
E.E.S. granules for suspension 200mg	EryPed
erythromycin base capsule DR	Ery-Tab DR 333 mg tablet
erythromycin base tablet DR 250, 500 mg tablet	Erythrocin
erythromycin ethylsuccinate suspension	erythromycin base tablet
Ery-Tab DR 250, 500 mg tablet	erythromycin base tablet DR 333 mg
	erythromycin ethylsuccinate 400 mg tablet
	erythromycin filmtab
	Zithromax

ANTIBIOTICS: OXAZOLIDINONES

Preferred Agents	Non-Preferred Agents
linezolid suspension ^{QL, MD}	Sivextro ^{QL}
linezolid tablet ^{CC, QL, MD}	Zyvox suspension ^{QL, MD}
	Zyvox tablet ^{QL, MD}

ANTIBIOTICS: PENICILLINS

Preferred Agents	Non-Preferred Agents
Amoxicillin	amoxicillin/clavulanate ER
amoxicillin/clavulanate chewable table, suspension, tablet	Augmentin
Ampicillin capsule	Augmentin XR
Dicloxacillin capsule	
penicillin V potassium tablet, suspension	

ANTIBIOTICS: QUINOLONES

Preferred Agents	Non-Preferred Agents
ciprofloxacin tablet	Baxdela ^{AE, QL}
levofloxacin tablet	ciprofloxacin suspension
	Cipro
	levofloxacin solution
	moxifloxacin
	ofloxacin

ANTIBIOTICS: SULFONAMIDES, FOLATE ANTAGONIST

Preferred Agents	Non-Preferred Agents
sulfamethoxazole/trimethoprim	Bactrim
Sulfatrim suspension	Bactrim DS
trimethoprim	sulfadiazine

ANTIBIOTICS: TETRACYCLINES

Preferred Agents	Non-Preferred Agents
demeclocycline	Doryx, Doryx DR, Doryx MPC
doxycycline hyclate	doxycycline hyclate DR
doxycycline monohydrate 50 mg, 100 mg capsule	doxycycline IR-DR
doxycycline monohydrate suspension, tablet	doxycycline monohydrate 40, 75, 150 mg capsule
minocycline capsule	doxycycline "kit" or "pack"
tetracycline capsule	Lymepak
	minocycline tablet
	minocycline ER
	Minolira ER
	Morgidox capsule
	Morgidox Kit
	Nuzyra ^{AE, CC, QL}
	Oracea
	tetracycline tablet
	Vibramycin

ANTI-INFECTIVE: VAGINAL ANTIBIOTICS

GUIDELINES FOR USE

Approval Duration: Date of Service

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial of appropriate duration and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Cleocin Ovule	Cleocin cream
clindamycin vaginal 2% cream	Clindesse vaginal cream
metronidazole vaginal 0.75% gel	Vandazole gel
Nuvessa gel	Xaciato gel

ANTIRETROVIRALS: HUMAN IMMUNODEFICIENCY VIRUS/ACQUIRED IMMUNODEFICIENCY SYNDROME (HIV/AIDS)

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Rukobia ^{AE, CC, QL}	<ul style="list-style-type: none">Diagnosis of human immunodeficiency virus (HIV); ANDPrescribed by, or in consultation with, an infectious disease specialist or HIV specialist (AAHIVS); ANDPrevious treatment with at least 3 drug classes (nucleoside reverse transcriptase inhibitors [NRTI], non-nucleoside reverse transcriptase inhibitors [NNRTI], or protease inhibitor [PI]); ANDDocumentation (e.g., progress note, lab report) of baseline viral load ≥ 400 copies/mL on current antiretroviral regimen; ANDUsed in combination with highly active antiretroviral therapy (HAART); ANDNot used in combination with strong cytochrome P450 (CYP)3A inducers. <p>Renewal Criteria</p> <ul style="list-style-type: none">Documentation (e.g., progress note, lab report) of a decrease in viral load from pretreatment baseline. <p>Age Limit: ≥ 18 years Quantity Limit: 2 per day</p>
Sunlenca tablet ^{AE, CC, QL}	<ul style="list-style-type: none">Diagnosis of human immunodeficiency virus type 1 (HIV-1) infection; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Prescribed by, or in consultation with, an infectious disease specialist or HIV specialist (AAHIVS); AND • Patient is heavily treatment-experienced with multidrug resistance HIV-1 infection (has documented resistance to ≥ 2 antiretroviral [ARV] medications from each of at least 3 of the 4 main classes: nucleoside reverse-transcriptase inhibitors [NRTIs], non-nucleoside reverse-transcriptase inhibitors [NNRTIs], protease inhibitors [PIs], and integrase strand-transfer inhibitors [INSTI]); AND • Patient has ≤ 2 fully active ARVs remaining from the 4 main classes that can be effectively combined; AND • Documentation (e.g., progress note, lab report) of baseline viral load ≥ 400 copies/mL on current antiretroviral regimen; AND • Patient has no history of treatment failure or known or suspected resistance to lenacapavir; AND • Patient will be taking with other antiretrovirals (optimized background regimen); AND • Not used in combination with strong cytochrome CYP3A inducers. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient has been adherent to their ARV treatment regimen; AND • Patient has not experienced virologic failure of lenacapavir and has documented clinical improvement and/or stabilization (e.g., disease response as indicated by a decrease in viral load from pretreatment baseline; increased or stabilized CD4+ counts). <p>Age Limit: ≥ 18 years Quantity Limits: 300 mg tablets: 5 tablets per fill</p>
Vocabria AE, CC, QL	<p>Pre-Exposure Prophylaxis</p> <ul style="list-style-type: none"> • Prescribed for pre-exposure prophylaxis (PrEP) of HIV; AND • Prescriber submits prior authorization request; AND • Used as an oral lead-in for Apretude (cabotegravir extended release injectable suspension) or for oral therapy for patients who will miss a planned injection of Apretude • Prescriber attests that: <ul style="list-style-type: none"> ○ Patient is considered high-risk for HIV infection; AND ○ Risk-reduction and medication adherence counseling were performed; AND ○ Negative HIV-1 test immediately prior to initiating. <p>Treatment of HIV Infection</p> <ul style="list-style-type: none"> • Patient has a diagnosis of human immunodeficiency virus type 1 (HIV-1) infection; AND • Patient is virologically suppressed with HIV-RNA < 50 copies/mL and is on a stable antiretroviral regimen; AND • Patient has no history of treatment failure or known or suspected resistance to cabotegravir or rilpivirine; AND • Patient will take rilpivirine concomitantly for 28 days; AND • Patient will be using cabotegravir as: <ul style="list-style-type: none"> ○ Oral lead-in to assess tolerability of cabotegravir prior to administration of the injectable extended-release formulations of cabotegravir/rilpivirine; OR

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ Oral therapy for patients who plan to miss a dose of their cabotegravir/rilpivirine injection • Patient will NOT receive concomitant therapy with ANY of the following medications that can result in significant decreases of cabotegravir and/or rilpivirine; AND <ul style="list-style-type: none"> ○ Carbamazepine ○ Oxcarbazepine ○ Phenobarbital ○ Phenytoin ○ Rifabutin ○ Rifampin ○ Rifapentine ○ Dexamethasone (more than a single-dose treatment) ○ St. John's wort • Prescribed by or in consultation with an infectious disease specialist or HIV specialist. <p>Age Limit: ≥ 12 years Quantity Limit: 1 per day</p>
Yeztugo tablets (lenacapavir) ^{AE, CC, QL}	<p>Approval Duration: 6 months initial; 12 months renewal</p> <p>Initial Criteria: Pre-Exposure Prophylaxis</p> <ul style="list-style-type: none"> • Prescribed for HIV-1 PrEP <ul style="list-style-type: none"> ○ Day 1 / Day 2 initiation along with injection; OR ○ Temporary oral bridging; OR ○ Supplemental oral dosing when needed for coverage with concomitant CYP3A inducers; AND • Documentation that patient's weight is ≥35 kg • Prior authorization request is submitted by the prescriber; AND • Prescriber attests that patient is considered high-risk for HIV infection; AND • Prescriber attests that risk-reduction and medication adherence counseling were performed; AND • Documentation of negative HIV-1 status immediately prior to initiating Yeztugo using an FDA-approved method. If an antigen/antibody test is used, results should be confirmed with an RNA assay, even if available after the first dose; AND • The requested dose does not exceed the FDA-approved dose for this indication <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Patient continues to meet the above criteria; AND • The requested dose does not exceed the FDA-approved dose for this indication <p>Age Limit: ≥16 years Quantity Limit: 4 tablets per fill; 8 tablets per year</p>
Yeztugo Injection (lenacapavir) ^{AE, CC}	<p>Approval Duration: 6 months initial; 12 months renewal</p> <p>Initial Criteria:</p> <ul style="list-style-type: none"> • Prescribed for pre-exposure prophylaxis (PrEP) of HIV <ul style="list-style-type: none"> ○ For Day 1/Day 2 Initiation along with tablets; OR ○ For ongoing maintenance therapy, AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Documentation that patient's weight is ≥ 35 kg; AND Prior authorization request is submitted by the prescriber; AND Prescriber attests that patient is considered high-risk for HIV infection; AND Prescriber attests that risk-reduction and medication adherence counseling were performed; AND Documentation of negative HIV-1 test immediately prior to each Yeztugo injection, including Day 1/Day 2 initiation (with tablets) and all maintenance doses to confirm HIV-negative status using an FDA-approved method. If an antigen/antibody test is used, results should be confirmed with an RNA assay, even if available after the first dose; AND The requested dose does not exceed the FDA-approved dose for this indication <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Patient continues to meet the above criteria; AND The requested dose does not exceed the FDA-approved dose for this indication. <p>Age Limit: ≥ 16 years</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
abacavir ^{QL}	Aptivus
abacavir-lamivudine	Atripla ^{QL}
atazanvir ^{QL}	Combivir
Biktarvy ^{QL}	Complera ^{QL}
Cimduo ^{QL}	Crixivan
Delstrigo ^{QL}	darunavir
Descovy ^{QL}	didanosine DR ^{QL}
Dovato ^{QL}	efavirenz/lamivudine/tenofovir disoproxil fumarate (generic for Symfi, Symfi Lo) ^{QL}
Edurant Ped tablet suspension	emtricitabine (generic for Emtriva) ^{QL}
Edurant	Epivir ^{QL}
efavirenz (generic for Sustiva)	Epzicom
efavirenz/emtricitabine/tenofovir disoproxil fumarate (generic for Atripla) ^{QL}	etravirine (generic for Intelence)
emtricitabine/tenofovir disoproxil fumarate (generic for Truvada) ^{QL}	fosamprenavir (generic for Lexiva)
emtricitabine/rilpivirine/tenofovir disoproxil fumarate (generic for Complera) ^{QL}	Fuzeon
Emtriva ^{QL}	Invirase
Evotaz ^{QL}	Kaletra solution, tablet
Genvoya ^{QL}	Lexiva
Intelence	Maraviroc (generic for Selzentry)
Isentress	nevirapine ^{QL}
Juluca ^{QL}	nevirapine ER ^{QL}
lamivudine ^{QL}	Norvir solution, tablet, powder packet
lamivudine-zidovudine	Prezcobix ^{QL}
lopinavir-ritonavir solution, tablet	Retrovir
Odefsey ^{QL}	Reyataz ^{QL}
Pifeltro ^{QL}	Rukobia ^{AE, CC, QL}
Prezista	stavudine capsule ^{QL}
ritonavir tablet	Sunlenca ^{AE, CC, QL}
Selzentry	Sustiva
Stribild ^{QL}	Temixys ^{QL}

Preferred Agents	Non-Preferred Agents
Symfi ^{QL}	Tivicay suspension
Symfi Lo ^{QL}	Triumeq suspension
Symtuza ^{QL}	Truvada ^{CC, QL}
tenofovir disoproxil fumarate tablet (generic for Viread) ^{QL}	Viracept
Tivicay tablets ^{QL}	Viramune ^{QL}
Triumeq tablet ^{QL}	Viramune XR ^{QL}
Trizivir	Viread powder packet
Tybost	Viread tablet ^{QL}
zidovudine syrup, tablet	Vocabria ^{AE, CC, QL}
	Yeztugo tablets ^{QL}
	Yeztugo vial
	Ziagen ^{QL}
	zidovudine capsule

HEPATITIS B AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Vemlidy ^{AE, CC, QL}	<ul style="list-style-type: none">Diagnosis of hepatitis B virus infection; ANDPrescribed by, or in consultation with, a hepatology/gastroenterology, infectious disease, transplant or other appropriate specialist; ANDPatient does NOT have decompensated cirrhosis (Child-Pugh B or C); ANDTrial and failure of, intolerance, or contraindication to, entecavir or (tenofovir disoproxil fumarate); ANDNOT concurrently taking any P-gp inducers (e.g., oxcarbazepine, phenobarbital, phenytoin, rifabutin, rifampin, rifapentine, or St. John's wort); ANDIf HIV-1 positive, NOT using Vemlidy as monotherapy. <p>Age Limit: ≥ 18 years Quantity Limit: 1 per day (allow 2 per day for drug interactions)</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Epivir-HBV solution	Adefovir tablet
lamivudine HBV tablet	Baraclude solution, tablet
	Hepsera tablet
	Vemlidy tablet ^{AE, CC, QL}

HEPATITIS C AGENTS: INTERFERONS AND RIBAVIRINS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
PEGASYS syringe, vial ^{CC, QL}	<ul style="list-style-type: none">Prescribed by a gastroenterologist, hepatologist, or infectious disease specialist for the treatment of chronic hepatitis C.Prescribed in combination with interferon or direct-acting antiviral for the treatment of chronic hepatitis C; ANDCriteria for the accompanying product have been met.
ribavirin ^{CC}	

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent in the same sub-class.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable

CURRENT PDL STATUS

HEPATITIS C: INTERFERONS

Preferred Agents	Non-Preferred Agents
PEGASYS syringe, vial ^{CC, QL}	

HEPATITIS C: RIBAVIRINS

Preferred Agents	Non-Preferred Agents
ribavirin capsule, tablet ^{CC}	None

HEPATITIS C AGENTS: DIRECT-ACTING ANTIVIRALS

GUIDELINES FOR USE

Approval Duration: Course of Therapy

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Mavyret ^{CC, QL} sofosbuvir/velpatasvir ^{CC, QL}	<ul style="list-style-type: none">• Simplified HCV Treatment Criteria below are met; OR• HCV Direct-Acting Antiviral Class Criteria (Non-Simplified) below are met; AND• If applicable, Additional Criteria for Patients Previously Treated with a Direct-Acting Antiviral below are met.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires clinical justification (e.g., allergy, contraindication, potential drug-drug interactions with other medications, or intolerance) as to why **Mavyret or sofosbuvir/velpatasvir** cannot be used or are not indicated. **HCV Direct-Acting Antiviral Class Criteria (Non-Simplified) must be met.**

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. TREATMENT CRITERIA

Treatment Criteria Category	Criteria for Approval
Simplified HCV Treatment Criteria (treatment-naïve, non-cirrhotic, uncomplicated cases)	<p>Approval Duration: Mavyret – 12 weeks; sofosbuvir/velpatasvir – 16 weeks</p> <ul style="list-style-type: none">• Diagnosis of chronic hepatitis C virus (HCV) infection; AND• Prescribed regimen is either of the following:<ul style="list-style-type: none">○ Mavyret 12 weeks; OR○ sofosbuvir/velpatasvir for 16 weeks; AND• Documentation (e.g., progress note, prior authorization form questions) of the following clinical data confirming simplified treatment eligibility:<ul style="list-style-type: none">○ Date of Hepatitis C diagnosis or earliest record of HCV infection; AND○ Recent (within 3 months) qualitative or quantitative HCV RNA level (HCV viral load); AND○ NOT pregnant; AND○ NOT previously treated for HCV; AND○ NOT cirrhotic based on FIB-4 score < 3.25 (https://www.heaptitisc.uw.edu/page/clinical-calculators/fib-4); AND○ Human immunodeficiency virus (HIV) negative; AND○ Hepatitis B surface antigen (HBsAg) negative; AND

Treatment Criteria Category	Criteria for Approval
	<ul style="list-style-type: none"> No history of liver transplant or hepatocellular carcinoma.
HCV Direct-Acting Antiviral Class Criteria (non-simplified)	<p>Approval Duration: Full course of treatment (varies by product and clinical factors)</p> <ul style="list-style-type: none"> Diagnosis of acute or chronic hepatitis C virus (HCV) infection; AND Prescribed treatment regimen is included in the requested drug's package insert and/or supported by current HCV guidelines for the patient's age/weight, and other clinical data requested below; AND Prescribed by, a gastroenterologist, hepatologist, infectious disease (including HIV specialist, AAHIVP), or transplant specialist OR prescriber completed/participates in an HCV academic/mentorship training program or network (e.g., KHAMP, ECHO); AND Documentation (e.g., progress note, prior authorization form questions) of the following clinical data: <ul style="list-style-type: none"> Date of acute or chronic HCV diagnosis or earliest record of HCV infection; AND Recent (within 3 months) qualitative or quantitative HCV RNA level (HCV viral load); AND HCV genotype, including subtype and resistance mutations (if known); AND If pregnant, prescriber attests that the benefits of HCV treatment outweigh potential risks to the fetus; AND If applicable, prior HCV treatment regimen(s); AND If cirrhotic, assessment of liver disease severity using the Child-Pugh score; AND Human immunodeficiency virus (HIV) status; AND Hepatitis B surface antigen (HBsAg) status; AND If applicable, any history of liver transplant or hepatocellular carcinoma.

Treatment Criteria Category	Criteria for Approval
Additional Criteria for Patients Previously Treated with a Direct-Acting Antiviral	<ul style="list-style-type: none"> Prescriber must answer the following questions: <ul style="list-style-type: none"> Is retreatment necessary due to treatment failure or reinfection? Was the patient compliant (e.g., few to no missed doses) with previous DAA therapy? If not, why? Were there any additional factors that led to DAA treatment failure? If so, describe these factors and how they have been addressed or are no longer relevant. Patient must be evaluated for alcohol and substance abuse using a validated screening tool; AND If the patient has a recent history (within the past 6 months) of alcohol or substance abuse, the following is required: <ul style="list-style-type: none"> Documentation that the patient has completed or is participating in a recovery program, receiving alcohol or substance abuse counseling services, or seeing an addiction specialist as part of HCV treatment; AND Documentation that the patient is not actively participating in illicit substance use or alcohol abuse with confirmatory laboratory testing (e.g., urine drug screen); AND Provider attests that they believe:

Treatment Criteria Category	Criteria for Approval
	<ul style="list-style-type: none"> ○ Patient is willing and able to comply with the requirements of the proposed retreatment plan; AND ○ Any factors that may have led to noncompliance with previous treatment(s) have been addressed; AND ○ Patient has received education regarding risk behaviors (e.g., IV drug use) associated with HCV infection.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Mavyret ^{AE, CC, QL}	Epclusa ^{AE, CC, QL}
sofosbuvir/velpatasvir ^{AE, CC, QL}	Harvoni ^{AE, CC, QL}
	ledipasvir/sofosbuvir ^{AE, CC, QL}
	Sovaldi ^{AE, CC, QL}
	Viekira Pak ^{AE, CC, QL}
	Vosevi ^{AE, CC, QL}
	Zepatier ^{AE, CC, QL}

DIABETES: INSULINS AND RELATED AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 3 month** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **2 preferred agents** in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Admelog vial and Solostar ^{CC}	<ul style="list-style-type: none">Trial and failure of ≥ 2 preferred insulins; ANDClinical rationale (e.g., intolerance to an inactive ingredient) that a preferred product cannot be used.
Fiasp vial, pump cartridge, and FlexTouch ^{CC} Lyumjev pen, Tempo Pen, and vial ^{CC}	<ul style="list-style-type: none">Trial and failure of ≥ 2 preferred insulins; ANDClinical rationale (e.g., intolerance to an inactive ingredient) that a preferred product cannot be used.
Symlin ^{AE, CC}	<ul style="list-style-type: none">Prescribed by, or in consultation with, an endocrinologist or other diabetes specialist; ANDTrial and failure of ≥ 1 rapid-acting insulin. <p>Age Limit: ≥ 18 years</p>
Basaglar KwikPen ^{CC} insulin glargine-yfgn pen and vial ^{CC} Semglee (yfgn) pen and vial ^{CC}	<ul style="list-style-type: none">Trial and failure of ≥ 2 preferred insulins, one of which must be insulin glargine or Lantus; ANDClinical rationale (e.g., intolerance to an inactive ingredient) that a preferred product cannot be used.

CURRENT PDL STATUS

RAPID- AND SHORT-ACTING INSULINS

Preferred Agents	Non-Preferred Agents
Humulin R vial	Admelog and Admelog Solostar ^{CC}
Humulin R U-500 KwikPen and vial	Afrezza
insulin aspart cartridge, pen, and vial	Apidra Solostar and vial
insulin lispro Junior (Jr) KwikPen	Fiasp FlexTouch, pen, pumpcart, and vial ^{CC}

Preferred Agents	Non-Preferred Agents
insulin lispro pen and vial	Humalog 200 unit/mL KwikPen
	Humalog cartridge, KwikPen, and vial
	Humalog Junior (Jr) KwikPen
	Humalog Tempo Pen
	Kirsty vial and pen
	Lyumjev pen, Tempo Pen, and vial ^{CC}
	Merilog Solostar and vial
	Novolin R pen and vial
	Novolog cartridge, FlexPen, and vial
	Symlin ^{AE, CC}

INTERMEDIATE-ACTING INSULINS

Preferred Agents	Non-Preferred Agents
Humalog Mix 50/50 KwikPen	Humalog Mix 75/25 KwikPen and vial
Humulin 70/30 KwikPen and vial	Humulin N KwikPen
Humulin N vial	Novolin 70/30 pen and vial
insulin aspart/insulin aspart protamine pen and vial	Novolin N pen
insulin lispro/insulin lispro protamine KwikPen	Novolog Mix vial
insulin lispro protamine mix	
Novolin N vial	
Novolog Mix FlexPen	

LONG-ACTING INSULINS

Preferred Agents	Non-Preferred Agents
insulin glargine Solostar U100 (generic for Lantus Solostar)	Basaglar KwikPen, Tempo Pen ^{CC}
insulin glargine vial	insulin degludec pen and vial
Lantus and Lantus Solostar	Insulin glargine Solostar and Max Solostar (generic for Toujeo)
Levemir FlexPen, FlexTouch, and vial	insulin glargine-yfgn pen and vial ^{CC}
	Rezvoglar Kwikpen
	Semglee (yfgn) pen and vial ^{CC}
	Toujeo Solostar and Max Solostar
	Tresiba FlexTouch and vial

DIABETES: GLUCAGON-LIKE PEPTIDE-1 (GLP-1) RECEPTOR AGONISTS

GUIDELINES FOR USE

Approval Duration: 6 months

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Byetta ^{CC, QL} Ozempic ^{AE, CC, QL} Trulicity ^{CC, QL} Victoza ^{CC, QL}	Initial Approval Criteria <ul style="list-style-type: none"> Diagnosis of Type II Diabetes Mellitus (T2DM) confirmed with clinical documentation including: <ul style="list-style-type: none"> ICD-10 diagnosis of T2DM (chart notes within the past 12 months) ; AND <ul style="list-style-type: none"> A1c lab value that correlates to a T2DM diagnosis (i.e., 6.5 or greater); OR Historical A1c that correlates to a T2DM diagnosis (i.e. 6.5 or greater), AND A1C within the past 6 months; AND No personal or family history of medullary thyroid carcinoma (MTC) or Multiple Endocrine Neoplasia syndrome type 2 (MEN 2); AND Not used in combination with another GLP-1 receptor agonist OR DPP4 UNLESS the member is changing therapy; AND The requested dose does not exceed the maximum FDA-approved dose for the treatment of diabetes mellitus. Renewal Criteria <ul style="list-style-type: none"> ICD-10 diagnosis of T2DM (chart notes within the past 12 months) AND Clinical documentation must be submitted demonstrating an A1c value within the past 6 months; AND The provider attests that the patient has been evaluated for safety (e.g. lacks treatment limiting adverse events) and demonstrates a positive response to therapy. AND No personal or family history of medullary thyroid carcinoma (MTC) or Multiple Endocrine Neoplasia syndrome type 2 (MEN 2); AND Not used in combination with another GLP-1 receptor agonist OR DPP4 UNLESS the member is changing therapy; AND The requested dose does not exceed the maximum FDA-approved dose for treating diabetes mellitus. <p>*Drugs used for anorexia, weight loss, or weight gain are excluded from coverage</p>

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires documentation of a **≥ 3 month** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **2 preferred agent**, unless otherwise specified. Additional drug specific criteria must also be met.

Agent(s) Subject to Criteria	Criteria for Approval
Mounjaro Rybelsus	Initial Approval Criteria <ul style="list-style-type: none"> Diagnosis of Type II Diabetes Mellitus (T2DM) confirmed with

Agent(s) Subject to Criteria	Criteria for Approval
	<p>clinical documentation including:</p> <ul style="list-style-type: none"> ○ ICD-10 diagnosis of T2DM (chart notes within the past 12 months) ; AND <ul style="list-style-type: none"> ▪ A1c lab value that correlates to a T2DM diagnosis (i.e., 6.5 or greater); OR ▪ Historical A1c that correlates to a T2DM diagnosis (i.e. 6.5 or greater), AND A1C within the past 6 months; AND • No personal or family history of medullary thyroid carcinoma (MTC) or Multiple Endocrine Neoplasia syndrome type 2 (MEN 2); AND • Not used in combination with another GLP-1 receptor agonist OR DPP4 UNLESS the member is changing therapy; AND • The requested dose does not exceed the maximum FDA-approved dose for the treatment of diabetes mellitus. • A ≥ 3-month trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified. (chart notes or claim history must confirm) <p>Renewal Criteria</p> <ul style="list-style-type: none"> • ICD-10 diagnosis of T2DM (chart notes within the past 12 months) AND • Demonstrate ONE of the following: <ul style="list-style-type: none"> ○ an A1c of less than or equal to 8% within the past 6 months OR ○ demonstration of improved A1c value OR ○ provider must submit clinical justification for continued therapy • The provider attests that the patient has been evaluated for safety (e.g. lacks treatment limiting adverse events) and demonstrates a positive response to therapy. AND • No personal or family history of medullary thyroid carcinoma (MTC) or Multiple Endocrine Neoplasia syndrome type 2 (MEN 2); AND • Not used in combination with another GLP-1 receptor agonist OR DPP4 UNLESS the member is changing therapy; AND • The requested dose does not exceed the maximum FDA-approved dose for treating diabetes mellitus. <p>*Drugs used for anorexia, weight loss, or weight gain are excluded from coverage</p>

3. BRAND MEDICALLY NECESSARY CRITERIA

Not applicable.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. ADDITIONAL DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
exenatide ^{CC, QL} liraglutide ^{CC, QL}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none"> • Diagnosis of Type II Diabetes Mellitus (T2DM) confirmed with

Agent(s) Subject to Criteria	Criteria for Approval
	<p>clinical documentation including:</p> <ul style="list-style-type: none"> • ICD-10 diagnosis of T2DM (chart notes within the past 12 months) ; AND <ul style="list-style-type: none"> ▪ A1c lab value that correlates to a T2DM diagnosis (i.e., 6.5 or greater); OR ▪ Historical A1c that correlates to a T2DM diagnosis (i.e. 6.5 or greater), AND A1C within the past 6 months; AND • No personal or family history of medullary thyroid carcinoma (MTC) or Multiple Endocrine Neoplasia syndrome type 2 (MEN 2); AND • Not used in combination with another GLP-1 receptor agonist OR DPP4 UNLESS the member is changing therapy; AND • The requested dose does not exceed the maximum FDA-approved dose for the treatment of diabetes mellitus. • A ≥ 3-month trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified. (chart notes or claim history must confirm) • A known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation <p>Renewal Criteria</p> <ul style="list-style-type: none"> • ICD-10 diagnosis of T2DM (chart notes within the past 12 months) AND • Demonstrate ONE of the following: <ul style="list-style-type: none"> ○ an A1c of less than or equal to 8% within the past 6 months OR ○ demonstration of improved A1c value OR ○ provider must submit clinical justification for continued therapy • The provider attests that the patient has been evaluated for safety (e.g. lacks treatment limiting adverse events) and demonstrates a positive response to therapy. AND • No personal or family history of medullary thyroid carcinoma (MTC) or Multiple Endocrine Neoplasia syndrome type 2 (MEN 2); AND • Not used in combination with another GLP-1 receptor agonist OR DPP4 UNLESS the member is changing therapy; AND • The requested dose does not exceed the maximum FDA-approved dose for treating diabetes mellitus. • A known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation <p>*Drugs used for anorexia, weight loss, or weight gain are excluded from coverage</p>
<p>Soliqua^{AE, CC, QL} Xultophy^{AE, CC, QL}</p>	<ul style="list-style-type: none"> • Trial and failure (e.g., non-compliance, need to reduce injections) of a long-acting insulin (e.g. insulin glargine) and a GLP-1 agonist (e.g., Victoza) used concurrently AND • Not used in combination with a DPP-4 inhibitors (such as Janumet or Tradjenta) UNLESS the member is changing therapy and the DPP-4 inhibitor medication will be discontinued <p>Age Limit: ≥ 18 years</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Byetta ^{CC, QL}	Bydureon BCise ^{QL}
Ozempic ^{AE, CC, QL}	exenatide ^{CC, QL}
Trulicity ^{CC, QL}	liraglutide ^{CC, QL}
Victoza ^{CC, QL}	Mounjaro ^{AE, QL}
	Rybelsus ^{CC, QL}
	Soliqua ^{AE, CC, QL}
	Xultophy ^{AE, CC, QL}

DIABETES: DIPEPTIDYL PEPTIDASE-4 (DPP-4) INHIBITORS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Janumet ^{CC, QL} Janumet XR ^{CC, QL} Januvia ^{CC, QL} Jentadueto ^{CC, QL} Jentadueto XR ^{CC, QL} Nesina ^{CC, QL} Tradjenta ^{CC, QL}	<ul style="list-style-type: none">• Diagnosis of Type II Diabetes Mellitus (without chronic kidney disease); AND<ul style="list-style-type: none">◦ Trial and failure (e.g., A1c goal not met) of, intolerance, or contraindication to metformin; OR• Diagnosis of Type II Diabetes Mellitus (with chronic kidney disease (ICD-10 Group N18)); AND<ul style="list-style-type: none">◦ Trial and failure of, intolerance, or contraindication to ≥ 1 SGLT2 inhibitor plus metformin. AND• Not used in combination with a GLP-1 receptor (such as Ozempic or Mounjaro) UNLESS the member is changing therapy and the GLP medication will be discontinued.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires ≥ 3 month trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 **preferred agent**, unless otherwise specified. Preferred with PA (PDP) criteria must be met.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
sitagliptin/metformin ER ^{CC, QL} Sitagliptin/Metformin ^{CC, QL} Zituvimet ^{CC, QL} Zituvimet XR ^{CC, QL}	<ul style="list-style-type: none">• Diagnosis of Type II Diabetes Mellitus (without chronic kidney disease); AND<ul style="list-style-type: none">◦ Trial and failure (e.g., A1c goal not met) of, intolerance, or contraindication to metformin; OR• Diagnosis of Type II Diabetes Mellitus (with chronic kidney disease (ICD-10 Group N18)); AND<ul style="list-style-type: none">◦ Trial and failure of, intolerance, or contraindication to ≥ 1 SGLT2 inhibitor plus metformin; AND• Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in Janumet or Janumet XR; AND• At least 3-month trial and failure, allergy,

Agent(s) Subject to Criteria	Criteria for Approval
	<p>contraindication (including potential drug-drug interactions with other medications), or intolerance to 1 preferred agent AND</p> <ul style="list-style-type: none"> Not used in combination with a GLP-1 receptor (such as Ozempic or Mounjaro) UNLESS the member is changing therapy and the GLP medication will be discontinued.
Zituvio ^{CC, QL}	<ul style="list-style-type: none"> Diagnosis of Type II Diabetes Mellitus (without chronic kidney disease); AND <ul style="list-style-type: none"> Trial and failure (e.g., A1c goal not met) of, intolerance, or contraindication to metformin; OR Diagnosis of Type II Diabetes Mellitus (with chronic kidney disease (ICD-10 Group N18)); AND <ul style="list-style-type: none"> Trial and failure of, intolerance, or contraindication to ≥ 1 SGLT2 inhibitor plus metformin; AND Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in Januvia; AND Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with at least 2 manufacturers (if available) of the corresponding generic; AND At least 3-month trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications), or intolerance to 1 preferred agent AND Not used in combination with a GLP-1 receptor (such as Ozempic or Mounjaro) UNLESS the member is changing therapy and the GLP medication will be discontinued.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Janumet ^{CC, QL}	alogliptin ^{QL}
Janumet XR ^{CC, QL}	alogliptin/metformin ^{QL}
Januvia ^{CC, QL}	alogliptin/pioglitazone ^{QL}
Jentadueto ^{CC, QL}	Brynovin ^{QL}
Jentadueto XR ^{CC, QL}	Glyxambi ^{QL}
linagliptin/metformin ^{QL}	Kazano ^{QL}
Nesina ^{CC, QL}	Kombiglyze XR ^{QL}
Tradjenta ^{CC, QL}	Onglyza ^{QL}
	Oseni ^{QL}
	Qtern ^{QL}
	saxagliptin ^{QL}
	saxagliptin/metformin ER ^{QL}
	sitagliptin ^{QL}
	sitagliptin/metformin ER ^{CC, QL}
	sitagliptin/metformin ^{CC, QL}
	Steglujan ^{AE, QL}
	Trijardy XR ^{QL}
	Zituvio ^{CC, QL}
	Zituvimet ^{CC, QL}
	Zituvimet XR ^{CC, QL}

DIABETES: SODIUM-GLUCOSE COTRANSPORTER-2 (SGLT₂) INHIBITORS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Synjardy ^{CC, QL}	<ul style="list-style-type: none">• Diagnosis of Type 2 Diabetes Mellitus; AND<ul style="list-style-type: none">○ Diagnosis of chronic kidney disease (ICD-10 Group N18); OR○ Diagnosis of atherosclerotic cardiovascular disease (ASCVD); OR○ Diagnosis of heart failure with reduced ejection fraction; OR○ Trial and failure (e.g., A1c goal not met) of, intolerance or contraindication to metformin.
Farxiga ^{CC, QL} Jardiance ^{CC, QL}	<ul style="list-style-type: none">• Diagnosis of Type 2 Diabetes Mellitus; AND<ul style="list-style-type: none">○ Diagnosis of atherosclerotic cardiovascular disease (ASCVD); OR○ Trial and failure (e.g., A1c goal not met) of, intolerance or contraindication to metformin; OR• Diagnosis of chronic kidney disease (ICD-10 Group N18); OR• Diagnosis of heart failure.
Xigduo XR ^{CC, QL}	<ul style="list-style-type: none">• Diagnosis of Type 2 Diabetes Mellitus; AND<ul style="list-style-type: none">○ Diagnosis of atherosclerotic cardiovascular disease (ASCVD); OR○ Trial and failure (e.g., A1c goal not met) of, intolerance or contraindication to metformin; OR• Diagnosis of heart failure with reduced ejection fraction.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 3 month** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent, unless otherwise specified. Preferred with PA (PDP) criteria must be met.

3. BRAND MEDICALLY NECESSARY CRITERIA

Not applicable.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Inpefa ^{CC, AE, QL}	<ul style="list-style-type: none">• Diagnosis of Type 2 Diabetes Mellitus; AND• Diagnosis of chronic kidney disease; AND• Patient has other cardiovascular risk factors; OR• Diagnosis of heart failure; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Patient has had ≥ 3 month trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.
	Quantity Limit: 30 tablets per 30 days

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Farxiga ^{CC, QL}	dapagliflozin ^{QL}
Jardiance ^{CC, QL}	dapagliflozin-metformin ER ^{QL}
Synjardy ^{CC, QL}	Inpefa ^{CC, AE, QL}
Xigduo XR ^{CC, QL}	Invokamet ^{CC, QL}
	Invokamet XR ^{QL}
	Invokana ^{CC, QL}
	Segluromet ^{AE, QL}
	Steglatro ^{AE, QL}
	Synjardy XR ^{QL}

DIABETES: ALPHA-GLUCOSIDASE INHIBITORS

GUIDELINES FOR USE

Approval Duration: 1 year

1. **PREFERRED WITH PA (PDP) CRITERIA**

Not applicable. All preferred agents are preferred without PA.

2. **NON-PREFERRED (NPD) CRITERIA**

Approval of non-preferred agents requires **≥ 3 month** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **1 preferred agent**.

3. **BRAND MEDICALLY NECESSARY CRITERIA**

Approval of NON-PREFERRED branded agents requires:

- a. Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. **GENERIC MEDICALLY NECESSARY CRITERIA**

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. **DRUG-SPECIFIC CLINICAL CRITERIA**

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
acarbose ^{QL}	miglitol ^{QL} Precose ^{QL}

DIABETES: METFORMINS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 3 month** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **2 preferred agents** (e.g., metformin IR and metformin ER).

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
metformin solution ^{CC} Riomet ^{CC} Riomet ER ^{CC}	<ul style="list-style-type: none">Unable to swallow metformin or metformin ER tablets.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
glyburide/metformin	glipizide/metformin
metformin 500 mg, 850 mg, 1000 mg tablet	metformin ER tablet (generic Fortamet, Glumetza)
metformin ER tablet (generic Glucophage XR)	metformin solution ^{CC}
	metformin 625 mg, 750 mg tablet
	Riomet solution ^{CC}
	Riomet ER suspension ^{CC}

DIABETES: MEGLITINIDES

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 3 month** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **1 preferred agent**.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
nateglinide ^{QL}	
repaglinide ^{QL}	

DIABETES: SULFONYLUREAS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **2 preferred agents**.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Glimepiride	Glucotrol XL
Glipizide	
glipizide ER	
glipizide XL	
Glyburide	
glyburide micronized	

DIABETES: THIAZOLIDINEDIONES (TZDS)

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **1 preferred agent**.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
pioglitazone ^{QL}	Actoplus Met ^{QL}
	Actos ^{QL}
	Duetact ^{QL}
	pioglitazone/glimepiride ^{QL}
	pioglitazone/metformin ^{QL}

ENDOCRINE AND METABOLIC AGENTS: GLUCAGON AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Baqsimi ^{CC}	<ul style="list-style-type: none">Intramuscular (IM) glucagon was dispensed in the past 180 days; ORPrescriber attestation that caregiver(s) or family member(s) would have or have had difficulty preparing and administering the IM injection in a correct and timely manner.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure **within the past 180 days**, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **1 preferred agent**.

3. BRAND MEDICALLY NECESSARY CRITERIA

Not applicable.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Baqsimi spray ^{CC}	diazoxide suspension
Gvoke autoinjector, syringe	Glucagon emergency kit
Proglycem suspension	Gvoke vial
Zegalogue autoinjector, syringe ^{AE}	

ENDOCRINE AND METABOLIC AGENTS: GROWTH HORMONES

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Genotropin ^{CC} Norditropin Flexpro ^{CC}	<ul style="list-style-type: none"> • Diagnosis (documented or reported) of one of the following conditions: <ul style="list-style-type: none"> ○ Hypofunction and other disorders of the pituitary gland (ICD-10 = E23.x); OR ○ Short stature due to endocrine disorder (ICD-10 = E34.3) or idiopathic short stature (ICD-10 = R62.52); OR ○ Post-procedural (iatrogenic) hypopituitarism (ICD-10 = E89.3); OR ○ Neoplasm of pituitary or craniopharyngeal duct (ICD-10 = C75.1, C75.2, D35.2, D35.3, D44.3, D44.4); OR ○ Turner's syndrome (ICD-10 = Q96); OR ○ Congenital malformation syndromes (e.g., Noonan syndrome, Prader-Willi syndrome) predominantly associated with short stature (ICD-10 = Q87.1); OR ○ End-stage renal disease (ICD-10 = N18.5, N18.6, N18.9); OR ○ Newborn light for gestational age (ICD-10 = P05.0y); OR ○ Cachexia (ICD-10 = R64). <p><i>x = a blank value or a number 1-7 that completes an ICD-10 code. ICD-10 Disease Group = E23</i></p> <p><i>y = value of 0-8 (based on member weight) that completes an ICD-10 code.</i></p> <ul style="list-style-type: none"> • For all diagnoses except HIV-associated wasting or cachexia (ICD-10: B22.2 or R64) and short bowel syndrome (ICD-10: K90.82), documentation or attestation of open epiphyses is required when appropriate.
Skytrofa ^{AE, CC}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none"> • Patient has growth failure secondary to growth hormone deficiency (GHD); AND • Patient must have tried and failed 1 preferred short-acting growth hormone products due to frequency of administration or adherence. • For all diagnoses except HIV-associated wasting or cachexia (ICD-10: B22.2 or R64) and short bowel syndrome (ICD-10: K90.82), documentation or attestation of open epiphyses is required when appropriate. • Patient must meet the minimum age recommended by the package insert for the FDA-approved indication <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient continues to meet the above criteria; AND • Patient has a positive response compared to pre-treatment

Agent(s) Subject to Criteria	Criteria for Approval
	baseline.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **2 preferred agents**. Preferred with PA (PDP) criteria must be met.

For all diagnoses except HIV-associated wasting or cachexia (ICD-10: B22.2 or R64) and short bowel syndrome (ICD-10: K90.82), documentation or attestation of open epiphyses is required when appropriate.

3. BRAND MEDICALLY NECESSARY CRITERIA

Not applicable.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Ngenla ^{CC, AE}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none"> • Diagnosis of growth hormone deficiency; AND • Pediatric patient must NOT have closed epiphyses if used for longitudinal growth promotion; AND • Patient does NOT have active malignancy; AND • Patient does NOT have active proliferative or severe non-proliferative diabetic retinopathy; AND • Patient does NOT have Prader-Willi syndrome with ≥ 1 of the following: <ul style="list-style-type: none"> ○ Severe obesity ○ History of upper airway obstruction or sleep apnea ○ Severe respiratory impairment ○ Unidentified respiratory infection; AND • Trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications), or intolerance of 2 preferred agents. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient continues to meet the above criteria; AND • Patient has a positive response compared to pre-treatment baseline. <p>Age Limit: ≥ 3 years</p>
Sogroya ^{CC, QL}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none"> • Patient will be at least 2.5 years old at the start of treatment; AND • Diagnosis of growth hormone deficiency; AND • Patient does NOT have a hypersensitivity to any somapacitan product or any of the excipients; AND • Pediatric patient must NOT have closed epiphyses if used for longitudinal growth promotion; AND • Patient does NOT have active malignancy; AND • Patient does NOT have active proliferative or severe non-proliferative diabetic retinopathy; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Patient does NOT have, or previously had, an intracranial tumor growth as confirmed by a sellar MRI scan with contrast; AND • Patient does NOT have Prader-Willi syndrome with > 1 of the following: <ul style="list-style-type: none"> • Severe obesity • History of upper airway obstruction or sleep apnea • Severe respiratory impairment • Unidentified respiratory infection; AND • Trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications), or intolerance of 2 preferred agents <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient continues to meet the above criteria; AND • Patient has not had unacceptable toxicity from the drug; AND • Patient has a positive response compared to pre-treatment baseline <p>Quantity Limit: 4 pens per 28 days</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Genotropin cartridge, syringe ^{CC}	Humatrope cartridge
Norditropin FlexPro ^{CC}	Ngenla ^{CC, AE}
Skytrofa cartridge ^{CC}	Nutropin AQ NuSpin ^{CC}
	Omnitrope cartridge, vial
	Serostim vial
	Sogroya ^{CC, QL}
	Zomacton vial

ENDOCRINE AND METABOLIC AGENTS: GLUCOCORTICOIDS, ORAL (ORAL STEROIDS)

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **2 preferred agents**, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
budesonide DR capsule ^{QL}	Alkindi Sprinkle capsule
budesonide EC capsule ^{QL}	Cortef tablet
dexamethasone elixir, solution, tablet	cortisone acetate tablet
hydrocortisone tablet	dexamethasone dose pack, Intensol drop
methylprednisolone dose pack, 4 mg, 32 mg tablet	Hemady tablet
prednisolone solution	Khindivi solution ^{AE}
prednisolone sodium phosphate solution 5 mg/5 mL, 15 mg/5 mL, 25 mg/5 mL	Medrol dose pack, tablet
prednisone dose pack, solution, tablet	methylprednisolone 8 mg, 16 mg tablet
	Millipred dose pack, tablet
	prednisolone tablet
	prednisolone sodium phosphate ODT, solution 10 mg/5 mL, 20 mg/5 mL
	prednisone Intensol oral concentrate
	Rayos DR tablet
	TaperDex dose pack
	Tarpeyo DR capsule

ENDOCRINE AND METABOLIC AGENTS: PANCREATIC ENZYMES

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **1 preferred agent**.

3. BRAND MEDICALLY NECESSARY CRITERIA

Not applicable.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Creon capsule	Pertzye capsule
Viokace tablet	
Zenpep capsule	

ENDOCRINE AND METABOLIC AGENTS: PROGESTINS FOR CACHEXIA

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2 manufacturers** (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agent
megestrol acetate 40 mg/mL suspension ^{QL} , tablet	megestrol acetate 625 mg/5 mL suspension

ENDOCRINE AND METABOLIC AGENTS: ANDROGENIC AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **2 preferred agents**.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Androgel gel pump	Androgel gel packet
testosterone gel pump (generic Androgel)	Natesto nasal pump
	Testim gel
	testosterone gel (generic Testim, Vogelxo)
	testosterone gel packet (generic Androgel)
	testosterone gel pump (generic Axiron, Fortesta, Vogelxo)
	Testosterone solution
	Vogelxo gel, gel packet, gel pump

ENDOCRINE AND METABOLIC AGENTS: BONE RESORPTION SUPPRESSION AND RELATED AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Forteo AE, CC, QL	<ul style="list-style-type: none"> Diagnosis one of the following: <ul style="list-style-type: none"> Female with postmenopausal osteoporosis, OR Male with primary or hypogonadal osteoporosis, OR Female or male with osteoporosis associated with systemic glucocorticoid therapy; AND Documented hip DXA (femoral neck or total hip) or lumbar spine T-score ≤ -2.5 standard deviations; AND Patient is at a high risk for fractures; AND Patient is not at increased risk for osteosarcoma (e.g., Paget's disease of bone, bone metastases or skeletal malignancies, etc.); AND Patient has not received therapy with parathyroid hormone analogs (e.g., abaloparatide, teriparatide) for 24 months or more (lifetime cumulative); AND Documented allergy, intolerance, contraindication, or therapeutic failure to a minimum 12-month trial of the following: <ul style="list-style-type: none"> Oral or IV bisphosphonate drug (e.g., alendronate, ibandronate, or risedronate); AND raloxifene tablets (Evista); OR calcitonin injections (Miacalcin) <p>Renewal Criteria</p> <ul style="list-style-type: none"> Documentation of disease response (e.g., absence of fractures); AND Total length of therapy will not exceed 24 months (lifetime cumulative). <p>Age Limit: ≥ 18 years of age Quantity Limit: 20 mcg per day</p>
Jubbonti AE, CC, QL	<p>Initial Approval Criteria:</p> <p>Osteoporosis</p> <ul style="list-style-type: none"> Patient is male; OR Patient is female and postmenopausal; AND Diagnosis of osteoporosis, confirmed by ONE of the following: <ul style="list-style-type: none"> Documented lumbar spine or total hip T-score ≤ -2.5 standard deviations; OR Documented lumbar spine or total hip T-score of -1.0 to -2.5 and patient is at a high risk for fractures based on FRAX score; OR Documented lumbar spine or total hip T-score of -1.0 to -2.5 and patient has a fragility fracture of humerus, pelvis, or forearm; AND Documented allergy, intolerance, contraindication, or therapeutic failure to a minimum 12-month trial of oral or IV

Agent(s) Subject to Criteria	Criteria for Approval
	<p>bisphosphonate drugs (e.g., alendronate, ibandronate, or risedronate); AND</p> <ul style="list-style-type: none"> • Provider attestation that any pre-existing hypocalcemia has been reviewed and corrected prior to initiation of denosumab therapy; AND • Prescriber attestation that the patient has been instructed to supplement with at least 1000 mg calcium and at least 400 IU vitamin D daily. <p>Bone Loss Related to Non-Metastatic Prostate Cancer</p> <ul style="list-style-type: none"> • Diagnosis of non-metastatic prostate cancer; AND • Patient is receiving androgen deprivation therapy (ADT) with ONE of the following: <ul style="list-style-type: none"> ◦ Luteinizing hormone-releasing hormone (LHRH) or gonadotropin releasing hormone (GnRH) agonist (e.g., Lupron); OR ◦ Bilateral orchiectomy; AND • Diagnosis of secondary osteoporosis related to ADT, confirmed by ONE of the following: <ul style="list-style-type: none"> ◦ Documented lumbar spine or total hip T-score ≤ -1.0 standard deviations; OR ◦ History of fragility fracture; AND • Documented allergy, intolerance, contraindication, or therapeutic failure to a minimum 12-month trial of oral or IV bisphosphonate drugs (e.g., alendronate, ibandronate, or risedronate); AND • Provider attestation that any pre-existing hypocalcemia has been reviewed and corrected prior to initiation of denosumab therapy; AND • Prescriber attestation that the patient has been instructed to supplement with at least 1000 mg calcium and at least 400 IU vitamin D daily. <p>Bone Loss Related to Adjuvant Aromatase Inhibitor Therapy for Breast Cancer</p> <ul style="list-style-type: none"> • Diagnosis of breast cancer; AND • Patient is receiving adjuvant aromatase inhibitor therapy (AIT) (e.g., anastrozole, exemestane, letrozole); AND • Diagnosis of secondary osteoporosis related to AIT, confirmed by ONE of the following: <ul style="list-style-type: none"> ◦ Documented lumbar spine or total hip T-score ≤ -1.0 standard deviations; OR ◦ History of fragility fracture; AND • Documented allergy, intolerance, contraindication, or therapeutic failure to a minimum 12-month trial of oral or IV bisphosphonate drugs (e.g., alendronate, ibandronate, or risedronate); AND • Provider attestation that any pre-existing hypocalcemia has been reviewed and corrected prior to initiation of denosumab therapy; AND • Prescriber attestation that the patient has been instructed to supplement with at least 1000 mg calcium and at least 400 IU vitamin D daily. <p>Osteoporosis Associated with Systemic Glucocorticoid Therapy</p> <ul style="list-style-type: none"> • Patient is receiving ONE of the following:

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ Glucocorticoid treatment of ≥ 7.5 mg per day of prednisone (or equivalent) and is expected to maintain treatment for at least 6 months; OR ○ Glucocorticoid treatment of ≥ 30 mg per day; OR ○ Cumulative glucocorticoid treatment of ≥ 5 g per year; AND • Diagnosis of secondary osteoporosis related to glucocorticoid therapy; AND • Patient is at a high risk for fracture, confirmed by ONE of the following: <ul style="list-style-type: none"> ○ Documented lumbar spine or total hip T-score ≤ -2.5 standard deviations; OR ○ Documented lumbar spine or total hip T-score of -1.0 to -2.5 and patient is at a high risk for fractures based on FRAX score; OR ○ Documented lumbar spine or total hip T-score of -1.0 to -2.5 and patient has a fragility fracture of humerus, pelvis, or forearm; AND • Documented allergy, intolerance, contraindication, or therapeutic failure to a minimum 12-month trial of oral or IV bisphosphonate drugs (e.g., alendronate, ibandronate, or risedronate); AND • Provider attestation that any pre-existing hypocalcemia has been reviewed and corrected prior to initiation of denosumab therapy; AND • Prescriber attestation that the patient has been instructed to supplement with at least 1000 mg calcium and at least 400 IU vitamin D daily. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Patient must continue to meet initial approval criteria; AND • Documentation of positive disease response to therapy (e.g., improved or stabilized BMD, absence of fractures). <p>Age Limit: ≥ 18 years of age Quantity Limit: 60 mg every 6 months</p>

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires ≥ 12 month trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **2 preferred agents**.

Agent(s) Subject to Criteria	Criteria for Approval
Prolia ^{AE, CC, QL}	<p>Initial Approval Criteria:</p> <p>Osteoporosis</p> <ul style="list-style-type: none"> • Patient is male; OR • Patient is female and postmenopausal; AND • Diagnosis of osteoporosis, confirmed by ONE of the following: <ul style="list-style-type: none"> ○ Documented lumbar spine or total hip T-score ≤ -2.5 standard deviations; OR ○ Documented lumbar spine or total hip T-score of -1.0 to -2.5 and patient is at a high risk for fractures based on FRAX score; OR ○ Documented lumbar spine or total hip T-score of -1.0 to -2.5 and patient has a fragility fracture of humerus, pelvis, or forearm; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Documented allergy, intolerance, contraindication, or therapeutic failure to a minimum 12-month trial of oral or IV bisphosphonate drugs (e.g., alendronate, ibandronate, or risedronate); AND Provider attestation that any pre-existing hypocalcemia has been reviewed and corrected prior to initiation of denosumab therapy; AND Prescriber attestation that the patient has been instructed to supplement with at least 1000 mg calcium and at least 400 IU vitamin D daily; AND Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to Jubbonti. <p>Bone Loss Related to Non-Metastatic Prostate Cancer</p> <ul style="list-style-type: none"> Diagnosis of non-metastatic prostate cancer; AND Patient is receiving androgen deprivation therapy (ADT) with ONE of the following: <ul style="list-style-type: none"> Luteinizing hormone-releasing hormone (LHRH) or gonadotropin releasing hormone (GnRH) agonist (e.g., Lupron); OR Bilateral orchiectomy; AND Diagnosis of secondary osteoporosis related to ADT, confirmed by ONE of the following: <ul style="list-style-type: none"> Documented lumbar spine or total hip T-score ≤ -1.0 standard deviations; OR History of fragility fracture; AND Documented allergy, intolerance, contraindication, or therapeutic failure to a minimum 12-month trial of oral or IV bisphosphonate drugs (e.g., alendronate, ibandronate, or risedronate); AND Provider attestation that any pre-existing hypocalcemia has been reviewed and corrected prior to initiation of denosumab therapy; AND Prescriber attestation that the patient has been instructed to supplement with at least 1000 mg calcium and at least 400 IU vitamin D daily; AND Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to Jubbonti. <p>Bone Loss Related to Adjuvant Aromatase Inhibitor Therapy for Breast Cancer</p> <ul style="list-style-type: none"> Diagnosis of breast cancer; AND Patient is receiving adjuvant aromatase inhibitor therapy (AIT) (e.g. anastrozole, exemestane, letrozole); AND Diagnosis of secondary osteoporosis related to AIT, confirmed by ONE of the following: <ul style="list-style-type: none"> Documented lumbar spine or total hip T-score ≤ -1.0 standard deviations; OR History of fragility fracture; AND Documented allergy, intolerance, contraindication, or therapeutic failure to a minimum 12-month trial of oral or IV bisphosphonate drugs (e.g., alendronate, ibandronate, or risedronate); AND Provider attestation that any pre-existing hypocalcemia has been reviewed and corrected prior to initiation of denosumab therapy; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Prescriber attestation that the patient has been instructed to supplement with at least 1000 mg calcium and at least 400 IU vitamin D daily; AND • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to Jubbonti. <p>Osteoporosis Associated with Systemic Glucocorticoid Therapy</p> <ul style="list-style-type: none"> • Patient is receiving ONE of the following: <ul style="list-style-type: none"> ○ Glucocorticoid treatment of ≥ 7.5 mg per day of prednisone (or equivalent) and is expected to maintain treatment for at least 6 months; OR ○ Glucocorticoid treatment of ≥ 30 mg per day; OR ○ Cumulative glucocorticoid treatment of ≥ 5 g per year; AND • Diagnosis of secondary osteoporosis related to glucocorticoid therapy; AND • Patient is at a high risk for fracture, confirmed by ONE of the following: <ul style="list-style-type: none"> ○ Documented lumbar spine or total hip T-score ≤ -2.5 standard deviations; OR ○ Documented lumbar spine or total hip T-score of -1.0 to -2.5 and patient is at a high risk for fractures based on FRAX score; OR ○ Documented lumbar spine or total hip T-score of -1.0 to -2.5 and patient has a fragility fracture of humerus, pelvis, or forearm; AND • Documented allergy, intolerance, contraindication, or therapeutic failure to a minimum 12-month trial of oral or IV bisphosphonate drugs (e.g., alendronate, ibandronate, or risedronate); AND • Provider attestation that any pre-existing hypocalcemia has been reviewed and corrected prior to initiation of denosumab therapy; AND • Prescriber attestation that the patient has been instructed to supplement with at least 1000 mg calcium and at least 400 IU vitamin D daily; AND • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to Jubbonti <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Patient must continue to meet initial approval criteria; AND • Documentation of positive disease response to therapy (e.g., improved or stabilized BMD, absence of fractures). <p>Age Limit: ≥ 18 years of age Quantity Limit: 60 mg every 6 months</p>

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2 manufacturers** (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Evenity ^{AE, CC, QL}	<ul style="list-style-type: none">Documented intolerance, contraindication or treatment failure/ineffective response to a minimum 12-month trial on previous therapy with teriparatide.
teriperatide ^{AE, CC, QL}	<ul style="list-style-type: none">Diagnosis of one of the following:<ul style="list-style-type: none">Female with postmenopausal osteoporosis, ORMale with primary or hypogonadal osteoporosis, ORFemale or male with osteoporosis associated with systemic glucocorticoid therapy; ANDDocumented hip DXA (femoral neck or total hip) or lumbar spine T-score ≤ -2.5 (standard deviations); ANDPatient is at a high risk for fractures; ANDPatient is not at increased risk for osteosarcoma (e.g., Paget's disease of bone, bone metastases or skeletal malignancies, etc.); ANDPatient has not received therapy with parathyroid hormone analogs (e.g., abaloparatide, teriparatide) for 24 months or more (lifetime cumulative); ANDDocumented allergy, intolerance, contraindication, or therapeutic failure to a minimum 12-month trial of the following:<ul style="list-style-type: none">oral or IV bisphosphonate drug (e.g., alendronate, ibandronate, or risedronate), ANDraloxifene tablets (Evista), ORcalcitonin injections (Miacalcin); ANDPatient has a known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. <p>Renewal Criteria</p> <ul style="list-style-type: none">Documentation of disease response (e.g., absence of fractures); ANDTotal length of therapy will not exceed 24 months (lifetime cumulative); ANDPatient has a known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. <p>Age Limit: ≥ 18 years of age Quantity Limit: 20 mcg per day</p>
Tymlos ^{AE, CC, QL}	<ul style="list-style-type: none">Documented intolerance, contraindication or treatment failure/ineffective response to a minimum 12-month trial on previous therapy with teriparatide.

CURRENT PDL STATUS

Preferred ORAL and NASAL Agents	Non-Preferred ORAL and NASAL Agents
alendronate tablet ^{QL}	Actonel tablet ^{QL}
ibandronate tablet	Alendronate oral solution ^{QL}
raloxifene tablet	Atelvia DR tablet ^{QL}
	Binosto tablet ^{QL}
	Boniva tablet ^{QL}
	calcitonin-salmon nasal spray, vial
	Evista tablet
	Fosamax tablet ^{QL}
	Fosamax Plus D tablet ^{QL}
	risedronate sodium tablet ^{QL}
	Risedronate DR tablet ^{QL}

Preferred INJECTABLE Agents	Non-Preferred Agents
Forteo pen ^{AE, CC, QL}	Bildyos
Jubbonti syringe ^{AE,CC,QL}	calcitonin-salmon vial
	Conexxence
	Evenity syringe ^{AE, CC, QL}
	Miacalcin vial
	Prolia syringe
	Reclast solution
	Stoboclo syringe
	teriperatide pen ^{AE, CC, QL}
	Tymlos pen ^{AE, CC, QL}
	zoledronic acid bag, bottle, vial

ENDOCRINE AND METABOLIC AGENTS: UTERINE DISORDER TREATMENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **1 preferred agent** with the same indication for use.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2 manufacturers** (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Myfembree	
Oriahnn	
Orilissa	

IMMUNOSUPPRESSANTS

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **2 preferred agents**.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2 manufacturers** (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Rezurock ^{AE, CC, QL}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none">• Patient is post-allogeneic stem cell transplants (generally 3 or more months); AND• Patient has diagnosis of chronic graft-versus-host disease (cGVHD); AND• Patient does not have histologic relapse of underlying cancer or post-transplant lymphoproliferative disease; AND• Patient has had a trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents; AND• Will be used in combination with stable doses of systemic therapies for GVHD which must include, but are not limited to, corticosteroids, calcineurin inhibitors (cyclosporine; tacrolimus), sirolimus, mycophenolate mofetil, methotrexate, or rituximab; AND• Belumosudil will not be used in combination with ibrutinib (subsequent therapy is allowed). <p>Renewal Criteria</p> <ul style="list-style-type: none">• Patient continues to meet the above criteria; AND• Patient has not had unacceptable toxicity from the drug (e.g., grade 4 hepatotoxicity); AND• Patient has had a positive response to therapy. <p>Age Limit: ≥ 12 years old Quantity Limit: 1 per day</p>

Agent(s) Subject to CriteriaTavneos^{AE, CC, QL}**Criteria for Approval****Approval Duration: 6 months initial, 1 year renewal****Initial Approval Criteria**

- Patient has severe active antineutrophil cytoplasmic autoantibody (ANCA)-associated vasculitis; **AND**
 - Patient has autoantibodies for proteinase 3 (PR3) or myeloperoxidase (MPO), as detected using indirect immunofluorescence (IIF) assay or antigen-specific enzyme linked immunosorbent assays (ELISAs); **OR**
 - Disease is confirmed by tissue biopsy at the site of active disease; **AND**
- Patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment; **AND**
- Physician has assessed disease severity utilizing an objective measure/tool (e.g., Birmingham Vasculitis Activity Score [BVAS]) and patient has a baseline score of ≥ 16 with 1 of the following:
 - Patient has 1 major item; **OR**
 - Patient has ≥ 3 non-major items; **OR**
 - Patient has ≥ 2 renal items of proteinuria and hematuria; **AND**
- Patient does NOT have an active infection, including clinically important localized infections; **AND**
- Patient has failed on ≥ 1 of the following regimens:
 - Patient has failed immunosuppressant therapy (e.g., cyclophosphamide, azathioprine, methotrexate, mycophenolate), unless contraindicated or intolerant; **OR**
 - Patient has failed on anti-CD20 monoclonal antibody therapy (e.g., rituximab), unless contraindicated or intolerant; **AND**
- Avacopan (Tavneos) will be used as adjunctive therapy in combination with standard therapy (e.g., corticosteroids, cyclophosphamide, azathioprine, mycophenolate, rituximab).

Renewal Criteria

- Disease response from pre-treatment baseline as indicated by the following:
 - Absence of new symptoms; **AND**
 - Minimal use of glucocorticoids (e.g., < 5 mg of prednisone or equivalent); **AND**
 - One or more of the following:
 - Decrease in relapses/flares and/or ANCA levels; **OR**
 - Improvement in organ manifestations (e.g., those with pulmonary renal syndrome should improve in PFTs, proteinuria, creatinine); **OR**
 - Remission (defined as a composite scoring index of 0 on the BVAS)

Age Limit: ≥ 18 years**Quantity Limit: 6 capsules per day****CURRENT PDL STATUS**

Preferred Agents	Non-Preferred Agents
azathioprine tablet	Astagraf XL capsule
CellCept suspension	CellCept capsule, tablet
cyclosporine capsule, modified capsule, modified solution	Envarsus XR tablet
cyclosporine modified	

Preferred Agents	Non-Preferred Agents
Gengraf capsule, solution	everolimus tablet
mycophenolate mofetil capsule, tablet	Imuran tablet
mycophenolic acid tablet	mycophenolate mofetil suspension
sirolimus solution, tablet	Myfortic DR tablet
tacrolimus capsule	Myhibbin suspension
	Neoral capsule, solution
	Prograf capsule, gran pack
	Rapamune solution, tablet
	Rezurock tablet ^{AE, CC, QL}
	Sandimmune capsule, solution
	Tavneos capsule ^{AE, CC, QL}
	Zortress tablet

IMMUNOLOGIC AND GENETIC: MULTIPLE SCLEROSIS AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Avonex ^{CC, QL} Betaseron ^{CC, QL} Copaxone 20 mg ^{CC, QL} dimethyl fumarate ^{CC, QL} fingolimod ^{CC, QL} Kesimpta ^{AE, CC, QL}	<ul style="list-style-type: none">• Diagnosis of multiple sclerosis (ICD-10 Disease Group G35). <ul style="list-style-type: none">• Initially prescribed by or in consultation with a neurologist or multiple sclerosis specialists; AND• Diagnosis of a relapsing form of multiple sclerosis (MS): relapsing-remitting MS (RRMS) active secondary progressive MS (SPMS), or clinically isolated syndrome (CIS); AND• Patient has had an inadequate response to, or unable to tolerate, 1 or more preferred MS agent; AND• NOT have active Hepatitis B, or other clinically significant active infection; AND• Baseline serum immunoglobulin measurement has been or will be performed; AND• NOT used in combination with any other MS agent; AND• Patient does NOT have current systemic or clinically significant local infection. Renewal Criteria <ul style="list-style-type: none">• Documentation of response to therapy (e.g., progress note); AND• Documentation (e.g., lab results) of ongoing serum immunoglobulin monitoring.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 1 month** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **2 preferred agents**, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2 manufacturers** (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Gilenya ^{CC, QL}	<ul style="list-style-type: none"> • Diagnosis of multiple sclerosis (ICD-10 Disease Group G35); • Patient has had an inadequate response to, or is unable to tolerate, 2 or more preferred MS agent; AND • Patient has had a trial and failure (e.g., allergy or intolerance to an inactive ingredient) with at least TWO manufacturers (if available) of the corresponding generic.
Mavenclad ^{AE, CC, QL}	<p>Approval Duration: 35 days initial; one 35-day renewal</p> <ul style="list-style-type: none"> • Initially prescribed by or in consultation with a neurologist or multiple sclerosis specialists; AND • Diagnosis of a relapsing form of multiple sclerosis (MS), relapsing-remitting MS (RRMS), or active secondary progressive MS (SPMS); AND • Patient has had an inadequate response to, or is unable to tolerate, 2 or more preferred MS agents; AND • Not used in combination with any other MS agent; AND • Patient does not meet any of the following conditions: <ul style="list-style-type: none"> ○ Human immunodeficiency virus (HIV), hepatitis B or C, or tuberculosis (TB) infection; ○ Current cancer or malignancy; ○ Current systemic, or clinically significant local, infection; ○ Use of any other antineoplastic, immunosuppressive or immunomodulator drugs to treat other conditions; <p>AND</p> <ul style="list-style-type: none"> • Patient has had or will have all of the following: <ul style="list-style-type: none"> ○ Screening for hepatitis B/C, HIV, and TB infections; AND ○ Testing for antibodies to the varicella zoster virus (VZV) OR have received immunization for VZV at least 4 to 6 weeks prior to beginning therapy; AND ○ Baseline MRI \leq 3 months before initiating the first treatment course; AND ○ For women of childbearing potential, a negative pregnancy test and counseling on contraception use during therapy. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • At least 43 weeks has/will have elapsed since the end of the first treatment course; AND • Continue to meet initial approval criteria; AND • Documentation of response to therapy (e.g., progress note).
Mayzent ^{AE, CC, QL}	<ul style="list-style-type: none"> • Initially prescribed by or in consultation with a neurologist or multiple sclerosis specialist; AND • Patient has a diagnosis of a relapsing form of multiple sclerosis (MS): relapsing-remitting MS (RRMS) active secondary progressive MS (SPMS), or clinically isolated syndrome (CIS); AND • Patient has had an inadequate response to, or is unable to tolerate, 2 or more preferred MS agents; AND • NOT used in combination with another MS agent; AND • Patient does not meet any of the following conditions: <ul style="list-style-type: none"> ○ Presence of contraindicated cardiovascular comorbidities (e.g., recent heart attack or stroke, heart failure); ○ Current systemic or clinically significant local infection; ○ Use of any other antineoplastic, immunosuppressive or immunomodulating drugs to treat other conditions; ○ Prior use of alemtuzumab; AND • Patient has had or will have all of the following: <ul style="list-style-type: none"> ○ CYP2C9 variant genotyping testing to guide dosing; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ Screening for clinically significant drug interactions; AND ○ Baseline electrocardiogram (ECG), liver function tests (LFTs) and ophthalmic evaluation; AND ○ If pre-existing non-contraindicated cardiac disease (e.g., arrhythmia), cardiology consultation and follow-up will be conducted prior to and during treatment; AND ○ Testing for antibodies to the varicella zoster virus (VZV) OR have received immunization for VZV at least 4 to 6 weeks prior to beginning therapy. <p>Renewal Criteria</p> <ul style="list-style-type: none"> ● Continue to meet initial approval criteria; AND ● Documentation of response to therapy (e.g., progress note).
Ponvory AE, CC, QL	<ul style="list-style-type: none"> ● Initially prescribed by or in consultation with a neurologist or multiple sclerosis specialists; AND ● Patient has a diagnosis of a relapsing form of multiple sclerosis (MS): relapsing-remitting MS (RRMS) active secondary progressive MS (SPMS), or clinically isolated syndrome (CIS); AND ● Patient has had an inadequate response to, or is unable to tolerate, 2 or more preferred MS agents; AND ● NOT used in combination with another MS agent; AND ● Patient has a baseline heart rate (HR) ≥ 55 beats per minute (bpm); AND ● If patient is of child-bearing potential, patient is taking effective contraception; AND ● Patient does not meet any of the following conditions: <ul style="list-style-type: none"> ○ Presence of contraindicated cardiovascular comorbidities (e.g., recent heart attack or stroke, heart failure); ○ Presence of Mobitz Type II second- or third-degree atrioventricular (AV) block, sick sinus syndrome, or sinoatrial block (unless treated with a functioning pacemaker); ○ Current systemic or clinically significant local infection; ○ Moderate to severe hepatic impairment (Child-Pugh B or C); ○ Use of any other antineoplastic, immunosuppressive or immunomodulating drugs to treat other conditions; ○ Prior use of alemtuzumab; AND ● Patient has had or will have all of the following: <ul style="list-style-type: none"> ○ Screening for clinically significant drug interactions; AND ○ Baseline electrocardiogram (ECG), liver function tests (LFTs) and ophthalmic evaluation; AND ○ Monitoring of respiratory function in patients with baseline respiratory conditions (e.g., pulmonary fibrosis, asthma, chronic obstructive pulmonary disease); AND ○ If pre-existing non-contraindicated cardiac disease (e.g., arrhythmia), cardiology consultation and follow-up will be conducted prior to and during treatment; AND ○ Testing for antibodies to the varicella zoster virus (VZV) OR have received immunization for VZV at least 4 weeks prior to beginning therapy. <p>Renewal Criteria</p> <ul style="list-style-type: none"> ● Continue to meet initial approval criteria; AND ● Documentation of response to therapy (e.g., progress note).
Rebif CC, QL	<ul style="list-style-type: none"> ● Diagnosis of multiple sclerosis (ICD-10 Disease Group G35).
Zeposia AE, CC, QL	Multiple Sclerosis:

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Initially prescribed by or in consultation with a neurologist or multiple sclerosis specialists; AND Patient has a diagnosis of a relapsing form of multiple sclerosis (MS): relapsing-remitting MS (RRMS) active secondary progressive MS (SPMS), or clinically isolated syndrome (CIS); AND Patient has had an inadequate response to, or is unable to tolerate, 2 or more preferred MS agents; AND NOT used in combination with another MS agent; AND Patient does not meet any of the following conditions: <ul style="list-style-type: none"> Presence of contraindicated cardiovascular comorbidities (e.g., recent heart attack or stroke, heart failure); Current systemic or clinically significant local infection; Use of any other antineoplastic, immunosuppressive or immunomodulating drugs to treat other conditions; Prior use of alemtuzumab; AND Patient has had or will have all of the following: <ul style="list-style-type: none"> Screening for clinically significant drug interactions; AND Baseline electrocardiogram (ECG), liver function tests (LFTs) and ophthalmic evaluation; AND If pre-existing non-contraindicated cardiac disease (e.g., arrhythmia), cardiology consultation and follow-up will be conducted prior to and during treatment; AND Testing for antibodies to the varicella zoster virus (VZV) OR have received immunization for VZV at least 4 to 6 weeks prior to beginning therapy.
	<p>Renewal Criteria</p> <ul style="list-style-type: none"> Continue to meet initial approval criteria; AND Documentation of response to therapy (e.g., progress note).
	<p>Ulcerative Colitis:</p> <p>Approval Duration: 6 months initial; 1 year renewal</p> <ul style="list-style-type: none"> Diagnosis of moderate to severely active ulcerative colitis (UC); AND Prescribed by or in consultation with a gastroenterologist or other specialist in the treatment of UC; AND Patient does not meet any of the following conditions: <ul style="list-style-type: none"> Presence of contraindicated cardiovascular comorbidities (e.g., recent heart attack or stroke, heart failure); Current systemic or clinically significant local infection; Use of any other antineoplastic, immunosuppressive or immunomodulating drugs to treat other conditions; Prior use of alemtuzumab; AND Patient has had or will have all of the following: <ul style="list-style-type: none"> CYP2C9 variant genotyping testing to guide dosing; AND Screening for clinically significant drug interactions; AND Baseline electrocardiogram (ECG), liver function tests (LFTs) and ophthalmic evaluation; AND If pre-existing non-contraindicated cardiac disease (e.g., arrhythmia), cardiology consultation and follow-up will be conducted prior to and during treatment; AND Testing for antibodies to the varicella zoster virus (VZV) OR have received immunization for VZV at least 4 to 6 weeks prior to beginning therapy; AND Patient has trial and failure (at least 3 months) of ≥ 1 of the following conventional therapies: <ul style="list-style-type: none"> Oral/rectal 5-aminosalicylic acid agents (e.g., balsalazide, Lialda, mesalamine, sulfasalazine); OR

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ Oral/rectal steroids (e.g., budesonide, hydrocortisone, prednisone); OR ○ Immunosuppressant (e.g., azathioprine, mercaptopurine); OR • Patient is deemed high-risk for intestinal complications or post-operative recurrence; AND • NOT used in combination with any other biologic agent; AND • Patient has had a ≥ 3 month trial and therapeutic failure, allergy, contraindication (including potential drug-drug interaction) or intolerance to a preferred anti-TNF therapy indicated for ulcerative colitis.
	Renewal Criteria <ul style="list-style-type: none"> • Documentation of response to therapy (e.g., progress note).

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Avonex ^{CC, QL}	Ampyra tablet ^{QL}
Avonex pen ^{QL} , syringe ^{QL} , syringe kit ^{QL}	Aubagio tablet ^{QL}
Betaseron ^{CC, QL}	Bafiertam capsule ^{AE, QL}
Betaseron kit ^{QL} , vial ^{QL}	Copaxone 40 mg syringe ^{QL}
Copaxone 20 mg syringe ^{CC, QL}	Extavia kit ^{QL} , vial ^{QL}
dalfampridine ER tablet ^{QL}	Gilenya capsule ^{CC, QL}
dimethyl fumarate DR capsule ^{CC, QL}	glatiramer acetate syringe ^{QL}
fingolimod capsule ^{CC, QL}	Glatopa syringe ^{QL}
Kesimpta pen ^{AE, CC, QL}	Mavenclad tablet ^{AE, CC, QL}
teriflunomide tablet ^{QL}	Mayzent tablet ^{AE, CC, QL} , tablet dose pack ^{AE, CC, QL}
	Plegridy pen ^{QL} , syringe ^{QL}
	Ponvory tablet ^{AE, CC, QL} , tablet dose pack ^{AE, CC, QL}
	Rebif Rebidose autoinjector ^{CC, QL}
	Rebif syringe ^{CC, QL}
	Tascenso ODT ^{QL}
	Tecfidera capsule ^{QL}
	Vumerity capsule ^{AE, QL}
	Zeposia capsule ^{AE, CC, QL}

IMMUNOLOGIC AND GENETIC: CYTOKINE AND CAM ANTAGONISTS

GUIDELINES FOR USE

Approval Duration: 6 months initial, 1 year renewal

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Cosentyx ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Ankylosing Spondylitis (AS) and nonradiographic axial spondylarthritis (nr-axSpA) Clinical Criteria • Hidradenitis Suppurativa (HS) Clinical Criteria • Plaque Psoriasis Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria <p>OR</p> <ul style="list-style-type: none"> • Diagnosis of active enthesitis-related arthritis <ul style="list-style-type: none"> ◦ Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Juvenile Psoriatic Arthritis Criteria:</p> <p>Initial Criteria</p> <ul style="list-style-type: none"> • Diagnosis of juvenile psoriatic arthritis; AND • Prescribed by, or in consultation with, a rheumatologist or other specialist in the treatment of juvenile psoriatic arthritis; AND • At least 2 years of age; AND • Trial and failure (at least 3 months), contraindication or intolerance to, ≥ 1 disease-modifying anti-rheumatic drug (DMARD), such as methotrexate; AND • NOT used in combination with any other biologic agent; AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Documentation (e.g., progress note) of response to therapy.
Enbrel ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Ankylosing Spondylitis (AS) Clinical Criteria • Juvenile Idiopathic Arthritis (JIA) Clinical Criteria • Plaque Psoriasis Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria • Rheumatoid Arthritis (RA) Clinical Criteria
Humira (and adalimumab biosimilars) ^{CC, QL} Hadlima ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Ankylosing Spondylitis (AS) and nonradiographic axial spondylarthritis (nr-axSpA) Clinical Criteria • Crohn's Disease (CD) or Ulcerative Colitis (UC) Clinical Criteria • Hidradenitis Suppurativa (HS) Clinical Criteria • Juvenile Idiopathic Arthritis (JIA) Clinical Criteria • Plaque Psoriasis Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria • Rheumatoid Arthritis (RA) Clinical Criteria

Agent(s) Subject to Criteria	Criteria for Approval
<p>Otezla^{CC, QL}</p> <p>Otezla XR^{CC, QL}</p>	<ul style="list-style-type: none"> Uveitis Clinical Criteria <p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> Psoriatic Arthritis (PsA) Clinical Criteria <p>Plaque Psoriasis Criteria:</p> <p>Initial Criteria</p> <ul style="list-style-type: none"> Diagnosis of plaque psoriasis (mild, moderate, or severe); AND Prescribed by, or in consultation with, a dermatologist, rheumatologist or other specialist in the treatment of psoriasis; AND Trial and failure (at least 3 months) of ≥ 1 conventional therapy: <ul style="list-style-type: none"> Disease-modifying anti-rheumatic drug (DMARD), such as methotrexate Immunosuppressant (e.g., cyclosporine) Oral retinoid (e.g., acitretin); AND NOT used in combination with any other biologic agent; AND Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> Documentation (e.g., progress note) of response to therapy compared to baseline, such as redness, thickness, scaliness, amount of surface area involvement, and/or PASI score. <p>Behcet's Disease Criteria:</p> <p>Initial Criteria</p> <ul style="list-style-type: none"> Prescribed for the treatment of oral ulcers in a patient with Behcet's disease; AND Prescribed by, or in consultation with a rheumatologist or other specialist in the treatment of <i>Behçet's Disease</i>; AND Patient must meet the minimum age recommended by the package insert for this FDA-approved indication; AND Trial and failure of ≥ 1 of the following conventional therapies <ul style="list-style-type: none"> Azathioprine Sulfasalazine Colchicine Topical or oral steroids (e.g., triamcinolone, prednisone) <p>Renewal Criteria</p> <ul style="list-style-type: none"> Documentation (e.g., progress note) of response to therapy compared to baseline.
<p>Rinvoq^{AE, CC, QL}</p>	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> Ankylosing Spondylitis (AS) or nonradiographic axial spondylarthritis (nr-axSpA) Clinical Criteria Crohn's Disease (CD) Clinical Criteria Giant Cell Arthritis (GCA) Clinical Criteria Juvenile Idiopathic Arthritis (JIA) Clinical Criteria Psoriatic Arthritis (PsA) Clinical Criteria Rheumatoid Arthritis (RA) Clinical Criteria Ulcerative Colitis (UC) Clinical Criteria <p>Treatment of Refractory, Moderate-to-Severe Atopic Dermatitis Initial Criteria:</p>

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Patient has moderate-to-severe atopic dermatitis (AD) defined by ≥ 1 of the following: <ul style="list-style-type: none"> ◦ Involvement of $\geq 10\%$ of body surface area (BSA); OR ◦ Eczema Area and Severity Index (EASI) score of ≥ 16; OR ◦ Investigator's Global Assessment (IGA) score of ≥ 3; OR ◦ Scoring Atopic Dermatitis (SCORAD) score of ≥ 25; OR ◦ Pruritus Numerical Rating Scale (NRS) score of ≥ 4; OR ◦ Incapacitation due to AD lesion location (head and neck, palms, soles, or genitalia); AND • Prescribed by, or in consultation with, a dermatologist, allergist/immunologist, or other specialist in the treatment of atopic dermatitis; AND • Trial and failure, contraindication or intolerance to, ≥ 1 agent in each of the following categories: <ul style="list-style-type: none"> ◦ Topical corticosteroid of medium to high potency (e.g., mometasone, fluocinolone) unless inappropriate for the location (e.g., face, groin); AND ◦ Topical calcineurin inhibitor (i.e., tacrolimus or pimecrolimus); AND ◦ Immunomodulating systemic agent (e.g., cyclosporine, azathioprine, methotrexate, mycophenolate mofetil, etc.); AND • Not used in combination with other JAK inhibitors, biologic immunomodulators, or with other immunosuppressants; AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Documentation (e.g., progress note) of response to therapy relative to baseline measure(s) (e.g., BSA involvement, EASI, IGA, SCORAD).
Rinvoq LQ ^{AE, CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Juvenile Idiopathic Arthritis (JIA) Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria <p>Age Limit: 2-17 years of age Quantity Limit: 360 mL per 30 days</p>
Tyenne ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Juvenile Idiopathic Arthritis (JIA) Clinical Criteria • Rheumatoid Arthritis (RA) Clinical Criteria <p>Giant Cell Arteritis (GCA) Criteria:</p> <ul style="list-style-type: none"> • Prescribed by a rheumatologist, vascular medicine, or other specialist in the diagnosis and treatment of GCA; AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Quantity Limit: 162 mg (1 pen or 1 syringe) per week</p>
Xeljanz ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Ankylosing Spondylitis (AS) or nonradiographic axial spondyloarthritis (nr-axSpA) Clinical Criteria • Juvenile Idiopathic Arthritis (JIA) Clinical Criteria

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Psoriatic Arthritis (PsA) Clinical Criteria • Rheumatoid Arthritis (RA) Clinical Criteria • Ulcerative Colitis (UC) Clinical Criteria

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 3 month** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **1 preferred agent**, unless otherwise specified (e.g., no preferred agents are indicated). Biosimilar agents must also meet PREFERRED WITH PA (PDP) OR DRUG-SPECIFIC CRITERIA for the reference product.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. Trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2 manufacturers** (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. ANKYLOSING SPONDYLITIS (AS) OR NONRADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA) CLINICAL CRITERIA

- Diagnosis of Ankylosing Spondylitis (AS) or nonradiographic axial spondyloarthritis (nr-axSpA); **AND**
- Prescribed by, or in consultation with, a rheumatologist or other specialist in the treatment of AS/nr-axSpA; **AND**
- Trial and failure of, contraindication or intolerance to, **≥ 1** non-steroidal anti-inflammatory drug (NSAID); **AND**
- NOT used in combination with any other biologic agent; **AND**
- For non-preferred agents: 3-month trial and failure of, contraindication or intolerance to, **≥ 1** preferred cytokine or CAM antagonist indicated for the treatment of this condition; **AND**
- Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.

Renewal Criteria:

- Documentation (e.g., progress note) of response to therapy.

6. CROHN'S DISEASE (CD) OR ULCERATIVE COLITIS (UC) CLINICAL CRITERIA

- Diagnosis of Crohn's disease (CD) or Ulcerative Colitis (UC); **AND**
- Patient is 18 years or older, and medication is prescribed by, or in consultation with, a gastroenterologist or other specialist in the treatment of CD/UC; **AND**
- Trial and failure of **≥ 1** of the following conventional therapies:
 - Oral/rectal 5-aminosalicylic acid agents (e.g., balsalazide, Lialda, mesalamine, sulfasalazine)

- Oral/rectal steroids (e.g., budesonide, hydrocortisone, prednisone)
- Immunosuppressant (e.g., azathioprine, mercaptopurine); **OR**
- Member is less than 18 years old and prescriber is a **pediatric** gastroenterologist/CD/UC specialist; **AND**
- Patient is deemed high-risk for intestinal complications or post-operative recurrence; **AND**
- NOT used in combination with any other biologic agent; **AND**
- For non-preferred agents: 3-month trial and failure of, contraindication or intolerance to, ≥ 1 preferred cytokine or CAM antagonist indicated for the treatment of this condition; **AND**
- Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.

Renewal Criteria:

- Documentation (e.g., progress note) of response to therapy.

7. HIDRADENITIS SUPPURATIVA (HS) CLINICAL CRITERIA

- Diagnosis of moderate to severe hidradenitis suppurativa (HS); **AND**
- Prescribed by, or in consultation with, a dermatologist, rheumatologist or other specialist in the treatment of HS; **AND**
- Trial and failure (at least 3 months) of ≥ 1 non-biologic therapy:
 - Contraceptives (e.g., ethinyl estradiol/norgestimate)
 - Oral retinoid (e.g., acitretin)
 - Systemic antibiotic (e.g., clindamycin, minocycline, doxycycline, rifampin) ; **AND**
- NOT used in combination with any other biologic agent; **AND**
- For non-preferred agents: **3-month** trial and failure of, contraindication or intolerance to, ≥ 1 preferred cytokine or CAM antagonist indicated for the treatment of this condition; **AND**
- Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.

Renewal Criteria:

- Documentation (e.g., progress note) of response to therapy.

8. JUVENILE IDIOPATHIC ARTHRITIS (JIA) CLINICAL CRITERIA

- Diagnosis of active polyarticular or systemic juvenile idiopathic arthritis (JIA); **AND**
- Prescribed by, or in consultation with, a rheumatologist or other specialist in the treatment of JIA; **AND**
- Trial and failure (at least 3 months), contraindication or intolerance to, ≥ 1 disease-modifying anti-rheumatic drug (DMARD), such as methotrexate; **AND**
- NOT used in combination with any other biologic agent; **AND**
- For non-preferred agents: 3-month trial and failure of, contraindication or intolerance to, ≥ 1 preferred cytokine or CAM antagonist indicated for the treatment of this condition; **AND**
- Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.

Renewal Criteria:

- Documentation (e.g., progress note) of response to therapy.

9. PLAQUE PSORIASIS CLINICAL CRITERIA

- Diagnosis of moderate to severe plaque psoriasis; **AND**
- Prescribed by, or in consultation with, a dermatologist, rheumatologist or other specialist in the treatment of psoriasis; **AND**
- Symptoms persistent for ≥ 6 months with at least 1 of the following:
 - Involvement of at least 3% of body surface area (BSA); **OR**
 - Psoriasis Area and Severity Index (PASI) score of 10 or greater; **OR**
 - Incapacitation due to plaque location (i.e., head and neck, palms, soles, or genitalia); **AND**
- Trial and failure (at least 3 months) of ≥ 1 conventional therapy:
 - Disease-modifying anti-rheumatic drug (DMARD), such as methotrexate
 - Immunosuppressant (e.g., cyclosporine)
 - Oral retinoid (e.g., acitretin); **AND**
- NOT used in combination with any other biologic agent; **AND**
- For non-preferred agents: 3-month trial and failure of, contraindication or intolerance to, ≥ 1 preferred cytokine or CAM antagonist indicated for the treatment of this condition; **AND**
- Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.

Renewal Criteria:

- Documentation (e.g., progress note) of response to therapy compared to baseline, such as redness, thickness, scaliness, amount of surface area involvement, and/or PASI score.

10. PSORIATIC ARTHRITIS (PSA) CLINICAL CRITERIA

- Diagnosis of psoriatic arthritis (PsA); **AND**
- Prescribed by, or in consultation with, a dermatologist, rheumatologist or other specialist in the treatment of PsA; **AND**

- NOT used in combination with any other biologic agent; **AND**
- For non-preferred agents: 3-month trial and failure of, contraindication or intolerance to, ≥ 1 preferred cytokine or CAM antagonist indicated for the treatment of this condition; **AND**
- Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.

Renewal Criteria:

- Documentation (e.g., progress note) of response to therapy.

11. RHEUMATOID ARTHRITIS (RA) CLINICAL CRITERIA

- Diagnosis of rheumatoid arthritis (RA) based on the American College of Rheumatology (ACR) criteria; **AND**
- Prescribed by, or in consultation with, a rheumatologist or other specialist in the treatment of RA; **AND**
- Documentation (e.g., progress note) of baseline RA disease activity measure using the clinical disease activity index, Disease Activity Score in 28 Joints with Erythrocyte Sedimentation Rate or C-Reactive Protein Level, Simplified Disease Activity Index, Routine Assessment of Patient Index Data 3, or Patient Activity Scale-II; **AND**
- Trial and failure (at least 3 months), contraindication or intolerance to, ≥ 1 disease-modifying anti-rheumatic drug (DMARD), such as methotrexate; **AND**
- NOT used in combination with any other biologic agent; **AND**
- For non-preferred agents: 3-month trial and failure of, contraindication or intolerance to, ≥ 1 preferred cytokine or CAM antagonist indicated for the treatment of this condition; **AND**
- Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.

Renewal Criteria:

- Documentation (e.g., progress note) of improved RA disease activity measure from baseline while on therapy.

12. UVEITIS CLINICAL CRITERIA

- Diagnosis of non-infectious intermediate, posterior, or panuveitis; **AND**
- Prescribed by, or in consultation with, a rheumatologist, ophthalmologist or other specialist in the treatment of uveitis; **AND**
- Failure of a ≥ 2 week trial of a systemic corticosteroid (e.g., prednisone) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced; **AND**
- Failure of a trial of a non-biologic immunosuppressive therapy (e.g., azathioprine, methotrexate, mycophenolate mofetil, cyclosporine, tacrolimus, cyclophosphamide, chlorambucil) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced; **AND**
- NOT used in combination with any other biologic agent; **AND**
- For non-preferred agents: 3-month trial and failure of, contraindication or intolerance to, ≥ 1 preferred cytokine or CAM antagonist indicated for the treatment of this condition; **AND**
- Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.

Renewal Criteria:

- Documentation (e.g., progress note) of response to therapy.

13. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Actemra syringe ^{CC, QL} Actemra Actpen ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Juvenile Idiopathic Arthritis (JIA) Clinical Criteria • Rheumatoid Arthritis (RA) Clinical Criteria <p>Giant Cell Arteritis (GCA) Criteria:</p> <ul style="list-style-type: none"> • Prescribed by a rheumatologist, vascular medicine, or other specialist in the diagnosis and treatment of GCA; AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Systemic Sclerosis-Associated Interstitial Lung Disease (SSc-ILD) Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of Systemic Sclerosis-Associated Interstitial Lung Disease (SSc-ILD) • Prescribed by a pulmonologist, or other specialist in the diagnosis and treatment of SSc-ILD; AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.
Adalimumab biosimilars ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Ankylosing Spondylitis (AS) and nonradiographic axial spondylarthritis (nr-axSpA) Clinical Criteria • Crohn's Disease (CD) or Ulcerative Colitis (UC) Clinical Criteria • Hidradenitis Suppurativa (HS) Clinical Criteria • Juvenile Idiopathic Arthritis (JIA) Clinical Criteria • Plaque Psoriasis Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria • Rheumatoid Arthritis (RA) Clinical Criteria • Uveitis Clinical Criteria
Bimzelx ^{AE, CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Ankylosing Spondylitis (AS) or nonradiographic axial spondyloarthritis (nr-axSpA) Clinical Criteria • Plaque Psoriasis Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria • Hidradenitis Suppurativa (HS) clinical criteria <p>Age Limit: ≥ 18 years of age Quantity Limit: 2 mL per 28 days</p>
Cibinqo ^{CC, QL}	<ul style="list-style-type: none"> • Patient has moderate-to-severe atopic dermatitis (AD) defined by ≥ 1 of the following: <ul style="list-style-type: none"> ○ Involvement of ≥ 10% of body surface area (BSA); OR ○ Eczema Area and Severity Index (EASI) score of ≥ 16; OR ○ Investigator's Global Assessment (IGA) score of ≥ 3; OR ○ Scoring Atopic Dermatitis (SCORAD) score of ≥ 25; OR ○ Pruritus Numerical Rating Scale (NRS) score of ≥ 4; OR ○ Incapacitation due to AD lesion location (head and neck, palms, soles, or genitalia); AND • Prescribed by, or in consultation with, a dermatologist, rheumatologist or other specialist in the treatment of atopic dermatitis; AND • Patient is up to date with all vaccinations, in accordance with current vaccination guidelines, prior to initiating therapy; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Patient will NOT receive live vaccines during therapy; AND • The medication will NOT be used in combination with other monoclonal antibody biologics; AND • Patient is NOT on concomitant antiplatelet therapies during the first 3 months of treatment (Note: excludes the use of low-dose aspirin) AND • Patient does NOT have any clinically relevant laboratory abnormalities (e.g., platelet count <150,000/mm³, an absolute lymphocyte count <500/mm³, an absolute neutrophil count <1,000/mm³, or a hemoglobin value <8 g/dL); AND • Patient has had a ≥ 3 month trial and failure, contraindication, or intolerance to ≥ 1 agent in each of the following categories: <ul style="list-style-type: none"> ○ Topical corticosteroid of medium to high potency (e.g., mometasone, fluocinolone) unless inappropriate for the location (e.g., face, groin); AND ○ Topical calcineurin inhibitor (i.e., tacrolimus or pimecrolimus); AND ○ Immunomodulating systemic agent (e.g., cyclosporine, azathioprine, methotrexate, mycophenolate mofetil, dupilumab); AND • Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Patient has disease response as indicated by improvement in signs and symptoms compared to baseline in ≥ 1 of the following: pruritus, the amount of surface area involvement, EASI, IGA, SCORAD, and/or NRS; AND <ul style="list-style-type: none"> ○ Patient has achieved clear or almost clear skin defined as achievement of an IGA 0/1 or EASI-75 at week 16; OR ○ Patient has had an inadequate response to standard doses of therapy after an adequate trial of ≥ 12 weeks OR patient experienced a disease flare and will require higher dosing; AND ○ Patient requires an increase in dose, in accordance with prescribing information recommended dosages (e.g., up to 200 mg daily); AND • Patient has NOT experienced a myocardial infarction or stroke <p>Quantity Limit: 1 per day</p>
Cimzia ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Ankylosing Spondylitis (AS) or nonradiographic axial spondyloarthritis (nr-axSpA) Clinical Criteria • Crohn's Disease (CD) Clinical Criteria • Juvenile Idiopathic Arthritis (JIA) Clinical Criteria • Plaque Psoriasis Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria • Rheumatoid Arthritis (RA) Clinical Criteria
Enspryng ^{AE, CC, QL}	<ul style="list-style-type: none"> • Diagnosis of anti-aquaporin-4 (AQP4) antibody positive neuromyelitis optica spectrum disorder (NMOSD) • Prescribed by a specialist (e.g., immunologist, neurologist, ophthalmologist, etc.) experienced in the diagnosis and treatment of neuromyelitis optica spectrum disorder (NMOSD); AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. • Diagnosis of NMOSD confirmed by the following: <ul style="list-style-type: none"> ○ Seropositive for aquaporin-4 (AQP4) IgG antibodies; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ Presence of ≥ 1 core clinical characteristic (e.g., optic neuritis, acute myelitis, area postrema syndrome, acute brainstem syndrome, symptomatic narcolepsy or acute diencephalic clinical syndrome with NMOSD-typical diencephalic MRI lesions, symptomatic cerebral syndrome with NMOSD-typical brain lesions); AND ○ Alternative diagnoses have been excluded (e.g., multiple sclerosis, sarcoidosis, cancer, chronic infection); AND ● Patient meets ALL of the following conditions: <ul style="list-style-type: none"> ○ History of ≥ 1 relapse(s) that required rescue therapy within the prior year or ≥ 2 relapses that required rescue therapy within the prior 2 years; AND ○ Expanded Disability Status Score (EDSS) of ≤ 6.5 (e.g., requires 2 walking aids [pair of canes, crutches, etc.] to walk about 20 m without resting); AND ○ At risk of having a disabling relapse of NMOSD for which oral agents (e.g., corticosteroids and immunosuppressants such as azathioprine and mycophenolate) alone are inadequate and biologic therapy is necessary; AND ○ Screening for and absence of Hepatitis B, tuberculosis (TB), and other active infections prior to therapy initiation; AND ● NOT previously treated with prolonged immunosuppressive therapy with alemtuzumab, cladribine, cyclophosphamide or mitoxantrone OR immunosuppressant procedures (e.g., bone marrow transplant, total lymphoid irradiation); AND ● NOT to be used in combination with any of the following: <ul style="list-style-type: none"> ○ Multiple sclerosis agents (e.g., interferon, dimethyl fumarate, fingolimod, glatiramer, etc.) within 6 months of therapy initiation; AND ○ Other biologics used for the treatment of NMOSD (e.g., eculizumab, inebilizumab, rituximab). <p>Renewal Criteria:</p> <ul style="list-style-type: none"> ● Disease response as indicated by stabilization/improvement in any of the following: neurologic symptoms as evidenced by a decrease in acute relapses, stability, or improvement in EDSS, reduced hospitalizations, reduction/discontinuation in plasma exchange treatments, and/or reduction/discontinuation of corticosteroids without relapse.
Entyvio Pen ^{CC, QL} Entyvio vial ^{CC}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> ● Crohn's Disease (CD) or Ulcerative Colitis (UC) Clinical Criteria
Ilaris ^{CC, QL}	<p>Quantity Limit (pens only): 2 pens per 28 days</p> <p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> ● Juvenile Idiopathic Arthritis (JIA) Clinical Criteria <p>Diagnosis of Cryopyrin-Associated Periodic Syndromes (CAPS)</p> <ul style="list-style-type: none"> ● Patient must have a diagnosis of Cryopyrin-Associated Periodic Syndromes (CAPS), including: <ul style="list-style-type: none"> ○ Familial Cold Auto-inflammatory Syndrome (FCAS); OR ○ Muckle-Wells Syndrome (MWS); AND ● Patient must meet the minimum age recommended by the package insert for this FDA-approved indication; AND ● Must be prescribed by or in consultation with a rheumatologist or other specialist in the diagnosis and treatment of CAPS; AND ● Patient is not on concurrent therapy with other IL-1 blocking agents (e.g., anakinra, rilonacept); AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Patient is not on concurrent treatment with a TNF inhibitor, biologic response modifier, or other non-biologic immunomodulating agent (i.e., apremilast, tofacitinib, baricitinib); AND • Patient has documented baseline serum levels of inflammatory proteins (C-Reactive Protein [CRP] and Serum Amyloid A [SAA]); AND • Patient has documented laboratory evidence of a genetic mutation in the Cold Induced Autoinflammatory Syndrome 1, also known as NLRP3; AND • Other causes for recurrent fever have been excluded (e.g. recurrent bacterial/viral infection, other autoinflammatory disease, cancer, cyclic neutropenia, interferonopathies); AND • Patient has ≥2 of any of the CAPS-typical symptoms: <ul style="list-style-type: none"> ○ Urticaria-like rash ○ Cold-triggered episodes ○ Sensorineural hearing loss ○ Musculoskeletal symptoms ○ Chronic aseptic meningitis ○ Skeletal abnormalities <p>Diagnosis of Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS)</p> <ul style="list-style-type: none"> • Patient has a diagnosis of Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS); AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication; AND • Prescribed by or in consultation with a rheumatologist or other specialist in the diagnosis and treatment of TRAPS; AND • Patient is not on concurrent therapy with other IL-1 blocking agents (e.g., anakinra, rilonacept); AND • Patient is not on concurrent treatment with a TNF inhibitor, biologic response modifier, or other non-biologic immunomodulating agent (i.e., apremilast, tofacitinib, baricitinib); AND • Patient has the presence of the TNFRSF1A mutation; AND • Patient has chronic or recurrent disease (defined as > 6 flares per year); AND • Other causes for recurrent fever have been excluded (e.g. recurrent bacterial/viral infection, other autoinflammatory diseases, cancer, cyclic neutropenia, interferonopathies). <p>Diagnosis of Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD)</p> <ul style="list-style-type: none"> • Patient has a diagnosis of Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD); AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication; AND • Prescribed by or in consultation with a rheumatologist or other specialist in the diagnosis and treatment of MKD; AND • Patient is not on concurrent therapy with other IL-1 blocking agents (e.g., anakinra, rilonacept); AND • Patient is not on concurrent treatment with a TNF inhibitor, biologic response modifier, or other non-biologic immunomodulating agent (i.e., apremilast, tofacitinib, baricitinib); AND • Patient has a confirmed diagnosis based on elevated serum IgD levels and mevalonate kinase (MVK) gene mutation testing, if IgD levels are normal; AND • Patient has tried and failed nonsteroidal anti-inflammatory drugs (NSAIDs) and glucocorticoids; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Patient has a documented history of at least three (3) febrile episodes within a 6 month period or is steroid dependent; AND • Other causes have been excluded for recurrent fever (e.g. bacterial/viral/fungal infection, sarcoidosis, cancer,) and/or recurrent abdominal pain and/or elevated IgD. Documentation maybe requested. <p>Diagnosis of Familial Mediterranean Fever (FMF)</p> <ul style="list-style-type: none"> • Patient has a diagnosis of Familial Mediterranean Fever (FMF); AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication; AND • Prescribed by or in consultation with a rheumatologist or other specialist in the diagnosis and treatment of FMF; AND • Patient is not on concurrent therapy with other IL-1 blocking agents (e.g., anakinra, rilonacept); AND • Patient is not on concurrent treatment with a TNF inhibitor, biologic response modifier, or other non-biologic immunomodulating agent (i.e., apremilast, tofacitinib, baricitinib); AND • Patient has a confirmed diagnosis based on of the following: <ul style="list-style-type: none"> ○ Patient continues to have one or more attacks monthly after a six-month compliant trial of colchicine at maximum tolerated doses.; OR ○ Patient has AA amyloidosis while on maximum tolerated doses of colchicine; OR ○ Patient has an intolerance or contraindication to colchicine therapy.; AND • Other causes for recurrent fever have been excluded (e.g. bacterial/viral infection, other autoinflammatory diseases, cancer, other causes of abdominal pain). Documentation maybe requested. <p>Diagnosis of Still's Disease (Adult-Onset Still's Disease [AOSD])</p> <ul style="list-style-type: none"> • Patient has a diagnosis of Still's Disease; AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication; AND • Prescribed by or in consultation with a rheumatologist or other specialist with expertise in treating the diagnosis for AOSD; AND • Patient is not on concurrent therapy with other IL-1 blocking agents (e.g., anakinra, rilonacept); AND • Patient is not on concurrent treatment with a TNF inhibitor, biologic response modifier, or other non-biologic immunomodulating agent (i.e., apremilast, tofacitinib, baricitinib); AND • Patient has active disease; AND • Physician has assessed baseline disease severity utilizing an objective measure/tool; AND • Patient has had at least a 1-month trial and failure (unless contraindicated or intolerant) of previous therapy with either oral non-steroidal anti-inflammatory drugs (NSAIDs) or a systemic glucocorticoid (e.g., prednisone, methylprednisolone); AND • Other causes for recurrent fever have been excluded (e.g., bacterial/viral infection, other autoinflammatory diseases, cancer, sarcoidosis). Documentation maybe requested. <p>Diagnosis of Gout Flare Approval Duration: 3 months</p> <ul style="list-style-type: none"> • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication; AND • Prescribed by or in consultation with a rheumatologist; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Patient is not on concurrent therapy with other IL-1 blocking agents (e.g., anakinra, rilonacept); AND • Patient has had three or more flares in the past 12 months; AND • Patient is currently experiencing a gout flare; AND • Patient has tried and failed all the following unless contraindicated or intolerant: <ul style="list-style-type: none"> ○ Non-steroidal anti-inflammatory drugs (NSAIDs); AND ○ Systemic corticosteroids; AND ○ Colchicine; AND • Patient must be taking a medication for prophylactic treatment of gout. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Absence of unacceptable toxicity from the drug; AND • Canakinumab will not be used concurrently with a TNF inhibitor, biologic response modifier, or other non-biologic immunomodulating agent (i.e., apremilast, tofacitinib, baricitinib); AND • Cryopyrin-Associated Periodic Syndromes: Documentation submitted showing disease improvement or stabilization as symptom assessment and improvement in serum levels of inflammatory proteins (e.g., C-Reactive Protein [CRP] and/or Serum Amyloid A [SAA]) as compared to baseline; OR • Tumor Necrosis Factor Receptor Associated Periodic Syndrome; Hyperimmunoglobulin D Syndrome/Mevalonate Kinase Deficiency; Familial Mediterranean Fever: Documentation submitted showing disease improvement or stabilization as indicated by symptom assessment and changes in any clinically relevant lab values as compared to baseline assessments.; OR • Adult-Onset Still's Disease/Systemic Juvenile Idiopathic Arthritis: Disease response as indicated by improvement in signs and symptoms compared to baseline such as the number of tender and swollen joint counts and/or an improvement on a disease activity scoring tool (e.g., an improvement on a composite scoring index such as Juvenile Arthritis Disease Activity Score [JADAS] or the American College of Rheumatology [ACR] Pediatric [ACR-Pedi 30] of at least 30% improvement from baseline in three of six variables). • Gout Flare: must meet initial criteria for approval. <p>Quantity Limit: 300 mg every 4 weeks</p>
Ilumya ^{AE, CC, QL}	<p>The following diagnosis-based clinical criteria for an FDA-approved has been met:</p> <ul style="list-style-type: none"> • Plaque Psoriasis Clinical Criteria must be met.
Imuldosa ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compend</p> <ul style="list-style-type: none"> • Crohn's Disease (CD) Clinical Criteria • Plaque Psoriasis Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria • Ulcerative Colitis (UC) Clinical Criteria
Infliximab vial (and biosimilars) ^{CC}	<p>Patient has a confirmed diagnosis of one of the following:</p> <ul style="list-style-type: none"> • Crohn's Disease, Fistulizing <p>One of the following diagnosis-based clinical criteria for an FDA-approved- indication has been met:</p> <ul style="list-style-type: none"> • Ankylosing Spondylitis (AS) Clinical Criteria • Crohn's Disease (CD) or Ulcerative Colitis (UC) Clinical Criteria • Juvenile Idiopathic Arthritis (JIA) Clinical Criteria • Plaque Psoriasis Clinical Criteria

Agent(s) Subject to Criteria	Criteria for Approval
Kevzara ^{AE, CC, QL}	<ul style="list-style-type: none"> Psoriatic Arthritis (PsA) Clinical Criteria Rheumatoid Arthritis (RA) Clinical Criteria <p>The following diagnosis-based clinical criteria for an FDA-approved:</p> <ul style="list-style-type: none"> Rheumatoid Arthritis (RA) Clinical Criteria Juvenile Idiopathic Arthritis (JIA) Clinical Criteria <p>Polymyalgia Rheumatica (PMR) Criteria</p> <ul style="list-style-type: none"> Diagnosis of polymyalgia rheumatica (PMR); AND Prescribed by a rheumatologist, or other specialist in the diagnosis and treatment of PMR; AND Patient has steroid-resistant active disease; AND Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.
Kineret ^{CC, QL}	<p>The following diagnosis-based clinical criteria for an FDA-approved:</p> <ul style="list-style-type: none"> Rheumatoid Arthritis (RA) Clinical Criteria must be met; OR <p>DIRA Criteria</p> <ul style="list-style-type: none"> Diagnosis of deficiency of Interleukin-1 Receptor Antagonist (DIRA); AND Prescribed by, or in consultation with, a pediatric rheumatologist, geneticist or other specialist in the diagnosis and treatment of DIRA; AND Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>NOMID Criteria</p> <ul style="list-style-type: none"> Diagnosis of neonatal-onset multisystem inflammatory disease (NOMID); AND Prescribed by, or in consultation with, a rheumatologist or other specialist in the diagnosis and treatment of NOMID; AND Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.
Olumiant ^{AE, CC, QL}	<p>The following diagnosis-based clinical criteria for an FDA-approved:</p> <ul style="list-style-type: none"> Rheumatoid Arthritis (RA) Clinical Criteria must be met.
OmvoH ^{AE, CC, QL}	<p>The following diagnosis-based clinical criteria for an FDA-approved-indication has been met:</p> <ul style="list-style-type: none"> Crohn's Disease (CD) or Ulcerative Colitis (UC) Clinical Criteria <p>Age Limit: ≥ 18 years of age Quantity Limit:</p> <ul style="list-style-type: none"> 1 vial per 28 days 100 mg: 2 per 28 days 200 mg, 300 mg: 1 per 28 days
Orencia ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved:</p> <ul style="list-style-type: none"> Juvenile Idiopathic Arthritis (JIA) Clinical Criteria Psoriatic Arthritis (PsA) Clinical Criteria Rheumatoid Arthritis (RA) Clinical Criteria <p>AND</p> <ul style="list-style-type: none"> Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.
Otufi ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> Crohn's Disease (CD) Clinical Criteria Plaque Psoriasis Clinical Criteria Psoriatic Arthritis (PsA) Clinical Criteria

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Ulcerative Colitis (UC) Clinical Criteria
Pyzchiva ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> Crohn's Disease (CD) Clinical Criteria Plaque Psoriasis Clinical Criteria Psoriatic Arthritis (PsA) Clinical Criteria Ulcerative Colitis (UC) Clinical Criteria
Remicade vial ^{CC}	<p>Patient has a confirmed diagnosis of one of the following:</p> <ul style="list-style-type: none"> Crohn's Disease, Fistulizing <p>One of the following diagnosis-based clinical criteria for an FDA-approved- indication has been met:</p> <ul style="list-style-type: none"> Ankylosing Spondylitis (AS) Clinical Criteria Crohn's Disease (CD) or Ulcerative Colitis (UC) Clinical Criteria Plaque Psoriasis Clinical Criteria Psoriatic Arthritis (PsA) Clinical Criteria Rheumatoid Arthritis (RA) Clinical Criteria
Selarsdi ^{CC, QL} ustekinumab-aekn ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> Crohn's Disease (CD) Clinical Criteria Plaque Psoriasis Clinical Criteria Psoriatic Arthritis (PsA) Clinical Criteria Ulcerative Colitis (UC) Clinical Criteria
Sotyktu ^{AE, CC, QL} Simponi ^{CC, QL}	<ul style="list-style-type: none"> Plaque Psoriasis Clinical Criteria must be met. <p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> Ankylosing Spondylitis (AS) Clinical Criteria Psoriatic Arthritis (PsA) Clinical Criteria Rheumatoid Arthritis (RA) Clinical Criteria Ulcerative Colitis (UC) Clinical Criteria
Simponi Aria ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> Ankylosing Spondylitis (AS) Clinical Criteria Juvenile Idiopathic Arthritis (JIA) Clinical Criteria Psoriatic Arthritis (PsA) Clinical Criteria Rheumatoid Arthritis (RA) Clinical Criteria <p>AND</p> <ul style="list-style-type: none"> Patient must meet the minimum age recommended by the package insert for this FDA-approved indication.
Skyrizi ^{AE, CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> Crohn's Disease (CD) Clinical Criteria Plaque Psoriasis Clinical Criteria Psoriatic Arthritis (PsA) Clinical Criteria Ulcerative Colitis (UC) Clinical Criteria
Stelara ^{CC, QL} ustekinumab ^{CC, QL} ustekinumab-hmny ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> Crohn's Disease (CD) Clinical Criteria Plaque Psoriasis Clinical Criteria Psoriatic Arthritis (PsA) Clinical Criteria Ulcerative Colitis (UC) Clinical Criteria

Agent(s) Subject to Criteria	Criteria for Approval
Steqeyma ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Crohn's Disease (CD) Clinical Criteria • Plaque Psoriasis Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria • Ulcerative Colitis (UC) Clinical Criteria
Taltz ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Ankylosing Spondylitis (AS) or nonradiographic axial spondyloarthritis (nr-axSpA) Clinical Criteria • Plaque Psoriasis Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria
Tremfya ^{AE, CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Crohn's Disease (CD) Clinical Criteria • Plaque Psoriasis Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria • Ulcerative Colitis (UC) Clinical Criteria
ustekinumab-ttwe ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Crohn's Disease (CD) Clinical Criteria • Plaque Psoriasis Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria • Ulcerative Colitis (UC) Clinical Criteria
Velsipity ^{AE, CC, QL}	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of moderate to severe ulcerative colitis (UC); AND • Prescribed by, or in consultation with, a gastroenterologist or other specialist in the treatment of UC; AND • Patient has had a trial and failure of ≥ 1 of the following conventional therapies: <ul style="list-style-type: none"> ○ Oral/rectal 5-aminosalicylic acid agents (e.g., balsalazide, Lialda, mesalamine, sulfasalazine) ○ Oral/rectal steroids (e.g., budesonide, hydrocortisone, prednisone) ○ Immunosuppressant (e.g., azathioprine, mercaptopurine); OR • Patient is deemed high-risk for intestinal complications or post-operative recurrence; AND • NOT used in combination with any other biologic agent; AND • Patient has had a 3-month trial and failure of, or contraindication or intolerance to, ≥ 1 preferred cytokine or CAM antagonist indicated for the treatment of UC; AND • Patient meets the minimum age recommended by the package insert for use in UC. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Documentation (e.g., progress notes) of response to therapy compared to baseline. <p>Age Limit: ≥ 18 years of age Quantity Limit: 1 tablet per day</p>
Xeljanz XR ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> • Ankylosing Spondylitis (AS) or nonradiographic axial spondyloarthritis (nr-axSpA) Clinical Criteria • Juvenile Idiopathic Arthritis (JIA) Clinical Criteria • Psoriatic Arthritis (PsA) Clinical Criteria • Rheumatoid Arthritis (RA) Clinical Criteria

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Ulcerative Colitis (UC) Clinical Criteria
Yesintek ^{CC, QL}	<p>One of the following diagnosis-based clinical criteria for an FDA-approved or compendia-supported indication have been met:</p> <ul style="list-style-type: none"> Crohn's Disease (CD) Clinical Criteria Plaque Psoriasis Clinical Criteria Psoriatic Arthritis (PsA) Clinical Criteria Ulcerative Colitis (UC) Clinical Criteria
Zymfentra ^{CC, QL}	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> Diagnosis of moderate to severe Crohn's disease (CD) or ulcerative colitis (UC); AND Patient has undergone induction therapy with intravenous infliximab; AND Prescribed by, or in consultation with, a gastroenterologist or other specialist in the treatment of CD or UC; AND Patient has had a trial and failure of ≥ 1 of the following conventional therapies: <ul style="list-style-type: none"> Oral/rectal 5-aminosalicylic acid agents (e.g., Apriso, balsalazide, Lialda, mesalamine, sulfasalazine) Oral/rectal steroids (e.g., budesonide, hydrocortisone, prednisone) Immunosuppressant (e.g., azathioprine, mercaptopurine); OR Patient is deemed high-risk for intestinal complications or post-operative recurrence; AND NOT used in combination with any other biologic agent; AND Patient has had a 3-month trial and failure of, or contraindication or intolerance to, ≥ 1 preferred cytokine or CAM antagonist indicated for the treatment of CD or UC; AND Patient meets the minimum age recommended by the package insert for use in CD or UC. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Documentation (e.g., progress notes) of response to therapy compared to baseline. <p>Quantity Limit: 2 pens or syringes per month</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Cosentyx ^{CC, QL}	Abrilada ^{CC, QL}
Enbrel ^{CC, QL}	Actemra ^{CC, QL}
Hadlima ^{CC, QL}	adalimumab-aacf ^{CC, QL}
Humira ^{CC, QL}	adalimumab-aaty ^{CC, QL}
Otezla ^{CC, QL}	adalimumab-adaz ^{CC, QL}
Otezla XR ^{CC, QL}	adalimumab-adbm ^{CC, QL}
Rinvoq ^{AE, CC, QL}	adalimumab-fjkg ^{CC, QL}
Rinvoq LQ ^{AE, CC, QL}	adalimumab-ryvk ^{CC, QL}
Tyenne ^{CC, QL}	Amjevita ^{CC, QL}
Xeljanz ^{CC, QL}	Avsola vial ^{CC}
	Bimzelx ^{AE, CC, QL}
	Cibinqo ^{CC, QL}
	Cimzia ^{CC, QL}
	Cyltezo ^{CC, QL}
	Enspryng ^{AE, CC, QL}
	Entyvio pen ^{CC, QL}

Preferred Agents	Non-Preferred Agents
	Entyvio vial ^{CC}
	Hulio ^{CC, QL}
	Hyrimoz ^{CC, QL}
	Idacio ^{CC, QL}
	Ilaris ^{CC, QL}
	Ilumya ^{AE, CC, QL}
	Imuldosa ^{CC, QL}
	Inflectra vial ^{CC}
	Infliximab vial ^{CC}
	Kevzara ^{AE, CC, QL}
	Kineret ^{CC, QL}
	Olumiant ^{AE, CC, QL}
	OmvoH ^{AE, CC, QL}
	Orencia ^{CC, QL}
	Otulf ^{CC, QL}
	Pyzchiva ^{CC, QL}
	Remicade vial ^{CC}
	Renflexis vial ^{CC}
	Selarsdi ^{CC, QL}
	Simponi ^{CC, QL}
	Simponi Aria ^{AE, CC, QL}
	Simlandi ^{CC, QL}
	Skyrizi ^{AE, CC, QL}
	Sotyktu ^{AE, CC, QL}
	Stelara ^{CC, QL}
	Steqeyma ^{CC, QL}
	Taltz ^{CC, QL}
	Tremfya ^{AE, CC, QL}
	ustekinumab ^{CC, QL}
	ustekinumab-aekn ^{CC, QL}
	ustekinumab-ttwe ^{CC, QL}
	Velsipity ^{AE, CC, QL}
	Xeljanz XR ^{CC, QL}
	Yesintek ^{CC, QL}
	Yuflyma ^{CC, QL}
	Yusimry ^{CC, QL}
	Zymfentra ^{CC, QL}

IMMUNOLOGIC AND GENETIC: IMMUNOMODULATORS, ASTHMA

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Fasenra ^{CC, QL, AE}	<p>Approval Duration: 6 months Initial, 1 year Renewal</p> <p>Asthma Initial Criteria</p> <ul style="list-style-type: none"> • Diagnosis of eosinophilic asthma; AND • Patient's asthma is classified as severe, as defined by one of the following: <ul style="list-style-type: none"> ○ Uncontrolled symptoms while on dose optimized high dose ICS-LABA therapy; OR ○ Loss of symptom control when high dose ICS-LABA therapy is decreased; OR ○ Use of oral steroids ≥ 2 times in the past year; AND • Fasenra will be used as adjunct therapy with inhaled maintenance asthma therapy; AND • Prescribed by, or in consultation with, an allergist, immunologist, pulmonologist, or other applicable specialist in the diagnosis and treatment of eosinophilic asthma; AND • NOT used in combination with any other biologic agent; AND • Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient has experienced disease improvement and/or stabilization based on an objective measure such as (but not limited to): <ul style="list-style-type: none"> ○ Improved or maintained FEV₁ ○ Reduced number of asthma exacerbations ○ Reduced number of missed days of school/work; AND • Fasenra will be used as adjunct therapy with inhaled maintenance asthma therapy. <p>Eosinophilic Granulomatosis with Polyangiitis (Churg-Strauss) Initial Criteria</p> <ul style="list-style-type: none"> • Diagnosis of eosinophilic granulomatosis with polyangiitis (EGPA) also known as Churg-Strauss; AND • Patient has had a ≥ 90 days of treatment with one of the following agents: <ul style="list-style-type: none"> ○ Systemic glucocorticoids ○ Azathioprine ○ Methotrexate ○ Cyclophosphamide ○ Mycophenolate • Prescribed by, or in consultation with, an allergist, immunologist, or other applicable specialist in the diagnosis and treatment of eosinophilic granulomatosis with polyangiitis; AND • NOT used in combination with any other biologic agent; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient has experienced disease improvement and/or stabilization based on an objective measure
Nucala ^{CC, QL, AE}	<p>Approval Duration: 6 months Initial, 1 year Renewal</p> <p>Asthma</p> <p>Initial Criteria</p> <ul style="list-style-type: none"> • Diagnosis of eosinophilic asthma; AND • Patient's asthma is classified as severe, as defined by one of the following: <ul style="list-style-type: none"> ○ Uncontrolled symptoms while on dose optimized ICS-LABA therapy; OR ○ Loss of symptom control when high dose ICS-LABA therapy is decreased; OR ○ Use of oral steroids ≥ 2 times in the past year; AND • Nucala will be used as adjunct therapy with inhaled maintenance asthma therapy; AND • Prescribed by, or in consultation with, an allergist, immunologist, pulmonologist, or other applicable specialist in the diagnosis and treatment of eosinophilic asthma; AND • NOT used in combination with any other biologic agent; AND • Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient has experienced disease improvement and/or stabilization based on an objective measure such as (but not limited to): <ul style="list-style-type: none"> ○ Improved or maintained FEV₁ ○ Reduced number of asthma exacerbations ○ Reduced number of missed days of school/work; AND • Nucala will be used as adjunct therapy with inhaled asthma therapy. <p>Chronic Obstructive Pulmonary Disease (COPD)</p> <p>Initial Criteria</p> <ul style="list-style-type: none"> • Diagnosis of chronic obstructive pulmonary disease (COPD) with eosinophilic phenotype; AND • At least 2 moderate exacerbations or 1 severe exacerbation in the past 12 months despite receiving triple therapy consistent of all of the following: <ul style="list-style-type: none"> ○ 1 long-acting muscarinic antagonist (LAMA) (e.g. Spiriva Handihaler); AND ○ 1 long-acting beta agonist (LABA) (e.g. Serevent); AND ○ 1 inhaled corticosteroid (ICS) (e.g. budesonide, fluticasone propionate); OR • Clinical documentation indicated that the patient has a contraindication or intolerance to drugs in one of the following classes: <ul style="list-style-type: none"> ○ 1 long-acting muscarinic antagonist (LAMA) (e.g. Spiriva Handihaler); AND ○ 1 long-acting beta agonist (LABA) (e.g. Serevent); AND ○ 1 inhaled corticosteroid (ICS) (e.g. budesonide, fluticasone propionate); AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Medication will be used as an add-on maintenance therapy; AND Prescribed by, or in consultation with, a pulmonologist or other specialist in the treatment of COPD; AND Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> Patient has experienced disease improvement and/or stabilization based on an objective measure such as (but not limited to): <ul style="list-style-type: none"> Improvement from baseline in FEV1; OR Reduced use of systemic corticosteroids; OR Reduced use of antibiotics; OR Reduced number of hospitalizations, ER visits, or unscheduled visits to healthcare provider due to condition. <p><i>Chronic Rhinosinusitis with Nasal Polyps</i></p> <p>Initial Criteria</p> <ul style="list-style-type: none"> Diagnosis of chronic rhinosinusitis with nasal polyps; AND Trial and failure ≥ 1 intranasal corticosteroid, unless intolerant or otherwise ineligible; AND Nucala will be used as adjunct therapy with an intranasal corticosteroid for maintenance, unless intolerant or otherwise ineligible asthma therapy; AND Prescribed by, or in consultation with, an allergist; immunologist; ear, nose, and throat (ENT), or other applicable specialist in the diagnosis and treatment of chronic rhinosinusitis with nasal polyps; AND NOT used in combination with any other biologic agent; AND Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> Patient has experienced disease improvement and/or stabilization based on an objective measure. <p><i>Eosinophilic Granulomatosis with Polyangiitis (Churg-Strauss)</i></p> <p>Initial Criteria</p> <ul style="list-style-type: none"> Diagnosis of eosinophilic granulomatosis with polyangiitis (EGPA) also known as Churg-Strauss; AND Patient has had a ≥ 90 days of treatment with one of the following agents: <ul style="list-style-type: none"> Systemic glucocorticoids Azathioprine Methotrexate Cyclophosphamide Mycophenolate Prescribed by, or in consultation with, an allergist, immunologist, or other applicable specialist in the diagnosis and treatment of eosinophilic granulomatosis with polyangiitis; AND NOT used in combination with any other biologic agent; AND Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Renewal Criteria</p>

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Patient has experienced disease improvement and/or stabilization based on an objective measure. <p><i>Hypereosinophilic Syndrome (HES)</i></p> <p>Initial Criteria</p> <ul style="list-style-type: none"> • Diagnosis of hypereosinophilic syndrome (HES) ; AND • Symptoms have been present for > 6 months; AND • Provider attests that other underlying causes have been ruled out; AND • Prescribed by, or in consultation with, an allergist, immunologist, or other applicable specialist in the diagnosis and treatment of hypereosinophilic syndrome; AND • NOT used in combination with any other biologic agent; AND • Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient has experienced disease improvement and/or stabilization based on an objective measure.
Xolair CC, QL, AE	<p>Approval Duration: 6 months Initial, 1 year Renewal</p> <p><i>Asthma</i></p> <p>Initial Criteria</p> <ul style="list-style-type: none"> • Diagnosis of moderate to severe persistent asthma; AND • Patient has had one of the following: <ul style="list-style-type: none"> ○ Positive skin test to a perennial aeroallergen; OR ○ Positive in vitro reactivity to a perennial aeroallergen; AND • Patient has experienced inadequate symptom control with inhaled corticosteroids; AND • Prescribed by, or in consultation with, an allergist, immunologist, pulmonologist, or other specialist in the treatment of asthma; AND • NOT used in combination with any other biologic agent; AND • Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient has experienced disease improvement and/or stabilization based on an objective measure such as (but not limited to): <ul style="list-style-type: none"> ○ Improved or maintained FEV₁ ○ Reduced number of asthma exacerbations ○ Reduced number of missed days of school/work. <p><i>Chronic Rhinosinusitis with Nasal Polyps</i></p> <p>Initial Criteria</p> <ul style="list-style-type: none"> • Diagnosis of chronic rhinosinusitis with nasal polyps; AND • Inadequate symptom control with use of ≥ 1 intranasal corticosteroid, unless intolerant or otherwise ineligible; AND • Prescribed by, or in consultation with, an allergist; immunologist; ear, nose, and throat (ENT), or other applicable specialist in the diagnosis and treatment of chronic rhinosinusitis with nasal polyps; AND • NOT used in combination with any other biologic agent; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient has experienced disease improvement and/or stabilization based on an objective measure. <p><i>Chronic Idiopathic Urticaria</i></p> <p>Initial Criteria</p> <ul style="list-style-type: none"> • Diagnosis of chronic idiopathic urticaria; AND • Patient has had a ≥ 14 day trial and failure of therapy with an H1-receptor antagonist; AND • Prescribed by, or in consultation with, an allergist, immunologist, or other applicable specialist in the diagnosis and treatment of chronic idiopathic urticaria; AND • NOT used in combination with any other biologic agent; AND • Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient has experienced disease improvement and/or stabilization based on an objective measure. <p><i>IgE-Mediated Food Allergy</i></p> <p>Initial Criteria</p> <ul style="list-style-type: none"> • Diagnosis of IgE-mediated food allergy; AND • Patient will avoid further contact with food allergen; AND • Prescribed by, or in consultation with, an allergist, immunologist, or other applicable specialist in the diagnosis and treatment of IgE-mediated food allergy; AND • NOT used in combination with any other biologic agent; AND • Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient has experienced disease improvement and/or stabilization based on an objective measure.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires ≥ 90 day trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Tezspire ^{CC, AE, QL}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none"> • Patient must have a diagnosis of severe asthma; AND • Must be used for add-on maintenance treatment in patients regularly receiving BOTH of the following: <ul style="list-style-type: none"> ○ Medium-to-high dose inhaled corticosteroids; AND ○ An additional controller medication (e.g., long-acting beta agonist, leukotriene modifiers); AND • Patient must have had, in the previous year, at least 2 exacerbations requiring oral or injectable corticosteroid treatment (in addition to the regular maintenance therapy defined above) OR one exacerbation resulting in a hospitalization; AND • Baseline measurement of ≥ 1 of the following for assessment of clinical status: <ul style="list-style-type: none"> ○ Use of systemic corticosteroids; OR ○ Use of inhaled corticosteroids; OR ○ Number of hospitalizations, ER visits, or unscheduled visits to healthcare provider due to condition; OR ○ FEV1; AND • Must not be used in combination with anti-IgE, anti-IL4, or anti-IL5 monoclonal antibody agents (e.g., benralizumab, omalizumab, mepolizumab, reslizumab, dupilumab); AND • Patient does not have an active or untreated helminth infection; AND • Will not be administered concurrently with live vaccines; AND • Patient has had a trial and failure, contraindication, or intolerance to at least 1 preferred agent. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Improvement in asthma symptoms, asthma exacerbations, or airway function as evidenced by decrease in ≥ 1 of the following: <ul style="list-style-type: none"> ○ Use of systemic corticosteroids; OR ○ Two-fold or greater decrease in inhaled corticosteroid use for at least 3 days; OR ○ Hospitalizations; OR ○ ER visits; OR ○ Unscheduled visits to healthcare provider; OR ○ Improvement from baseline in FEV1 of $\geq 15\%$; AND <p>Age Limit: ≥ 12 years old Quantity Limit: 1 prefilled syringe per 28 days (0.07 mL per day)</p>

6. THERAPEUTIC DUPLICATION

Approval Duration: Date of Service Only

Patients are limited to one immunomodulator used to treat asthma at a time within the quantity/dosing limits. Therapeutic duplication can be approved for DOS if patients are switching from one agent to another.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Fasenra pen autoinjector ^{CC, AE, QL}	Tezspire pen ^{CC, AE, QL}
Fasenra syringe ^{CC, AE, QL}	Tezspire syringe ^{CC, AE, QL}
Nucala autoinjector ^{CC, AE, QL}	
Nucala syringe ^{CC, AE, QL}	

Preferred Agents		Non-Preferred Agents	
Nucala vial	CC, AE, QL		
Xolair autoinjector	CC, AE, QL		
Xolair syringe	CC, AE, QL		
Xolair vial	CC, AE, QL		

IMMUNOLOGIC AND GENETIC: MUSCULAR DYSTROPHY AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Emflaza tablets ^{AE, CC, QL} Emflaza suspension ^{AE, CC}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none">• Diagnosis of Duchenne muscular dystrophy (DMD); AND• Patient is currently receiving, or planning to receive, physical therapy; AND• Patient has experienced 1 of the following adverse reactions directly attributable to previous therapy with prednisone:<ul style="list-style-type: none">○ Significant behavioral changes negatively impacting function at school, home, day care, etc.; OR○ Significant weight gain (e.g., crossing 2 percentiles and/or reaching 98th percentile for age and sex). <p>Renewal Criteria</p> <ul style="list-style-type: none">• Patient continues to receive physical therapy; AND• Patient has received benefit from therapy, which may include 1 or more of the following supported by documentation (e.g., progress notes):<ul style="list-style-type: none">○ Stability, improvement or slowing of decline in motor function;○ Stability, improvement or slowing of decline in respiratory function;○ Stability, improvement or slowing of decline in sequelae related to diminished strength of stabilizing musculature (e.g., scoliosis, etc.);○ Stability, improvement or slowing of decline in quality of life. <p>Age Limit: ≥ 2 years Quantity Limits: 2 tablets per day</p>

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 90 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Agamree ^{AE, CC, QL}	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none">Diagnosis of Duchenne Muscular Dystrophy (DMD); ANDPatient is currently receiving, or planning to receive, physical therapy; ANDPatient has had a trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications), or intolerance to at least 1 preferred agent; ANDPatient has tried prednisone or prednisolone for at least 6 months; ORPatient has experienced 1 of the following adverse reactions directly attributable to previous therapy with prednisone or prednisolone:<ul style="list-style-type: none">Significant behavioral changes negatively impacting function at school, home, day care, etc.; ORSignificant weight gain (e.g., crossing 2 percentiles and/or reaching 98th percentile for age and sex); AND <p>Renewal Criteria:</p> <ul style="list-style-type: none">Patient continues to receive physical therapy; ANDPatient has received benefit from therapy (i.e. stability, improvement or slowing of decline) in one or more of the following areas of assessment:<ul style="list-style-type: none">Motor function (North Star Ambulatory Assessment (NSAA))CardiologyEndocrinologyOrthopedics (e.g., scoliosis)Pulmonary function. <p>Age Limit: ≥ 2 years of age Quantity Limit: 7.5 mL per day</p>
deflazacort tablet ^{AE, CC, QL} deflazacort suspension ^{AE, CC}	<p>Initial Approval Criteria</p> <ul style="list-style-type: none">Diagnosis of Duchenne muscular dystrophy (DMD); ANDPatient is currently receiving, or planning to receive, physical

Agent(s) Subject to Criteria	Criteria for Approval
	<p>therapy; AND</p> <ul style="list-style-type: none"> • Patient has experienced 1 of the following adverse reactions directly attributable to previous therapy with prednisone: <ul style="list-style-type: none"> ○ Significant behavioral changes negatively impacting function at school, home, day care, etc.; OR ○ Significant weight gain (e.g., crossing 2 percentiles and/or reaching 98th percentile for age and sex); <p>AND</p> <ul style="list-style-type: none"> • Patient has a known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. <p>Renewal Criteria</p> <ul style="list-style-type: none"> • Patient continues to receive physical therapy; AND • Patient has received benefit from therapy, which may include 1 or more of the following supported by documentation (e.g., progress notes): <ul style="list-style-type: none"> ○ Stability, improvement or slowing of decline in motor function; ○ Stability, improvement or slowing of decline in respiratory function; ○ Stability, improvement or slowing of decline in sequelae related to diminished strength of stabilizing musculature (e.g., scoliosis, etc.); ○ Stability, improvement or slowing of decline in quality of life. <p>Age Limit: ≥ 2 years</p> <p>Quantity Limits: 2 tablets per day</p>
	<p>Approval Duration: 6 months initial, 1 year renewal</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of Duchenne muscular dystrophy (DMD) [G71.01]; AND • Platelet count within the last 30 days equals to or is greater than 150 x 10⁹/L; AND • Prescribed by, or in consultation with, a neuromuscular specialist with expertise in the treatment of DMD; AND • Patient is ambulatory (e.g., ability to walk with or without assistive devices, not wheelchair dependent); AND • Patient's baseline ambulatory function has been or will be assessed prior to therapy initiation; AND • Patient has been on a stable systemic corticosteroid therapy for at least 6 months and will continue to be on the systemic corticosteroid therapy unless contraindicated or clinically significant adverse effects are experienced; AND • Prescriber provides a patient weight obtained within the past 3 months; AND • The requested dose meets the FDA-approved dosing recommendation. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Documentation (e.g., progress note) of stabilized or improved ambulatory function from baseline; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Patient will continue systemic corticosteroid therapy unless contraindicated or clinically significant adverse effects are experienced; AND • Prescriber provides a patient weight obtained within the past 3 months; AND • The requested dose meets the FDA-approved dosing recommendation. <p>Age Limit: 6 years of age or older</p> <p>Quantity Limit: 12 mL per day</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Emflaza suspension ^{AE, CC}	Agamree suspension ^{AE, CC, QL}
Emflaza tablet ^{AE, CC, QL}	deflazacort suspension ^{AE, CC}
	deflazacort tablet ^{AE, CC, QL}
	Duvyzat ^{AE, CC, QL}

IMMUNOMODULATORS, ATOPIC DERMATITIS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Adbry Syringe ^{AE, CC, QL} Adbry Autoinjector ^{AE, CC, QL}	<p>Approval Duration: 16 weeks initial, 1 year renewal</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none">• Diagnosis of moderate to severe atopic dermatitis with at least 1 of the following:<ul style="list-style-type: none">○ Involvement of at least 10% of body surface area (BSA); OR○ Eczema Area and Severity Index (EASI) score of 16 or greater; OR○ Investigator's Global Assessment (IGA) score of 3 or more; OR○ Scoring Atopic Dermatitis (SCORAD) score of 25 or more; OR○ Incapacitation due to AD lesion location (i.e., head and neck, palms, soles, or genitalia); AND• Prescribed by, or in consultation with, a dermatologist, allergist/immunologist, or other specialist in the treatment of atopic dermatitis; AND• Patient has had a trial and failure, contraindication, or intolerance to at least 1 agent from ≥ 2 of the following classes:<ul style="list-style-type: none">○ Prescription strength topical corticosteroids (e.g., mometasone, fluocinolone) unless inappropriate for the location (e.g., face, groin); OR○ Topical calcineurin inhibitor (e.g., pimecrolimus or tacrolimus); OR○ Topical phosphodiesterase-4 inhibitor (e.g., crisaborole); OR○ Topical Janus kinase inhibitor (e.g., ruxolitinib); OR○ Immunomodulating systemic agent (e.g., cyclosporine, azathioprine, methotrexate, mycophenolate mofetil, dupilumab) <p>Renewal Criteria:</p> <ul style="list-style-type: none">• Patient must have disease improvement and/or stabilization from baseline <p>Age Limit:</p> <ul style="list-style-type: none">• Adbry autoinjector: ≥ 18 years• Adbry syringe: ≥ 12 years <p>Quantity Limit:</p> <ul style="list-style-type: none">• Adbry autoinjector: 2 injectors per 28 days• Adbry syringe: 4 syringes per 28 days (300 mg every other week)
Dupixent ^{CC, QL}	<p>Atopic Dermatitis</p> <ul style="list-style-type: none">• Diagnosis of moderate-to-severe atopic dermatitis (AD) with ≥ 1 of the following:<ul style="list-style-type: none">○ Involvement of at least 10% of body surface area (BSA); OR○ Scoring Atopic Dermatitis (SCORAD) score of 25 or more; OR○ Investigator's Global Assessment (IGA) with a score ≥ 3; OR○ Eczema Area and Severity Index (EASI) score of ≥ 16; OR○ Incapacitation due to AD lesion location (e.g., head and neck, palms, soles, or genitalia); AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Prescribed by, or in consultation with, a dermatologist, allergist/immunologist, or other specialist in the treatment of atopic dermatitis; AND • Trial and failure, contraindication, or intolerance to ≥ 1 agent in 2 or more of the following categories (total prior agent use of ≥ 90 days): <ul style="list-style-type: none"> ○ Topical corticosteroid of medium to high potency (e.g., mometasone, fluocinolone) unless inappropriate for the location (e.g., face, groin); AND ○ Topical calcineurin inhibitor (i.e., tacrolimus or pimecrolimus); OR ○ Immunosuppressive systemic agent (e.g., cyclosporine, azathioprine, methotrexate, mycophenolate mofetil, etc.) • Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Asthma</p> <ul style="list-style-type: none"> • Diagnosis of moderate-to-severe asthma; AND • Eosinophilic phenotype; OR • Use of oral steroids ≥ 2 times in the past year; AND • Prescribed by, or in consultation with, an allergist, immunologist, pulmonologist or other specialist in the treatment of asthma; AND • Patient is ≥ 6 years of age. <p>Bullous Pemphigoid</p> <ul style="list-style-type: none"> • Diagnosis of bullous pemphigoid (BP); AND • Bullous Pemphigoid Disease Area Index (BPDAI) activity score of ≥ 20 at baseline; AND • Trial and failure, contraindication, or intolerance to ≥ 1 agent in the following categories: Systemic corticosteroids (e.g. prednisone); OR <ul style="list-style-type: none"> ○ Immunosuppressive agents (e.g., azathioprine, methotrexate, mycophenolate mofetil); OR ○ Other BP-directed therapies (e.g., dapsone, doxycycline); ○ Prescribed by, or in consultation with, a dermatologist, allergist/immunologist, or other specialist in the treatment of bullous pemphigoid; AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Chronic Obstructive Pulmonary Disease</p> <ul style="list-style-type: none"> • Diagnosis of chronic obstructive pulmonary disease (COPD) with eosinophilic phenotype; AND • At least 2 moderate exacerbations or 1 severe exacerbation in the past 12 months despite receiving maintenance triple therapy consistent of all of the following: <ul style="list-style-type: none"> ○ 1 long-acting muscarinic antagonist (LAMA) (e.g., Spiriva Handihaler); AND ○ 1 long-acting beta agonist (LABA) (e.g., Serevent); AND ○ 1 inhaled corticosteroid (ICS) (e.g., budesonide, fluticasone propionate); OR • Clinical documentation indicated that the patient has a contraindication or intolerance to drugs in one of the following classes: <ul style="list-style-type: none"> ○ 1 long-acting muscarinic antagonist (LAMA) (e.g., Spiriva Handihaler); AND ○ 1 long-acting beta agonist (LABA) (e.g., Serevent); AND ○ 1 inhaled corticosteroid (ICS) (e.g., budesonide, fluticasone propionate); AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Symptoms of chronic productive cough for at least 3 months in the past 12 months; AND Medication will be used as an add-on maintenance therapy; AND Prescribed by, or in consultation with, a pulmonologist or other specialist in the treatment of COPD; AND Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Chronic Rhinosinusitis with Nasal Polyposis</p> <ul style="list-style-type: none"> Diagnosis of chronic rhinosinusitis with nasal polyposis; AND Patient is ≥ 12 years of age; AND Prescribed by or in consultation with an allergist; immunologist; ear, nose, and throat (ENT), or other applicable specialist; AND Trial and failure (and continued use of) ≥ 1 intranasal corticosteroid, unless intolerant or otherwise ineligible. <p>Chronic Spontaneous Urticaria</p> <ul style="list-style-type: none"> Diagnosis of chronic spontaneous urticaria (CSU); AND Trial and failure of (> 14-day treatment course), contraindication, or intolerance to histamine-1 antihistamine (e.g. diphenhydramine, hydroxyzine); AND Prescribed by, or in consultation with, an allergist, immunologist, or other applicable specialist in the diagnosis and treatment of chronic spontaneous urticaria; AND Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Eosinophilic Esophagitis</p> <ul style="list-style-type: none"> Diagnosis of eosinophilic esophagitis; AND Prescribed by, or in consultation with, an allergist, immunologist, gastroenterologist, or other specialist in the treatment of eosinophilic esophagitis; AND Patient has tried and failed at least 8 weeks of treatment with a topical glucocorticoid; AND Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Prurigo Nodularis</p> <ul style="list-style-type: none"> Diagnosis of prurigo nodularis; AND Patient has severe pruritus based on an objective measure; AND At least 20 nodular lesions; AND Other causes of pruritus have been ruled out; AND Trial and failure, contraindication, or intolerance to one of the following: <ul style="list-style-type: none"> Moderate to super potent topical corticosteroids [e.g., betamethasone dipropionate, (augmented), fluocinonide 0.1%, flurandrenolide, betamethasone dipropionate 0.05%, clobetasol propionate 0.025%, or desoximetasone 0.05%] for a minimum of 2 weeks; OR Narrowband ultraviolet B (NBUVB) phototherapy or psoralen plus ultraviolet A (PUVA) phototherapy; AND Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Patient must have disease improvement and/or stabilization based on an objective measure Diagnosis of atopic dermatitis; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> • Trial and failure of ≥ 1 agent from either of the following classes, unless trial is not appropriate: <ul style="list-style-type: none"> ○ Topical immunomodulator unless < 2 years of age; OR ○ Topical steroid (e.g., triamcinolone, etc.) unless inappropriate for the affected area (e.g., face, groin). <p>Age Limit: ≥ 3 months Quantity Limit: 300 g per 365 days</p>
Opzelura cream ^{AE, CC, QL}	<p>Approval Duration: 1 year</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Patient is not immunocompromised; AND • Diagnosis of mild to moderate atopic dermatitis; AND • Patient is 2 years of age or older; AND • Patient has a history of trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to 1 of the following classes: <ul style="list-style-type: none"> ○ Prescription topical corticosteroids ○ Topical calcineurin inhibitor (e.g., pimecrolimus or tacrolimus) ○ Topical phosphodiesterase-4 inhibitor (e.g., crisaborole); OR <ul style="list-style-type: none"> • Patient has a diagnosis of nonsegmental vitiligo; AND • Patient is 12 years of age or older; AND • Patient has a history of trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to mid- to high-potency topical corticosteroids or topical calcineurin inhibitors. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Patient must continue to meet the above criteria; AND • Patient must have disease improvement and/or stabilization <p>Quantity Limit: 240 grams per 365 days</p>

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Ebglyss ^{AE, CC, QL}	<p>Approval Duration: 4 months initial, 1 year renewal</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of moderate-to-severe atopic dermatitis (AD) with ≥ 1 of the following:

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ Involvement of at least 10% of body surface area (BSA); OR ○ Scoring Atopic Dermatitis (SCORAD) score of 25 or more; OR ○ Investigator's Global Assessment (IGA) with a score ≥ 3; OR ○ Eczema Area and Severity Index (EASI) score of ≥ 16; OR ○ Incapacitation due to AD lesion location (e.g., head and neck, palms, soles, or genitalia); AND • Prescribed by, or in consultation with, a dermatologist, allergist/immunologist, or other specialist in the treatment of atopic dermatitis; AND • Trial and failure, contraindication, or intolerance to ≥ 1 agent in 2 or more of the following categories (total prior agent use of ≥ 90 days): <ul style="list-style-type: none"> ○ Topical corticosteroid of medium to high potency (e.g., mometasone, fluocinolone) unless inappropriate for the location (e.g., face, groin); AND ○ Topical calcineurin inhibitor (i.e., tacrolimus or pimecrolimus); OR ○ Immunosuppressive systemic agent (e.g., cyclosporine, azathioprine, methotrexate, mycophenolate mofetil); AND • Trial and failure, contraindication, or intolerance to at least one preferred injectable agent (Adbry or Dupixent); AND • Patient must meet the minimum age and weight recommended by the package insert for the provided indication. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Patient must continue to meet initial approval criteria; AND • Patient must have disease improvement and/or stabilization based on an objective measure. <p>Age Limit: 12 years of age or older Quantity Limit: 1 pen/syringe (2 mL) per 28 days</p>
Nemluvio AE, CC, QL	<p>Approval Duration: 4 months initial, 1 year renewal</p> <p>Initial Approval Criteria:</p> <p>Atopic Dermatitis:</p> <ul style="list-style-type: none"> • Diagnosis of moderate-to-severe atopic dermatitis (AD) with ≥ 1 of the following: <ul style="list-style-type: none"> ○ Involvement of at least 10% of body surface area (BSA); OR ○ Investigator's Global Assessment (IGA) with a score ≥ 3; OR ○ Eczema Area and Severity Index (EASI) score of ≥ 16; OR ○ Peak Pruritis Numeric Rating Scale (PP-NRS) score ≥ 4; OR ○ Incapacitation due to AD lesion location (e.g., head and neck, palms, soles, or genitalia); AND • Prescribed by, or in consultation with, a dermatologist, allergist/immunologist, or other specialist in the treatment of atopic dermatitis; AND • Trial and failure, contraindication, or intolerance to ≥ 1 agent in 2 or more of the following categories (total prior agent use of ≥ 90 days):

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ Topical corticosteroid of medium to high potency (e.g., mometasone, fluocinolone) unless inappropriate for the location (e.g., face, groin); AND ○ Topical calcineurin inhibitor (i.e., tacrolimus or pimecrolimus); OR ○ Immunosuppressive systemic agents (e.g., cyclosporine, azathioprine, methotrexate, mycophenolate mofetil, etc.); AND • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred injectable (Adbry, Dupixent) agent; AND • Nemluvio will be taken with topical corticosteroids and/or calcineurin inhibitors (e.g., pimecrolimus, tacrolimus); AND • Patient must meet the minimum age recommended by the package insert for this FDA approved indication. <p>Prurigo Nodularis:</p> <ul style="list-style-type: none"> • Diagnosis of prurigo nodularis; AND • At least 20 nodular lesions; AND • Other causes of pruritis have been ruled out; AND • Trial and failure, contraindication, or intolerance to one of the following: <ul style="list-style-type: none"> ○ Moderate to super potent topical corticosteroids [e.g., betamethasone dipropionate, (augmented), fluocinonide 0.1%, flurandrenolide, betamethasone dipropionate 0.05%, clobetasol propionate 0.025%, or dexamethasone 0.05%] for a minimum of 2 weeks; OR ○ Narrowband ultraviolet B (NBUVB) phototherapy or psoralen plus ultraviolet A (PUVA) phototherapy; AND • Trial and failure, contraindication, or intolerance to Dupixent; AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Patient must continue to meet initial approval criteria; AND • Patient must have disease improvement and/or stabilization based on an objective measure <p>Quantity Limit: 2 pens (60 mg) per 28 days</p>
Vtama AE, CC, QL	<p>Plaque Psoriasis</p> <ul style="list-style-type: none"> • Diagnosis of plaque psoriasis; AND • Prescribed by or in consultation with a dermatologist or other disease state specialist; AND • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to two preferred agents in the Dermatologics: Topical Antipsoriatics drug class. <p>Atopic Dermatitis</p> <ul style="list-style-type: none"> • Diagnosis of atopic dermatitis; AND • Prescribed by or in consultation with a dermatologist or other disease state specialist; AND • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to two preferred agents in the Immunomodulators, Atopic Dermatitis drug class.

Agent(s) Subject to Criteria	Criteria for Approval
	Age Limit: ≥ 2 years of age Quantity Limit: 2 grams per day
Zoryve 0.15% cream ^{AE, CC, QL}	Atopic Dermatitis <ul style="list-style-type: none"> • Diagnosis of mild to moderate atopic dermatitis; AND • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications), or intolerance to ONE preferred agent; AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. Age Limit: ≥ 6 years of age Quantity Limit: 2 grams per day
Zoryve 0.05% cream ^{AE, CC, QL}	Atopic Dermatitis <ul style="list-style-type: none"> • Diagnosis of mild to moderate atopic dermatitis; AND • Trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications), or intolerance to ONE preferred agent; AND • Patient must meet the minimum age recommended by the package insert for this FDA-approved indication. Age Limit: 2 to 5 years of age Quantity Limit: 2 grams per day

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Adbry autoinjector ^{AE, CC, QL}	Ebglyss ^{AE, CC, QL}
Adbry syringe ^{AE, CC, QL}	Nemludio ^{AE, CC, QL}
Dupixent pen ^{CC, QL}	Vtama ^{AE, CC, QL}
Dupixent syringe ^{CC, QL}	Zoryve 0.15% cream ^{AE, CC, QL}
Eucrisa ^{CC, QL}	Zoryve 0.05% cream ^{AE, CC, QL}
Opzelura cream ^{AE, CC, QL}	
pimecrolimus cream	
tacrolimus ointment	

BLOOD MODIFIERS: ANTIHYPERURICEMICS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
colchicine tablets ^{CC}	<ul style="list-style-type: none">• Diagnosis of one of the following conditions:<ul style="list-style-type: none">○ Familial Mediterranean Fever (FMF) (ICD-10 = E85.0); OR○ Pericarditis; OR○ Gout prophylaxis; OR○ Gout (acute attack) WITH trial and failure of, or contraindication/intolerance to, at least 1 of the following:<ul style="list-style-type: none">▪ Non-steroidal anti-inflammatory drug (NSAID); OR▪ Oral steroid.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 90 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Colcrys ^{CC}	<ul style="list-style-type: none">• NPD Criteria above when used for gout prophylaxis; OR• Diagnosis of one of the following conditions:<ul style="list-style-type: none">○ Familial Mediterranean Fever (FMF) (ICD-10 = E85.0); OR○ Pericarditis; OR○ Gout (acute attack) WITH trial and failure of, or contraindication/intolerance to, at least 1 of the following:<ul style="list-style-type: none">▪ Non-steroidal anti-inflammatory drug (NSAID); OR▪ Oral steroid.
colchicine capsules ^{CC} Gloperba ^{CC} Mitigare ^{CC}	<ul style="list-style-type: none">• NPD Criteria above; AND• Used for prophylaxis of gout flares.
Uloric ^{QL}	<ul style="list-style-type: none">• NPD Criteria above:<ul style="list-style-type: none">○ Therapeutic failure is defined as serum urate/uric acid level ≥ 6.0 mg/dL.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
allopurinol tablet	colchicine capsule ^{CC}
colchicine tablet ^{CC}	Colcrys tablet ^{CC}
febuxostat tablet ^{QL}	Gloperba solution ^{CC}
probenecid tablet	Mitigare capsule ^{CC}
probenecid/colchicine tablet	Uloric tablet ^{QL}
	Zyloprim tablet

BLOOD MODIFIERS: COLONY STIMULATING FACTORS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Fulphila ^{CC, QL} Fylmetra ^{CC, QL} Neupogen ^{CC, QL} Releuko ^{CC, QL}	Diagnosis of one of the following conditions: <ul style="list-style-type: none">• Myelosuppressive chemotherapy; OR• Induction or consolidation chemotherapy in acute myeloid/myelogenous leukemia; OR• Bone marrow transplantation; OR• Bone marrow transplant failure or engraftment delay; OR• Peripheral blood progenitor cell collection and therapy; OR• Severe chronic neutropenia (ANC \leq 500 mm³); OR• Hematopoietic Subsyndrome of Acute Radiation Syndrome

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 7 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent. Preferred with PA (PDP) Criteria must be met.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Neulasta ^{CC, QL} Neulasta Onpro ^{CC, QL} Nyvepria ^{CC, QL} Stimufend ^{CC, QL} Udenyca ^{CC, QL} Ziextenzo ^{CC, QL}	<ul style="list-style-type: none">• PDP Criteria above; AND<ul style="list-style-type: none">○ NPD Criteria above; OR○ Member is < 18 years old; OR○ Prescriber is a pediatric oncologist.
Rolvedon ^{AE, CC, QL} Ryzneuta ^{AE, CC, QL}	<ul style="list-style-type: none">• The medication is being used for chemotherapy-induced neutropenia prophylaxis, to decrease the incidence of febrile neutropenia; AND• Patient has a non-myeloid malignancy and is receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia; AND• Patient has had at least a 7-day trial and therapeutic failure, allergy, contraindication or intolerance of 2 preferred agents. <p>Age Limit: \geq 18 years Quantity Limit: 1 syringe per 14 days</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Fulphila ^{CC, QL}	Granix ^{QL}
Fylnetra ^{CC, QL}	Leukine ^{QL}
Neupogen ^{CC, QL}	Neulasta ^{CC, QL}
Releuko ^{CC, QL}	Neulasta Onpro ^{CC, QL}
	Nivestym ^{QL}
	Nyvepria ^{AE, CC, QL}
	Rolvedon ^{AE, CC, QL}
	Ryzneuta ^{AE, CC, QL}
	Stimufend ^{QL}
	Udenyca ^{CC, QL}
	Zarxio ^{QL}
	Ziextenzo ^{CC, QL}

BLOOD MODIFIERS: ERYTHROPOIESIS STIMULATING AGENTS

GUIDELINES FOR USE

Approval Duration: 3 months initial, 1 year renewal

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Aranesp ^{CC} Retacrit ^{CC} Epogen ^{CC}	<ul style="list-style-type: none">Member is not receiving hemodialysis*; ANDDocumentation (e.g., progress note, laboratory report) of hemoglobin (Hgb) < 10 g/dL in the past 90 days; ANDPrescribed for one of the following diagnosis:<ul style="list-style-type: none">Anemia associated with chronic renal failure; ORAnemia associated with kidney transplantation; ORTreatment of chemotherapy-induced anemia for non-myeloid malignancies; ORDrug-induced anemia (e.g., Retrovir®, Combivir® or ribavirin) (Retacrit and Epogen only); ORAutologous blood donations by patients scheduled to undergo nonvascular surgery. (Retacrit and Epogen Only) <p>Renewal Criteria:</p> <ul style="list-style-type: none">Documentation (e.g., progress note, laboratory report) of response to therapy.
Mircera ^{CC}	<ul style="list-style-type: none">Member is not receiving hemodialysis*; ANDPrescribed for anemia associated with chronic renal failure; ANDDocumentation (e.g., progress note, laboratory report) of hemoglobin (Hgb) < 10 g/dL in the past 90 days. <p>Renewal Criteria:</p> <ul style="list-style-type: none">Documentation (e.g., progress note, laboratory report) of response to therapy.

*Providers should bill Medicare B if member is receiving hemodialysis.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Jesduvroq ^{CC}	<p>Approval Duration: 6 months</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of chronic kidney disease (N18.9); AND • Pretreatment hemoglobin level $\leq 11\text{g/dl}$; AND • Patient has been receiving dialysis for at least 4 months; AND • Patient is not receiving treatment with any other erythropoiesis stimulating agents. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Documentation (e.g., progress note, laboratory report) demonstrating a positive response to therapy. <p>Quantity Limit:</p> <ul style="list-style-type: none"> • 1 mg, 2 mg, & 4 mg: one daily • 6 mg: two daily • 8 mg: three daily
Reblozyl ^{CC, AE}	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Prescribed by, or in consultation with, a hematology or oncology specialist; AND • Diagnosis of beta thalassemia requiring regular red blood cell (RBC) transfusions; OR • Diagnosis of anemia that is associated with low-to-moderate-risk myelodysplastic syndromes; AND <ul style="list-style-type: none"> ◦ Member has required 2 or more RBC units over an 8-week period; AND ◦ Serum erythropoietin (EPO) $< 500\text{ mU/mL}$; OR • Diagnosis of anemia that is associated with low-to-moderate-risk myelodysplastic syndromes with ring sideroblasts or myelodysplastic/myeloproliferative neoplasm with ring sideroblasts and thrombocytosis; AND • Patient has required 2 or more RBC units over an 8-week period; AND <ul style="list-style-type: none"> ◦ Failure of an erythropoiesis stimulating agent (e.g., epoetin alfa); OR ◦ Serum erythropoietin (EPO) $> 500\text{ mU/mL}$. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Attestation or documentation (e.g., progress note) of a reduction in transfusion burden or other clinical benefit. <p>Age Limit: ≥ 18 years</p>
Vafseo ^{CC, QL}	<p>Approval Duration: 6 months</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> • Diagnosis of chronic kidney disease (N18.9); AND • Pretreatment hemoglobin level $\leq 11\text{g/dl}$; AND • Patient has been receiving dialysis for at least 3 months; AND • Patient does not have uncontrolled hypertension; AND • Patient is not receiving treatment with any other erythropoiesis stimulating agents; AND • Patient meets the minimum age recommended by the package insert. <p>Renewal Criteria:</p>

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Documentation (e.g., progress notes, laboratory report) of a positive response to therapy. <p>Quantity Limit:</p> <ul style="list-style-type: none"> 150 mg four tablets per day 300 mg two tablets per day

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Aranesp ^{CC}	Jesduvroq ^{CC, QL}
Epogen ^{CC}	Procrit
Mircera ^{CC}	Reblozyl ^{CC, AE}
Retacrit ^{CC}	Vafseo ^{CC, QL}

BLOOD MODIFIERS: PHOSPHATE BINDERS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 7 day** trial and failure within the past 90 days, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Xphozah ^{CC, AE, QL}	<ul style="list-style-type: none">Diagnosis of chronic kidney disease; ANDDiagnosis of elevated serum phosphorous; ANDPatient is on dialysis; ANDPatient has had a trial and failure, contraindication to, intolerance, or inadequate response to at least 2 preferred phosphate binders. <p>Age Limit: ≥ 18 years of age Quantity Limit: 2 tablets daily</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
calcium acetate capsule, tablet	Auryxia
Phoslyra solution	ferric citrate tablet
sevelamer carbonate powder packet, tablet	Fosrenol chewable tablet, powder packet
sevelamer tablet	lanthanum carbonate chewable tablet
	Renagel
	Renvela powder packet, tablet
	Velphoro
	Xphozah ^{CC, AE, QL}

BLOOD MODIFIERS: SICKLE CELL ANEMIA TREATMENTS

GUIDELINES FOR USE

Approval Duration: 1 year, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Endari ^{CC, AE, QL}	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none">• Diagnosis of sickle cell disease; AND• Prescribed by or consultation with a hematologist or a provider that specializes in sickle cell disease; AND• Documentation that the member has had at least two vaso-occlusive crises within the past 12 months; AND• Patient has tried hydroxyurea for at least 3 months, unless contraindicated or intolerant. <p>Renewal Criteria:</p> <ul style="list-style-type: none">• Patient must have disease improvement (decrease in the number of sickle cell crises) <p>Age Limit: ≥ 5 years old Quantity Limit: 6 packets (30 gm) per day</p>

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires ≥ **3-month** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ **2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
L-glutamine (generic for Endari) ^{CC, AE, QL}	Initial Approval Criteria: <ul style="list-style-type: none">• Diagnosis of sickle cell disease; AND• Prescribed by or consultation with a hematologist or a provider that specializes in sickle cell disease; AND• Documentation that the member has had at least two vaso-occlusive crises within the past 12 months; AND• Patient has tried hydroxyurea for at least 3 months, unless contraindicated or intolerant. Renewal Criteria: <ul style="list-style-type: none">• Patient must have disease improvement (decrease in the number of sickle cell crises) Age Limit: ≥ 5 years old Quantity Limit: 6 packets (30 gm) per day

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Droxia	L-glutamine (generic for Endari) ^{CC, AE, QL}
Endari ^{CC, AE, QL}	Xromi solution ^{AE}
Siklos	

BLOOD MODIFIERS: THROMBOPOIESIS STIMULATING PROTEINS

GUIDELINES FOR USE

Approval Duration: 6 months, unless otherwise specified

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Promacta tablets ^{CC, QL}	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none">• Prescribed by, or in consultation with, a hematologist or liver disease specialist; AND• Diagnosis of one of the following conditions:<ul style="list-style-type: none">○ Chronic immune (idiopathic) thrombocytopenic purpura (ITP); OR○ Treatment of thrombocytopenia in patients with chronic hepatitis C (to allow the initiation & maintenance of interferon-based therapy); OR○ Treatment of severe aplastic anemia in patients who have had an insufficient response to immunosuppressive therapy. <p>Renewal Criteria:</p> <ul style="list-style-type: none">• Documentation (e.g., progress note, laboratory report) of response to therapy.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure within the past 90 days, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Alvaiz ^{CC, AE, QL}	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none">• Prescribed by, or in consultation with, a hematologist or liver disease specialist; AND• Patient has one of the following indications:<ul style="list-style-type: none">○ Diagnosis of persistent or chronic immune thrombocytopenia (ITP) with an insufficient response to corticosteroids, immunoglobulins, or splenectomy; OR

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ Used for the treatment of thrombocytopenia in patients with chronic hepatitis C (to allow the initiation and maintenance of interferon-based therapy); OR ○ Diagnosis of severe aplastic anemia with an insufficient response to immunosuppressive therapy; AND ● Patient meets the minimum age recommended by the package insert for respective indications. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> ● Documentation (e.g., progress note, laboratory report) of response to therapy. <p>Age Limit: 6 years or older</p> <p>Quantity Limit: 9 mg: 1 per day 18 mg: 1 per day 36 mg: 3 per day 54 mg: 2 per day</p>
Doptelet ^{CC, AE, QL}	<p>Approval Duration: Date of service (chronic liver disease); 6 months (ITP)</p> <p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> ● Diagnosis of chronic liver disease; AND <ul style="list-style-type: none"> ○ Documentation of platelet count $< 50 \times 10^9/L$ within the past 14 days; AND ○ Prescribed per FDA-approved labeling (10 tablets per 5 days for platelets $\geq 40 \times 10^9/L$ or 15 tablets per 5 days for platelets $< 40 \times 10^9/L$); AND ○ Confirmation (e.g., attestation or progress note) of a scheduled invasive procedure occurring 5 to 8 days following the last dose of avatrombopag; AND ○ Patient must meet the minimum age recommended by the package insert for the FDA-approved indication; OR ● Diagnosis of persistent or chronic immune (idiopathic) thrombocytopenic purpura (ITP); AND <ul style="list-style-type: none"> ○ Prescribed by, or in consultation with a hematologist or immunologist, or other specialist in the treatment of ITP; AND ○ Documentation (e.g., progress note, laboratory report) of platelet count within the past 30 days; AND ○ Trial and failure (i.e., not achieved a platelet count $\geq 50 \times 10^9/L$) of at least one other therapy for chronic ITP, such as corticosteroids, IV immune globulin, RhO(D) immune globulin, thrombopoietin receptor antagonists, etc.; AND ○ Patient must meet the minimum age recommended by the package insert for the FDA-approved indication. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> ● Documentation (e.g., progress note, laboratory report) of response to therapy. <p>Quantity Limit: 2 per day (except where 15 tablet per 5-day course is indicated)</p>
Doptelet Sprinkle ^{AE, CC, QL}	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> ● Diagnosis of persistent or chronic immune thrombocytopenic purpura (ITP); AND ● Prescribed by, or in consultation with a hematologist or immunologist, or other specialist in the treatment of ITP; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Documentation (e.g., progress note, laboratory report) of platelet count within the past 30 days; AND Trial and failure (i.e., not achieved a platelet count $\geq 50 \times 10^9/L$) of at least one other therapy for ITP, such as corticosteroids, IV immune globulin, RhO(D) immune globulin, thrombopoietin receptor antagonists, etc. <p>Renewal Criteria</p> <ul style="list-style-type: none"> Documentation (e.g., progress note, laboratory report) of response to therapy. <p>Age limit: 1 to less than 6 years of age Quantity Limit: 2 sprinkle capsules per day (except where 15 tablet per 5-day course is indicated)</p>
eltrombopag olamine powder pack CC, QL eltrombopag olamine tablets CC, QL	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none"> Prescribed by, or in consultation with, a hematologist or liver disease specialist; AND Diagnosis of one of the following conditions: <ul style="list-style-type: none"> Chronic immune (idiopathic) thrombocytopenic purpura (ITP); OR Treatment of thrombocytopenia in patients with chronic hepatitis C (to allow the initiation & maintenance of interferon-based therapy); OR Treatment of severe aplastic anemia in patients who have had an insufficient response to immunosuppressive therapy. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Documentation (e.g., progress note, laboratory report) of response to therapy. <p>Quantity Limit: 12.5 mg and 25 mg powder pack: 3 per day 12.5 mg and 25 mg tablet: 3 per day 50 mg and 75 mg tablet: 2 per day</p>
Mulpleta CC, AE, QL	<p>Approval Duration: Date of service</p> <ul style="list-style-type: none"> Diagnosis of chronic liver disease; AND Documentation of platelet count $< 50 \times 10^9/L$ within the past 14 days; AND NOT have severe hepatic impairment (Child-Pugh class C), absence of hepatopetal blood flow, a prothrombotic condition other than CLD or a history of splenectomy, partial splenic embolization, or thrombosis; AND Confirmation (e.g., attestation or progress note) of a scheduled invasive procedure occurring 2 to 8 days following the last dose of lusutrombopag. <p>Age Limit: ≥ 18 years Quantity Limit: 7 tablets per fill; no renewals</p>
Tavalisse CC, AE, QL	<ul style="list-style-type: none"> Diagnosis of chronic immune (idiopathic) thrombocytopenic purpura (ITP); AND Prescribed by, or in consultation with, a hematologist or liver disease specialist; AND Documentation (e.g., progress note, laboratory report) of platelet count within the past 30 days; AND

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> Trial and failure (i.e., not achieved a platelet count $\geq 50 \times 10^9/L$) of at least one other therapy for chronic ITP, such as corticosteroids, IV immune globulin, RhO(D) immune globulin, thrombopoietin receptor antagonists, etc. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> Documentation (e.g., progress note, laboratory report) of response to therapy. <p>Age Limit: ≥ 18 years Quantity Limit: 2 per day</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Promacta tablet ^{CC}	Alvaiz ^{CC, AE, QL}
	Doptelet sprinkles, tablet ^{CC, AE, QL}
	eltrombopag olamine powder pack ^{CC, QL}
	eltrombopag olamine tablet ^{CC, QL}
	Mulpleta ^{CC, AE, QL}
	Nplate
	Promacta powder packet ^{QL}
	Tavalisse ^{CC, AE, QL}

OPHTHALMIC ANTIBIOTICS AND ANTIVIRALS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 3 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent within the same sub-class.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

OPHTHALMIC ANTIVIRALS

Preferred Agents	Non-Preferred Agents
trifluridine	Zirgan

OPHTHALMIC QUINOLONES

Preferred Agents	Non-Preferred Agents
ciprofloxacin	Besivance
moxifloxacin (generic Vigamox)	Ciloxan
ofloxacin	gatifloxacin
	levofloxacin
	moxifloxacin (generic Moxeza)
	Ocuflox
	Vigamox
	Zymaxid

OPHTHALMIC ANTIBIOTICS, NON-QUINOLONES

Preferred Agents	Non-Preferred Agents
bacitracin ointment	AzaSite
bacitracin/polymyxin B ointment	Natacyn
erythromycin 0.5% ointment	neomycin/polymyxin B/bacitracin ointment
gentamicin sulfate drops	neomycin/polymyxin B/gramicidin drops
Polycin	Neo/Polycin ointment
polymyxin B/trimethoprim drops	Polytrim

Preferred Agents	Non-Preferred Agents
sulfacetamide drops tobramycin drops	sulfacetamide ointment Tobrex

OPHTHALMIC ANTIBIOTIC-STEROID COMBINATIONS

Preferred Agents	Non-Preferred Agents
dexamethasone/neomycin sulfate/polymyxin B suspension	Blephamide S.O.P ointment
hydrocortisone/bacitracin zinc/neomycin sulfate/polymyxin B ointment, suspension	hydrocortisone/neomycin sulfate/polymyxin B suspension
Neo-Polycin hydrocortisone ointment	Maxitrol ointment, suspension
Tobradex ointment, suspension	prednisolone sodium phosphate/sulfacetamide sodium
tobramycin/dexamethasone suspension	Pred-G ointment
	Tobradex ST
	Zylet

OPHTHALMICS FOR ALLERGIC CONJUNCTIVITIS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

OPHTHALMIC ANTIHISTAMINES

Preferred Agents	Non-Preferred Agents
azelastine	bepotastine besilate
olopatadine 0.1% (generic Patanol)	Bepreve
olopatadine 0.2% (generic Pataday)	epinastine
	Zerviate

OPHTHALMIC MAST CELL STABILIZERS

Preferred Agents	Non-Preferred Agents
cromolyn sodium	Alocril
	Alomide

OPHTHALMICS: GLAUCOMA AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

OPHTHALMIC PROSTAGLANDIN AGONISTS

Preferred Agents	Non-Preferred Agents
latanoprost ^{QL}	bimatoprost ^{QL}
	Iyuzeh ^{QL}
	Lumigan ^{QL}
	Tafluprost ^{QL}
	Travatan Z
	Travoprost
	Vyzulta ^{AE, QL}
	Xalatan ^{QL}
	Xelpros
	Zioptan ^{QL}

OPHTHALMIC BETA BLOCKERS

Preferred Agents	Non-Preferred Agents
levobunolol	betaxolol
timolol maleate drops (except preservative free)	Betimol
	Betoptic S
	Carteolol
	Istalol
	timolol (generic Betimol)
	timolol maleate once daily (generic Istalol)

Preferred Agents	Non-Preferred Agents
	timolol PF (preservative-free)
	timolol maleate gel-solution

OPHTHALMIC CARBONIC ANHYDRASE INHIBITORS

Preferred Agents	Non-Preferred Agents
dorzolamide	Azopt
	brinzolamide

OPHTHALMIC COMBINATIONS FOR GLAUCOMA

Preferred Agents	Non-Preferred Agents
Combigan	brimonidine-timolol 0.2%-0.5%
dorzolamide/timolol (except preservative-free)	Cosopt
Simbrinza	Cosopt PF
	dorzolamide/timolol PF (preservative-free)

OPHTHALMIC SYMPATHOMIMETICS

Preferred Agents	Non-Preferred Agents
Alphagan P 0.15%	Alphagan P 0.1%
brimonidine 0.2% drops	apraclonidine
	brimonidine 0.1% drops, 0.15% drops
	lopidine

OPHTHALMICS, GLAUCOMA AGENTS (OTHER)

Preferred Agents	Non-Preferred Agents
Rhopressa	phospholine iodide
Rocklatan	pilocarpine
	Vuity

OPHTHALMICS: NON-STEROIDAL ANTI-INFLAMMATORY DRUGS (NSAIDS)

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
diclofenac sodium drops	Acular
flurbiprofen sodium drops	Acular LS
ketorolac tromethamine drops	Acuvail
	bromfenac sodium drops
	BromSite
	Ilevro
	Nevanac
	Prolensa

OPHTHALMICS: ANTI-INFLAMMATORY STEROIDS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
dexamethasone sodium phosphate drops	Alrex
Durezol	difluprednate
fluorometholone suspension	Eysuvis
Lotemax gel, ointment, suspension	Flarex
prednisolone acetate suspension	FML suspension, FML Forte suspension
prednisolone sodium phosphate drops	Inveltys
	Lotemax SM gel
	loteprednol etabonate gel, suspension
	Maxidex
	Pred Forte
	Pred Mild

OPHTHALMICS: IMMUNOMODULATORS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Restasis (Blister Pack) ^{CC} Xiidra ^{CC, AE, QL}	<ul style="list-style-type: none">Prescribed following corneal transplant; ORTrial and failure of ≥ 1 over-the-counter ophthalmic lubricant (e.g., polyvinyl alcohol).

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires ≥ 30 day trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Miebo ^{CC, QL}	<ul style="list-style-type: none">Trial and failure of ≥ 1 over-the-counter ophthalmic lubricant (e.g., polyvinyl alcohol); ANDAt least a 1-month trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents. <p>Quantity Limit: 0.4 mL (8 drops) per day</p>
Restasis Multidose ^{CC}	<ul style="list-style-type: none">Prescribed following corneal transplant; ORTrial and failure of ≥ 1 over-the-counter ophthalmic lubricant (e.g., polyvinyl alcohol).
Tyrvaya ^{CC, AE, QL}	<p>Initial Approval Criteria:</p> <ul style="list-style-type: none">Patient has diagnosis of dry eye disease (DED); ANDPrescribed by or in consultation with an ophthalmologist or optometrist; ANDPatient has had a trial and failure of preservative-free, nonprescription lubricating eye drops (e.g., artificial tears); ANDPatient has had ≥ 1 month trial and therapeutic failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents; ANDPrescriber has documented at least 1 of the following signs of DED:

Agent(s) Subject to Criteria	Criteria for Approval
	<ul style="list-style-type: none"> ○ Corneal fluorescein staining (CFS) score of ≥ 2 points in any field on a 0 to 4 point scale; OR ○ Schirmer tear test (STT) of 1 to 10 mm in 5 minutes. <p>Renewal Criteria:</p> <ul style="list-style-type: none"> • Patient continues to meet the above criteria; AND • Patient has improvement in signs of DED, as measured by at least 1 of the following: <ul style="list-style-type: none"> ○ Decrease in corneal fluorescein staining score; OR ○ Increase in number of mm per 5 minutes using Schirmer tear test. <p>Age Limit: ≥ 18 years old Quantity Limit: 1 carton (2 bottles)/ 30 days</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Restasis (Blister Pack) ^{CC}	Cequa
Xiidra ^{AE, CC, QL}	cyclosporine 0.05%
	Miebo ^{CC, QL}
	Restasis Multidose ^{CC}
	Tyrvaya ^{AE, CC, QL}
	Verkazia

OPHTHALMIC: MYDRIATIC & CYCLOPLEGICS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
atropine sulfate drops, ointment	Cyclogyl drops
atropine sulfate/PF dropperette	Cyclomydril drops
Cyclopentolate drops	Mydracil drops
phenylephrine drops	
Tropicamide drops	

OTICS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents (antibiotics) or 1 preferred agent (anesthetics and anti-inflammatories).

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable

CURRENT PDL STATUS

OTIC ANTIBIOTICS

Preferred Agents	Non-Preferred Agents
CiproDex Otic	ciprofloxacin 0.2% drops
ciprofloxacin/dexamethasone suspension	Cipro HC Otic
hydrocortisone/neomycin/polymyxin B suspension, solution	ciprofloxacin/fluocinolone
ofloxacin 0.3% solution	Cortisporin-TC suspension
	Otovel

OTIC ANESTHETICS AND ANTI-INFLAMMATORIES

Preferred Agents	Non-Preferred Agents
acetic acid	DermOtic
fluocinolone acetonide 0.01% oil	Flac Otic Oil
	hydrocortisone/acetic acid drops

RENAL AND GENITOURINARY: ALPHA BLOCKERS FOR BPH & 5-ALPHA REDUCTASE INHIBITORS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
finasteride 5 mg ^{CC}	<ul style="list-style-type: none">Diagnosis of benign prostatic hyperplasia (ICD-10 Disease Group N40).

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents in any sub-class, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

ALPHA BLOCKERS FOR BPH

Preferred Agents	Non-Preferred Agents
alfuzosin ER	Cardura
doxazosin	Cardura XL
tamsulosin	Flomax
terazosin	Rapaflo
	silodosin
	Tezruly ^{QL}

5-ALPHA REDUCTASE (5AR) INHIBITORS

Preferred Agents	Non-Preferred Agents
dutasteride	Avodart
finasteride 5 mg tablet ^{CC}	dutasteride/tamsulosin
	Entadfi
	finasteride-tadalafil ^{QL}
	Jalyn
	Proscar

BLADDER RELAXANTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Gemtesa ^{CC, AE, QL}	Initial Approval Criteria: <ul style="list-style-type: none">Patient is ≥ 18 years of age; ANDPatient has a diagnosis of overactive bladder (OAB) with symptoms of urge urinary incontinence, urgency, and urinary frequency; ANDPatient must not have hypersensitivity to vibegron or any component of the product; ANDPatient must have an adequate trial and failure of behavioral therapy (bladder training, bladder control strategies, pelvic floor muscle training, and fluid management); ANDPatient has tried and failed at least one month, or has an intolerance, or contraindication to at least two preferred medications; ANDPatient has tried and failed at least one month of treatment with mirabegron. Renewal Criteria: <ul style="list-style-type: none">Patient has not experienced urinary retention; ANDPatient has experienced disease response as indicated by a reduction in the daily number of micturitions and the average daily number of urge urinary incontinence (UUI) episodes.
Oxytrol ^{CC, QL}	<ul style="list-style-type: none">Patient is unable to swallow or tolerate oral medications; ORNPD criteria

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
oxybutynin (5mg tablet ^{QL} , solution ^{QL} , syrup ^{QL})	darifenacin ER ^{QL}
oxybutynin ER ^{QL}	Detrol ^{QL}
solifenacin ^{QL}	Detrol LA ^{QL}
tolterodine ^{QL}	fesoterodine ER ^{QL}
tolterodine ER ^{QL}	flavoxate ^{QL}
	Gemtesa ^{CC, AE, QL}
	mirabegron ER ^{QL}
	Myrbetriq ^{QL}
	oxybutynin 2.5mg tablet ^{QL}
	Oxytrol ^{CC, QL}
	Toviaz ER ^{QL}
	trospium ^{QL}
	trospium ER ^{QL}
	Vesicare ^{QL}
	Vesicare LS ^{QL}

DERMATOLOGICS: TOPICAL ANTIBIOTIC AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent, unless otherwise specified.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Xepi ^{CC, QL}	<ul style="list-style-type: none">Diagnosis of impetigo; ANDTrial and failure with a preferred agent (e.g., mupirocin ointment); ANDNot have an affected body surface area (BSA) exceeding 100 cm² or 2% of total BSA, whichever is greater; ANDWill not be used for more than 5 days <p>Quantity Limit: Up to 30 grams per fill</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
gentamicin cream ^{QL} , ointment	Centany ointment ^{QL} , Centany AT kit
mupirocin ointment ^{QL}	mupirocin cream ^{QL}
	Xepi ^{CC, QL}

DERMATOLOGICS: TOPICAL ANTIPARASITICS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Natroba	Crotan
permethrin 5% cream	Elimite
	Eurax
	Lindane
	malathion lotion
	Ovide
	spinosad

DERMATOLOGICS: ORAL ANTIPSORIATICS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
acitretin	methoxsalen

DERMATOLOGICS: ORAL ACNE AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation. [Documentation required]

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
Amnesteem	Absorica
Claravis	Absorica LD
Zenatane	isotretinoin capsule

DERMATOLOGICS: TOPICAL ACNE AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 30 day** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of **≥ 4** preferred or over-the-counter (OTC) agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an *inactive* ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
adapalene/benzoyl peroxide 0.3-2.5% (Mayne Pharma)	adapalene cream, gel, gel pump
clindamycin gel, medicated swab (pledget), solution	adapalene/benzoyl peroxide gel
clindamycin/benzoyl peroxide (generic BenzaClin or Duac; excluding pumps)	Avar, Avar E, Avar E LS, Avar LS
erythromycin solution	BP 10-1 cleanser
erythromycin/benzoyl peroxide	BP Cleansing Wash
	Cleocin-T
	Clindacin ETZ kit, medicated swab
	Clindacin foam
	Clindacin P medicated swab
	Clindacin PAC kit
	clindamycin foam, lotion
	clindamycin phosphate EQ 1% gel (generic Clindagel)
	clindamycin/benzoyl peroxide gel pump (generic Acanya)
	clindamycin/benzoyl peroxide gel pump
	clindamycin/tretinoin gel
	dapsone gel, gel pump
	Differin cream, gel pump, lotion
	Epiduo Forte
	Ery medicated swab
	Erygel

Preferred Agents	Non-Preferred Agents
	erythromycin gel
	Evoclin
	Fabior
	Neuac gel
	Neuac Kit
	Ovace Plus shampoo, wash, wash cleanser gel
	Rosula
	sodium sulfacetamide cleanser, cleanser gel, shampoo, suspension
	sodium sulfacetamide/sulfur cleanser, cream, medicated pad, suspension
	SSS 10-5 cream, foam
	Sumadan cleanser, kit
	Sumadan XLT cleanser cream
	Sumaxin, Sumaxin CP, Sumaxin TS
	tazarotene cream, foam, gel
	tretinoin cream, gel, microsphere gel, microsphere gel pump
	Twynéo
	Winlevi ^{AE}
	Zma Clear suspension

DERMATOLOGICS: TOPICAL ROSACEA AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Rhofade ^{CC, AE, QL}	<ul style="list-style-type: none">Diagnosis of rosacea or facial erythema; ANDTrial and failure of topical metronidazole; ANDTrial and failure of an oral antibiotic (e.g., doxycycline). <p>Quantity Limit: 60 grams per 30 days Age Limit: ≥ 18 years</p>

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
azelaic acid gel	brimonidine tartrate 0.33% gel pump
Finacea gel	Epsolay
metronidazole cream, gel, gel pump	Finacea foam
	ivermectin 1% cream
	MetroCream
	MetroGel
	metronidazole lotion
	Mirvaso
	Rhofade ^{CC, AE, QL}
	Rosadan
	Soolantra

DERMATOLOGICS: TOPICAL ANTIFUNGAL AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥1 week** trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Kerydin ^{CC}	<ul style="list-style-type: none">Diagnosis of toenail onychomycosisTrial and failure of ciclopirox 8% nail solution or allergy to ciclopirox

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
ciclopirox cream, solution	Ciclodan cream, kit, solution
clotrimazole cream, solution	ciclopirox gel, kit, shampoo, suspension
clotrimazole/betamethasone cream	clotrimazole/betamethasone lotion
econazole cream	econazole foam
ketoconazole cream ^{QL} , shampoo	Ertaczo
Nyamyc	Extina
nystatin cream, ointment, powder ^{QL}	ketoconazole foam
nystatin/triamcinolone cream, ointment	Ketodan
Nystop	Kerydin ^{CC}
tavaborole	Loprox cream, cream kit, suspension, suspension kit
	miconazole/zinc oxide/petrolatum
	naftifine cream, gel
	Naftin
	oxiconazole ^{QL}
	Oxistat ^{QL}
	Vusion

DERMATOLOGICS: TOPICAL ANTIVIRAL AGENTS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 1 preferred agent.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- a. trial and failure (e.g., allergy or intolerance to an inactive ingredient) with ≥ 2 manufacturers (if available and covered) of the corresponding generic; **AND**
- b. NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
acyclovir cream, ointment	Denavir penciclovir cream

DERMATOLOGICS: TOPICAL ANTIPSORIATICS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 1 week** trial and failure within the past 90 days, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Agent(s) Subject to Criteria	Criteria for Approval
Zoryve 0.3% cream ^{AE, CC, QL}	<ul style="list-style-type: none">Diagnosis of plaque psoriasis; ANDNPD Criteria; ANDPatient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Age Limit: ≥ 6 years Quantity Limit: 2 grams per day</p>
Zoryve 0.3% foam ^{AE, CC, QL}	<p>Plaque Psoriasis</p> <ul style="list-style-type: none">Diagnosis of plaque psoriasis; ANDTrial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to ONE preferred agent; ANDPatient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Seborrheic dermatitis</p> <ul style="list-style-type: none">Diagnosis of seborrheic dermatitis; ANDAt least a 4-week trial and failure, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance to ONE of the following agents for the treatment of seborrheic dermatitis:<ul style="list-style-type: none">Topical steroids (e.g., hydrocortisone butyrate), ORTopical antifungal (e.g., ketoconazole, ciclopirox); ANDPatient must meet the minimum age recommended by the package insert for this FDA-approved indication. <p>Age Limit:</p>

Agent(s) Subject to Criteria	Criteria for Approval
	≥12 years for plaque psoriasis ≥9 years for seborrheic dermatitis Quantity Limit: 2 grams per day

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
calcipotriene cream, ointment, solution	calcipotriene foam
calcipotriene/betamethasone suspension	calcipotriene/betamethasone ointment
salicylic acid gel, liquid film	calcitriol ointment
urea cream ^{QL}	Dermacure cream ^{QL}
	Enstilar ^{MD, AE}
	salicylic acid foam, ointment
	Sorilux
	Taclonex suspension
	Uramaxin foam
	urea foam
	Vertical ointment
	Zoryve 0.3% cream, 0.3% foam ^{AE, CC, QL}

DERMATOLOGICS: TOPICAL STEROIDS

GUIDELINES FOR USE

Approval Duration: 1 year

1. PREFERRED WITH PA (PDP) CRITERIA

Not applicable. All preferred agents are preferred without PA.

2. NON-PREFERRED (NPD) CRITERIA

Approval of non-preferred agents requires **≥ 1 week** trial and failure **within the past 90 days**, allergy, contraindication (including potential drug-drug interactions with other medications) or intolerance of 2 preferred agents.

3. BRAND MEDICALLY NECESSARY CRITERIA

Approval of NON-PREFERRED branded agents requires:

- trial and failure (e.g., allergy or intolerance to an inactive ingredient) with **≥ 2** manufacturers (if available and covered) of the corresponding generic; **AND**
- NPD Criteria above.

4. GENERIC MEDICALLY NECESSARY CRITERIA

Known or suspected allergy, intolerance, or contraindication to an inactive ingredient in the preferred brand formulation.

5. DRUG-SPECIFIC CLINICAL CRITERIA

Not applicable.

CURRENT PDL STATUS

Preferred Agents	Non-Preferred Agents
alclometasone dipropionate	amcinonide cream ^{QL}
betamethasone dipropionate cream, lotion	Ana-Lex ^{QL}
betamethasone dipropionate (augmented) cream	Apexicon E
betamethasone valerate cream, ointment	Beser
clobetasol propionate 0.05% cream, ointment, shampoo, solution	betamethasone dipropionate augmented ointment, lotion, gel
Clodan shampoo	betamethasone dipropionate ointment
Derma-Smoothe/FS	betamethasone valerate foam, lotion
desonide cream, ointment	Capex Shampoo
fluocinonide ointment, solution	clobetasol emollient, emulsion
fluticasone propionate cream, ointment	clobetasol propionate 0.025% cream
halobetasol propionate cream, ointment	clobetasol propionate foam, gel, lotion, spray
hydrocortisone cream, lotion, ointment	Clobex spray
mometasone furoate cream, ointment, solution	clocortolone cream
Procto-Med HC	Clodan shampoo kit
Proctosol-HC	desonide lotion
Proctozone-HC	desoximetasone cream, gel, ointment, spray
triamcinolone acetonide cream, lotion, ointment	diflorasone diacetate cream, ointment
	Diprolene AF
	fluocinolone acetonide cream, oil, ointment, solution
	fluocinonide emollient cream
	fluocinonide-E cream
	halcinonide cream, solution
	halobetasol propionate foam

Preferred Agents	Non-Preferred Agents
	Halog cream, ointment, solution
	hydrocortisone butyrate cream, lotion, ointment, solution
	hydrocortisone butyrate/emollient cream
	hydrocortisone solution
	hydrocortisone valerate cream, ointment
	Impeklo
	Kenalog
	Lexette
	Luxiq
	Olux
	Pandel
	prednicarbate ointment
	Synalar TS
	Temovate
	Topicort cream, gel, ointment, spray
	Tovet emollient foam, kit
	triamcinolone acetonide spray
	Ultravate



MedImpact Healthcare Systems, Inc.

10181 Scripps Gateway Ct. San Diego, CA 92131

Phone: 800.788.2949

[MedImpact.com](https://www.MedImpact.com)

Copyright © 2025 MedImpact Healthcare Systems, Inc. All rights reserved.

This document is confidential and proprietary to MedImpact and contains material MedImpact may consider Trade Secrets. This document is intended for specified use by Business Partners of MedImpact under permission by MedImpact and may not otherwise be used, reproduced, transmitted, published, or disclosed to others without prior written authorization. MedImpact maintains the sole and exclusive ownership, right, title, and interest in and to this document.

